THE ONTARIO GREAT LAKES COASTAL WETLAND ATLAS:

A Summary of Information (1983-1997)

Environment Canada Environnement Canada

Ontario

ISBN Number: 0-662-33822-7
Catalogue Number: CW66-221/1997E

© March, 2003
Printed in Ontario, Canada

For more information contact:

Brian Potter
Ontario Ministry of Natural Resources
Conservation and Planning Section, Lands and Waters Branch
300 Water St., PO Box 7000, Peterborough, ON K9J 8M5
Tel: (705) 755-1917; E-mail: brian.potter@mnr.gov.on.ca

Maggie Galloway
Canadian Wildlife Service, Environment Canada
4905 Dufferin St., Downsview, ON M3H 5T4
Tel: (416) 739-4950; E-mail: maggie.galloway@ec.gc.ca

Please note: The Atlas data is subject to Ontario Ministry of Natural Resources data-sharing polices and will not generally be made available electronically other than to Geo-Spatial Data Exchange partners who may contact OMNR's Information Access Group at 1-800-667-1940.

For additional copies of the Atlas contact:

Liz Sauer, Canadian Wildlife Service, Ontario Region
4905 Dufferin St., Downsview, ON M3H 5T4
E-mail: Wildlife.Ontario@ec.gc.ca; Tel: (416) 739-5830

Natural Resources Information Centre – Peterborough
Robinson Pl, 1st Flr N, 300 Water St.
PO Box 7000, Peterborough, ON K9J 8M5
Toll free: 1-800-667-1940

Natural Resources Information Centre – Toronto
Macdonald Block, Rm M1-73
900 Bay St., Toronto, ON M7A 2C1
Toll free: 1-800-667-1940

This report is by:

Environment Canada

Canadian Wildlife Service

The Canadian Wildlife Service, part of Environment Canada, manages wildlife matters that are the responsibility of the federal government. These include protection and management of migratory birds, nationally significant habitat and endangered species, as well as work on other wildlife issues of national and international importance. In addition, the Canadian Wildlife Service conducts research in many fields of wildlife biology and provides incentive programs for land stewardship and donation. For more information, visit: www.on.ec.gc.ca/wildlife.

Ontario Ministry of Natural Resources

Conservation and Planning Section, Lands and Waters Branch

The Ontario Ministry of Natural Resources (OMNR) manages fish and wildlife, land, water, forest, and provincial parks matters that are the responsibility of the Ontario government. It also conducts related scientific research, and shares information and technology. OMNR is also responsible for developing and conducting evaluations under the Ontario Wetland Evaluation System. For more information, visit: www.mnr.gov.on.ca/MNR/index.html.

Natural Heritage Information Centre

The Natural Heritage Information Centre is that part of the Ontario Ministry of Natural Resources that compiles, maintains and provides information on rare, threatened and endangered species and spaces in Ontario. This information is stored in a central repository containing a computerized database, map files and an information library, which are accessible for conservation applications, land use planning, park management, etc. The NHIC web site makes this information available through the world wide web. For more information, visit: www.mnr.gov.on.ca/MNR/nhic/nhic.cfm.

This report may be cited as:
Environment Canada and Ontario Ministry of Natural Resources. 2003. *The Ontario Great Lakes Coastal Wetland Atlas: A Summary of Information (1983-1997).*

Cover art: King Rail: Marlene Ross; Great Lakes: INTUVISION Multimedia

Canada

THE ONTARIO GREAT LAKES COASTAL WETLAND ATLAS:

A Summary of Information (1983-1997)

Environment Canada

- Canadian Wildlife Service

Ontario Ministry of Natural Resources

- Conservation and Planning Section - Lands and Waters Branch; and
- Natural Heritage Information Centre

March 2003

105-00068/

AUTHORS

(*) Formerly of Ontario Ministry of Natural Resources

Helen Ball
Ontario Ministry of Natural Resources, Peterborough

Jarmo Jalava*
Ontario Ministry of Natural Resources, Peterborough

Tayarna King*
Ontario Ministry of Natural Resources, Peterborough

Laurie Maynard
Canadian Wildlife Service, Environment Canada, London

Brian Potter
Ontario Ministry of Natural Resources, Peterborough

Tanya Pulfer*
Ontario Ministry of Natural Resources, Peterborough

ADDITIONAL CONTRIBUTORS

The contributions of the following people who provided information or participated in the writing and reviewing of this report are gratefully acknowledged. The affiliations listed reflect those in effect in the late 1990s, when the information for the Atlas was being compiled.

Garry Allen, Ontario Ministry of Natural Resources
Jim Atkinson, Ontario Ministry of Natural Resources
Wasyl Bakowsky, Natural Heritage Information Centre, Ontario Ministry of Natural Resources
Bonnie Bannerman, Natural Heritage Information Centre, Ontario Ministry of Natural Resources
Dave Bell, Ontario Ministry of Natural Resources
Glenn Birch, Ontario Ministry of Natural Resources
Ron Black, Ontario Ministry of Natural Resources
Karen Bray, Lake Superior Remedial Action Plan
Mike Cadman, Canadian Wildlife Service, Environment Canada
Victor Cairns, Fisheries and Oceans Canada
Jim Cameron, Ontario Ministry of Natural Resources
Daniel Campbell, Consultant
Ross Cholmondeley, Ontario Ministry of Natural Resources
Robin Craig, Ontario Ministry of Natural Resources
Ken Cullis, Lake Superior Remedial Action Plan
Alan Dextrase, Ontario Ministry of Natural Resources
Doug Dodge, Ontario Ministry of Natural Resources
Lesley Dunn, Canadian Wildlife Service, Environment Canada
Andre Dupont, Ontario Ministry of Natural Resources
Mike Eckersley, Ontario Ministry of Natural Resources
Helen Godschalk, Natural Heritage Information Centre, Ontario Ministry of Natural Resources
Tamara Gomer, Ontario Ministry of Natural Resources
Ruth Grant, Ontario Ministry of Natural Resources
Sue Greenwood, St. Marys River Remedial Action Plan
Larry Halyk, Ontario Ministry of Natural Resources
Karen Hartley, Moira River Conservation Authority
Cliff Hatchey, Ontario Ministry of Natural Resources
Mark Heaton, Ontario Ministry of Natural Resources
Don Hector, Ontario Ministry of Natural Resources
Glen Hooper, Ontario Ministry of Natural Resources
Pud Hunter, Ontario Ministry of Natural Resources
Fred Johnson, Ontario Ministry of Natural Resources
Scott Jones, Ontario Ministry of Natural Resources
Al Kalas, Ontario Ministry of Natural Resources
Donald Kirk, Ontario Ministry of Natural Resources
Harold Leadlay, Environment Canada
Jennifer Line, Natural Heritage Information Centre, Ontario Ministry of Natural Resources
Arunas Liskauskas, Ontario Ministry of Natural Resources
Kevin Loftus, Ontario Ministry of Natural Resources
Neil MacLean, Ontario Ministry of Natural Resources
Mike Malhiot, Ontario Ministry of Natural Resources
Dan Mansell, Ontario Ministry of Natural Resources
Alastair Mathers, Ontario Ministry of Natural Resources
Gary McCullough, Canadian Wildlife Service, Environment Canada
Andy McKee, Ontario Ministry of Natural Resources
Dave McLachlin, Ducks Unlimited Canada
Mike McMurtry, Natural Heritage Information Centre, Ontario Ministry of Natural Resources
Bruce McPherson, Hamilton Region Conservation Authority
Linda Melnyk-Ferguson, Ontario Ministry of Natural Resources
John Milcz, Ontario Ministry of Natural Resources
Al Murray, Ontario Ministry of Natural Resources
Todd Norris, Ontario Ministry of Natural Resources
Mike Oldham, Natural Heritage Information Centre, Ontario Ministry of Natural Resources
Nancy Patterson, Canadian Wildlife Service, Environment Canada
Dave Richards, Ontario Ministry of Natural Resources
Bridget Schulte-Hostedde, Consultant
Christine Selinger, Ontario Ministry of Natural Resources
Wayne Selinger, Ontario Ministry of Natural Resources
Elizabeth Snell, Snell and Cecile Environmental Research
Barry Snider, Ontario Ministry of Natural Resources
Steve Suke, Lakehead Region Conservation Authority
Don Sutherland, Natural Heritage Information Centre, Ontario Ministry of Natural Resources
Richard Thompson, Ontario Ministry of Natural Resources
Shawn Thompson, Ontario Ministry of Natural Resources
Ken Tuininga, Ontario Ministry of Natural Resources
Steve Varga, Ontario Ministry of Natural Resources
Donna Wales, Ontario Ministry of Natural Resources
Maggie Young, Canadian Wildlife Service, Environment Canada

Executive Summary

Information and needs assessments with respect to wetlands in the Great Lakes basin consistently call for digital information on the location, distribution, relative significance and status of wetlands. The Ontario Great Lakes Coastal Wetland Atlas was prepared in response to various initiatives such as the State of the Lakes Ecosystem Conference (SOLEC) and the Great Lakes Wetlands Conservation Action Plan (GLWCAP) that pointed to the need for a basin-wide, comprehensive and digital database of coastal wetlands in the Great Lakes basin.

The Atlas represents a compilation of information on coastal wetlands of the Great Lakes and connecting channels in Ontario and provides a framework for the development of a comprehensive digital coastal wetland database for Ontario. The best available sources at the time of Atlas preparation (late 1990s) were used to produce the Atlas, including Ontario Ministry of Natural Resources Wetland Evaluation reports, which provide a consistent approach to evaluating and mapping wetlands, the Natural Heritage Information Centre's Natural Areas Database and Environment Canada's Great Lakes Environmental Sensitivity Atlases. These sources and others were consolidated to provide data such as the location, area and type of wetlands, significance and status, significant species within the wetlands, as well as a qualitative assessment of stressors and major threats affecting coastal wetlands.

Summaries of coastal wetlands in each major lake and connecting channel of the basin are accompanied by detailed maps and appendices. Results indicate that numerous significant species use these primarily marsh-type wetlands, but that human-induced wetland stressors remain prevalent and incremental and site-specific losses continue. There is no comprehensive estimate of historical and recent wetland loss, nor have wetland gains been quantified.

While a great deal of coastal wetland information exists, spatial and temporal gaps in data are common. Spatially, gaps in wetland evaluations are more extensive in the upper lakes and connecting channels where Atlas methodologies have identified that numerous wetlands exist in these areas that have yet to be evaluated by the OMNR Wetland Evaluation System. It is suspected that many of these wetlands may be provincially significant, or contain features to support other conservation designations. There are also temporal gaps in the available data, in that the state of existing wetlands may differ from what the Atlas is able to show. Well over half of the OMNR evaluations took place prior to 1988, within the first five years of implementation of the evaluation system. These evaluations may no longer be accurate, especially in light of changing water levels and the incremental losses that continue to occur throughout the basin.

The uncertainty of loss rates, the known significance of evaluated wetlands, and the gaps in wetland evaluations reinforce the need to achieve the ultimate goal of a larger, interactive database for the future protection and management of basin wetlands.

Sommaire

Les évaluations de l'information et des besoins concernant les terres humides du bassin des Grands Lacs font sans cesse appel à de l'information numérique sur l'emplacement, la répartition, l'importance relative et la situation des terres humides. L'Atlas des terres humides riveraines de l'Ontario a été préparé en réaction à diverses initiatives telles que la Conférence sur l'état de l'écosystème des lacs (CEEL) et le Plan d'action en matière de conservation des terres humides des Grands Lacs (GLWCAP), qui ont souligné le besoin d'une base de données numérique exhaustive, à l'échelle du bassin, sur les terres humides côtières du bassin des Grands Lacs.

L'Atlas réunit des renseignements sur les terres humides côtières des Grands Lacs et les canaux interlacustres de l'Ontario, et offre un cadre pour la constitution d'une base de données numérique exhaustive sur les terres humides côtières de l'Ontario. Les meilleures sources disponibles au moment de la préparation de l'Atlas (fin des années 1990) ont servi à sa production, y compris les rapports d'évaluation des terres humides du ministère des Richesses naturelles de l'Ontario, qui utilisent une approche cohérente de l'évaluation et de la cartographie des terres humides; la base de données sur les zones naturelles du Centre d'information sur le patrimoine naturel; et les Atlas de sensibilité environnementale des Grands Lacs d'Environnement Canada. Ces sources ont été réunies, ainsi que d'autres, pour offrir des données telles que le lieu, la superficie et le type des terres humides, leur importance et leur état, les espèces importantes qu'elles accueillent, ainsi qu'une évaluation qualitative des facteurs de stress et les principales menaces qui affectent les terres humides côtières.

Des résumés sur les terres humides côtières de chaque lac et chaque canal interlacustre d'importance dans le bassin sont accompagnés de cartes et d'annexes détaillées. Les résultats indiquent que de nombreuses espèces d'importance utilisent ces terres humides composées surtout de marécages, mais que les facteurs d'agression anthropiques des terres humides demeurent prépondérants et que les pertes graduelles et sitospécifiques se poursuivent. Il n'y a pas d'estimation globale des pertes historiques et récentes des terres humides, pas plus qu'il n'y a de quantification des gains réalisés par les terres humides.

Bien qu'il existe une foule de renseignements sur les terres humides côtières, les données présentent couramment des lacunes sur les plans spatial et temporel. Sur le plan spatial, les lacunes dans les évaluations des terres humides sont plus prononcées dans les lacs ou les canaux interlacustres supérieurs où les méthodologies de l'Atlas ont déterminé qu'il reste encore de nombreuses terres humides à évaluer par le système d'évaluation des terres humides du MRLO. On soupçonne que bon nombre de ces terres humides sont d'importance provinciale, ou présentent des caractéristiques qui les rendent admissibles à d'autres types de désignation en matière de conservation. Il y a aussi des lacunes temporelles dans les données disponibles, dans la mesure où l'état des terres humides actuelles peut différer de ce que l'Atlas est en mesure de révéler. Bien au-delà de la moitié des évaluations du MRLO ont eu lieu avant 1988, dans les cinq premières années suivant la mise en place du système d'évaluation. Il se peut que ces évaluations ne soient plus à jour, compte tenu surtout de l'évolution du niveau des eaux et des pertes graduelles qui continuent de survenir dans l'ensemble du bassin.

L'incertitude quant aux taux de perte, l'importance connue des terres humides évaluées et les lacunes des évaluations des terres humides accentuent le besoin d'en arriver au but ultime, soit une base de données interactive plus vaste pour la protection et la gestion futures des terres humides du bassin.

TABLE OF CONTENTS

Acknowledgements i

Executive Summary ii

Sommaire iii

List of Appendices, Figures, and Tables iv

1.0 Introduction 1
2.0 Background 1

3.0 Methods 2
- 3.1 Background 2
- 3.2 Data Sources 3
- 3.3 Methodology 3
  - 3.3.1 Data Sources 3
  - 3.3.2 Data Compilation 5

4.0 Atlas Summary and Future Plans 6
- 4.1 Accomplishments 6
- 4.2 Key Findings
  - 4.2.1 General /Information/Significance 6
  - 4.2.2 Stressors 6
  - 4.2.3 Significant Species 7
  - 4.2.4 Identification of Gaps 7
- 4.3 Future Plans 7

5.0 Results 13
- 5.1 Lake Superior 13
  - 5.1.1 Setting 13
  - 5.1.2 Significant Features 13
  - 5.1.3 Wetland Status 14
- 5.2 St. Marys River 15
  - 5.2.1 Setting 15
  - 5.2.2 Significant Features 15
  - 5.2.3 Wetland Status 16
- 5.3 Lake Huron 17
  - 5.3.1 Setting 17
  - 5.3.2 Significant Features 18
  - 5.3.3 Wetland Status 19
- 5.4 St. Clair River 20
  - 5.4.1 Setting 20
  - 5.4.2 Significant Features 20
  - 5.4.3 Wetland Status 21
- 5.5 Lake St. Clair 22
  - 5.5.1 Setting 22
  - 5.5.2 Significant Features 23
  - 5.5.3 Wetland Status 24
- 5.6 Detroit River 25
  - 5.6.1 Setting 25
  - 5.6.2 Significant Features 25
  - 5.6.3 Wetland Status 26

5.7 Lake Erie 28
    5.7.1 Setting 28
    5.7.2 Significant Features 28
    5.7.3 Wetland Status 30
5.8 Niagara River 32
    5.8.1 Setting 32
    5.8.2 Significant Features 32
    5.8.3 Wetland Status 33
5.9 Lake Ontario 34
    5.9.1 Setting 34
    5.9.2 Significant Features 35
    5.9.3 Wetland Status 36
5.10 St. Lawrence River 38
    5.10.1 Setting 38
    5.10.2 Significant Features 39
    5.10.3 Wetland Status 39

6.0 References 42

Appendices ..... A-1 to V-1

Figures ..... envelope at back

LIST OF APPENDICES

Appendix A: 1996 Ontario Ministry of Natural Resources District/Area Offices Survey

Appendix B: A Guide to Appendices E, G, I, K, M and O – Evaluated Coastal Wetlands

Appendix C: A Guide to Appendices F, H, J, L, and N – Natural Areas, Environmental Sensitivity Atlas Wetlands and Other Wetlands with an Unevaluated Coastal Wetland Component

Appendix D: A Guide to Appendices P through U – Significant Species Reported in Ontario's Great Lakes Coastal Wetlands

Appendix E: Lake Superior: Evaluated Coastal Wetlands

Appendix F: Lake Superior: Natural Areas, Environmental Sensitivity Atlas Wetlands and Other Wetlands with an Unevaluated Coastal Wetland Component

Appendix G: Lake Huron and St. Marys River: Evaluated Coastal Wetlands

Appendix H: Lake Huron and St. Marys River: Natural Areas, Environmental Sensitivity Atlas Wetlands and Other Wetlands with an Unevaluated Coastal Wetland Component

Appendix I: Detroit River, Lake St. Clair and St. Clair River: Evaluated Coastal Wetlands

Appendix J: Detroit River, Lake St. Clair and St. Clair River: Natural Areas, Environmental Sensitivity Atlas Wetlands and Other Wetlands with an Unevaluated Coastal Wetland Component

Appendix K: Lake Erie: Evaluated Coastal Wetlands

Appendix L: Lake Erie: Natural Areas, Environmental Sensitivity Atlas Wetlands and Other Wetlands with an Unevaluated Coastal Wetland Component

Appendix M: Lake Ontario and Niagara River: Evaluated Coastal Wetlands

Appendix N: Lake Ontario and Niagara River: Natural Areas, Environmental Sensitivity Atlas Wetlands and Other Wetlands with an Unevaluated Coastal Wetland Component

Appendix O: St. Lawrence River: Evaluated Coastal Wetlands

Appendix P: Confirmed Records of Significant Vascular Plant Species Reported in Ontario's Great Lakes Coastal Wetland Evaluations

Appendix Q: Significant Fish Species Reported in Ontario's Great Lakes Coastal Wetland Evaluations

Appendix R: Significant Reptile and Amphibian Species Reported in Ontario's Great Lakes Coastal Wetland Evaluations

Appendix S: Significant Bird Species Reported in Ontario's Great Lakes Coastal Wetland Evaluations

Appendix T: Significant Lepidopteran Species Reported in Ontario's Great Lakes Coastal Wetland Evaluations

Appendix U: Unconfirmed Records of Significant Vascular Plant Species Reported in Ontario's Great Lakes Coastal Wetland Evaluations

Appendix V: Scientific Names of Common (Non-Significant) Species mentioned in the Atlas

LIST OF TABLES

Table 1: Summary of Ontario Great Lakes Coastal Wetland Atlas Results 10

Table 2: Summary of Evaluated Ontario Great Lakes Coastal Wetlands 11

Table 3: Summary of Unevaluated Ontario Great Lakes Coastal Wetlands 12

Table 4: Summary of Significant Species Reported in Ontario Great Lakes Coastal Wetlands 13

LIST OF FIGURES

** Please note the figures are located in an envelope at the end of the report.*

Figure 1: Lake Superior Coastal Wetlands

Figure 2: Lake Huron and St. Marys River Coastal Wetlands

Figure 3: Detroit River, Lake St. Clair and St. Clair River Coastal Wetlands

Figure 4: Lake Erie Coastal Wetlands

Figure 5: Lake Ontario and Niagara River Coastal Wetlands

Figure 6: St. Lawrence River Coastal Wetlands

1.0 Introduction

Despite initiatives undertaken by partnerships such as the Eastern Habitat Joint Venture and the Great Lakes Wetlands Conservation Action Plan, leading to advances in wetland policy, restoration and conservation, wetland area and quality and the number of significant wetland-dependent species continue to decline. The most recent account of wetland loss in southern Ontario (Snell, 1984) is over 20 years old, and although there have been numerous site-specific descriptions of wetland loss, a comprehensive review does not exist. To keep track of current wetlands and monitor future losses, a comprehensive database of wetlands in the Great Lakes basin is needed. The Ontario Great Lakes Coastal Wetland Atlas represents a significant step toward an interactive database of the status and trends of coastal and inland wetlands of the Great Lakes basin.

The goals of the Ontario Great Lakes Coastal Wetland Atlas and future applications include:

- developing a database with complete geographic coverage of coastal wetlands of the Great Lakes basin, with a consistent evaluation methodology and current information sources;
- developing a database that is digital, with each wetland geo-referenced, which will serve as a registry for new wetland identification or evaluation;
- identifying gaps in wetland data and strategically planning to fill these gaps;
- assessing and monitoring the contribution of wetlands to biodiversity conservation; and,
- mapping wetland polygons with complete geographic coverage of the Great Lakes basin with links to summary data for each wetland to provide a visual distribution of wetlands and their relative significance.

Information on wetlands is required by many national, provincial and local governments, non-government agencies and individuals interested in wetland conservation and restoration. The information from the Atlas will be used to identify inventory and information gaps, assess the status, losses of and changes to coastal wetlands, identify natural heritage areas for land use planning purposes and establish priority areas for conservation. From a management and enhancement perspective, the content will help develop habitat management strategies, target limited resources for conservation and restoration programs, and contribute to setting goals for biodiversity conservation and recovery of species at risk in wetlands.

2.0 Background

Several initiatives have documented the need for a complete inventory of wetlands in the Great Lakes basin in order to better track and monitor wetland status and change, loss and restoration efforts. In part, the Ontario Great Lakes Coastal Wetland Atlas was produced in response to the work of the 1996 State of the Lakes Ecosystem Conference (SOLEC) and the Great Lakes Wetlands Conservation Action Plan (GLWCAP). SOLEC is held biennially in support of the Great Lakes Water Quality Agreement (GLWQA), under which the United States and Canada committed to the restoration and maintenance of the chemical, physical and biological integrity of the Great Lakes basin ecosystem. The 1996 SOLEC examined the health of aquatic and terrestrial communities and the biological habitat of nearshore ecosystems of the Great Lakes. The background paper presented on coastal wetlands concluded that there was no comprehensive current inventory or evaluation of Great Lakes coastal wetlands (Maynard and Wilcox, 1997). A review of all available data sets identified in the Catalogue of Wetland Databases and Inventories for the Canadian Great Lakes basin (Environment Canada, 1995) supported this conclusion. This lack of information on Great Lakes wetlands was also recognized in the Great Lakes Wetlands Conservation Action Plan, which calls for development of an accessible, computerized, database for Great Lakes basin wetlands.

The Atlas responds to the need to develop a coastal wetlands database for the Great Lakes by updating and consolidating information on coastal wetlands from the best available sources including OMNR

District/Area Office Surveys and Wetland Evaluations, Environment Canada's Great Lakes Environmental Sensitivity Atlases, the Natural Heritage Information Centre's (NHIC) Natural Areas Database, the NHIC Coastal Meadow Marsh Summary and other site specific studies. This information in turn provides:

- a framework for the development of a comprehensive and current Ontario wetland database;
- detailed, spatially referenced information that can be analyzed at a variety of levels for Great Lakes and connecting channels;
- information on wetland type (marsh, fen, bog, swamp), site type (palustrine, lacustrine, riverine, isolated) and associated functions and values;
- information on wetland significance and status; and,
- a qualitative assessment of stressors affecting coastal wetlands and data on biodiversity with respect to significant species dependent on coastal wetlands.

Typical of such large-scale projects, the Ontario Great Lakes Coastal Wetland Atlas data and results have limitations that must be recognized. For example, trend through time information is not available because each record represents a single evaluation at one point in time. The wetland evaluations themselves are of varying quality and do not provide comprehensive biophysical inventories. The evaluations are not comprehensive because the evaluation system only requires that certain information is recorded for each wetland, such as the presence of provincially significant plant and wildlife species, while more general information, such as the presence of common species, is omitted. Further, although attempts were made to update wetland status where possible, this information was compiled in the late 1990s and dates back to 1983. For example, lower water levels since the late 1990s have likely resulted in many wetlands that were not previously identified.

Additionally, the limitations of statistical analysis of the data should be recognized and respected. The data gaps previously identified and the variations in effort (e.g., number of people conducting wetland evaluations, time spent at a site, varying experience of evaluators) that might contribute to inaccurate results must be taken into consideration upon analysis. The Atlas lays the foundation to facilitate the tracking of historical wetland evaluations and wetland changes, permit the identification of data gaps, and provide baseline data on coastal wetlands.

3.0 Methods

3.1 Background

An important issue that arose from SOLEC 1996 was that current mapping methods did not effectively capture changes to wetlands. For example, site-scale changes were not always captured by air photos, and Landsat Thematic Mapper data (30 m spatial resolution) did not provide accurate information on the location, size and type of wetlands. Participants in a 1997 workshop on remote sensing and wetlands concluded that information collected through site-level wetland evaluations and studies is critical for providing a large-scale summary of the location, characteristics, and status of Great Lakes wetlands (Riley and Snell, 1997).

For years, wetland scientists had identified that existing information on coastal wetlands in the Great Lakes basin was piecemeal and dated. In May 1996 a survey was sent to Ontario Ministry of Natural Resources District/Area Offices bordering the Great Lakes. The purpose of this survey was to help determine the current status of Great Lakes coastal wetlands. The results provided summaries of the main characteristics of these wetlands, identified data gaps and identified major stressors affecting specific coastal wetlands.

The findings of SOLEC 1996 (Maynard and Wilcox, 1997), the outcomes of the remote sensing workshop, the results of the OMNR survey, the ultimate goals of the Atlas, and the criteria outlined in the following section all contributed to the selection of the data sources.

3.2 Data Sources

The following general criteria were established for data selection:

- data sets chosen must result in full geographic coverage of the Ontario shoreline of the Great Lakes and connecting channels;
- data must show coastal wetlands within 2 kilometres of the shore;
- data must reflect aerial extent of wetlands;
- data must be at a meaningful scale to identify extent, location, and type;
- data must be accessible; and,
- the integrity of the data must be known (source, collection method, consistency).

The aim was to select the best information available that satisfied these criteria and the needs and goals of the Atlas. No one data set met all the criteria. The Atlas is the result of the compilation and integration of the following data sets.

1. OMNR Wetland Evaluation records (1983 to 1997)
2. Environment Canada's Great Lakes Environmental Sensitivity Atlases (1991 to 1994)
3. Natural Heritage Information Centre's Natural Areas Database
4. 1996 OMNR District/Area Offices survey
5. Natural Heritage Information Centre's Significant Species Lists (1999)
6. Coastal Meadow Marsh Summary (NHIC, 1995)

3.3 Methodology

For the purposes of the Atlas, coastal wetlands were considered to be those wetlands within 2 kilometres of the shoreline, greater than 2 hectares in size. Two hectares is the minimum area for a non-complexed wetland to be included in the OMNR Wetland Evaluation Protocol.

3.3.1 Data Sources

3.3.1.1 Ontario Ministry of Natural Resources Wetland Evaluation Records (1983 to 1997), Ontario Ministry of Natural Resources District Offices

The Wetland Evaluation records provided a consistent approach to evaluating and mapping wetlands across the basin, thereby enabling distribution and relative significance comparisons. They also provided information on size, type, site type, ownership data and significant species information. OMNR Wetland Evaluations are the most detailed surveys available for wetlands in Ontario. The Wetland Evaluation reports were collected from OMNR District Offices in June 1997. The evaluations ranged from 1983 to 1997, however the majority dated from the early to mid-1980s. Three editions of the Wetland Evaluation System were in use over this time period (OMNR, 1993a,b, 1984, 1983). Data in the evaluations were interpreted and verified (Appendix B).

For evaluated wetlands, only evaluation information completed pre-1997 is included in the Atlas. Information from other sources is presented for unevaluated wetlands. The data source, most recent year and method of evaluation were recorded for each wetland. The most recent evaluation was used wherever possible. "DU" after the year of evaluation indicates a desktop update edition of the evaluation. In the case of desktop updates, previous edition evaluation information is transferred to a new edition evaluation without new fieldwork or research. Thus a 1996DU evaluation does not have the same currency of information that an evaluation actually carried out in 1996 would.

Using UTM coordinates, each wetland was plotted in ARCVIEW as a point on a map using a pie-shaped symbol (Figures 1 to 6). Large symbols were used for Provincially Significant Wetlands, while smaller symbols represent Non-Provincially Significant Wetlands. Summary tables providing information such as wetland size, type and significance, based on evaluation results, were created in Excel (Appendices E, G, I, K, M, O). Detailed information for each OMNR-evaluated wetland was entered into the Natural Areas Database (Microsoft Access) and each wetland was assigned a Natural Area Number and cross-referenced to the digital maps (Figures 1 to 6). The NHIC's Natural Areas Database therefore includes all the information from the summary tables (Appendices E, G, I, K, M, O) along with lists of dominant vegetation forms and communities, biodiversity lists, and information on landforms/soils, ecological and other values, land use, offsite uses, wetland stressors, rating/scoring evaluation edition, chronology of evaluations and references. Publicly accessible NHIC Natural Areas data can be found at www.mnr.gov.on.ca/MNR/nhic/areas.cfm. Where vegetation community and plant information was provided, an assessment of plant community significance was made (based on W. Bakowsky, pers. com., 1998).

Unevaluated wetlands identified through other data sources (described below) were plotted in a similar manner, using alphanumeric characters based on the sources (Figures 1 to 6). Unevaluated wetlands are under-represented in the Atlas.

3.3.1.2 Environmental Sensitivity Atlases (1991 to 1994), Environmental Protection Branch, Environment Canada

The Great Lakes Environmental Sensitivity Atlases (GLESAs) provided information on wetland features that could be used to identify unevaluated coastal wetlands plus additional information on adjacent land uses. The information contained in the Sensitivity Atlases was compiled by Environment Canada between 1991 and 1994 for use in the preparation and response to spills of oil and other chemicals in the Great Lakes and connecting channels. They were created to provide a visual reference for general environmental protection initiatives and for increasing the awareness of sensitive shoreline areas. The data were compiled primarily from species surveys, OMNR Wetland Evaluations, aerial video photography of the shoreline, and ground surveys. The shoreline classification categories in the Sensitivity Atlases that correspond to wetlands are low vegetated banks, delta mudflats, fringing wetlands and broad wetlands.

Once the OMNR-evaluated wetlands were identified, the Sensitivity Atlases were reviewed for each of the Great Lakes and unevaluated wetlands were identified. Many shoreline areas showed extensive fringing and broad wetlands in the Sensitivity Atlases, but have not been evaluated formally as wetlands by any government agency. Therefore only wetlands that were identified in the Sensitivity Atlases and also recognized by these agencies were included in the Atlas. Where wetland evaluations did not exist for the wetlands identified in the Environmental Sensitivity Atlases, the UTM coordinates were obtained from topographic maps, added to the database, and the wetland was plotted in ARCVIEW and identified on the maps with a star-shaped symbol (Figures 1 to 6).

3.3.1.3 Natural Areas Database, Natural Heritage Information Centre

The Natural Heritage Information Centre has developed a Natural Areas Database (NAD) in which information on natural areas such as wetlands can be entered and stored. Currently, it holds more than 6,000 records and is being updated and revised on an ongoing basis. Many coastal wetlands support significant features and are therefore included in this database. In order to identify natural areas with coastal wetland features, all natural areas within 2 kilometres of the Great Lakes shoreline were identified using ARCVIEW, and the descriptions of each natural area (such as Areas of Natural and Scientific Interest (ANSI), International Biological Program Sites (IBP), Wildlife Areas, Conservation Areas, etc.), were reviewed to determine if they contained representative coastal wetland features (see Appendix C). Using the coordinates, these were plotted on a map using red squares (Figures 1 to 6). Information from the NAD was included in summary tables (Appendices E to O).

3.3.1.4 1996 Ontario Ministry of Natural Resources District/Area Offices survey

In 1996, as part of this project and as background for SOLEC 1996, a survey (Appendix A) was sent to OMNR District and Area Offices bordering the Great Lakes. The results presented in the tables, appendices, and throughout the Atlas augment the Wetland Evaluations and update the current status of Great Lakes coastal wetlands based on staff knowledge, where coastal wetland status had changed since evaluation. It also served to identify unevaluated wetlands (Table 3). The results of this survey also provided summaries of the main characteristics of Ontario coastal wetlands, and identified data gaps and major stressors affecting specific coastal wetlands.

3.3.1.5 Significant Species Lists, Natural Heritage Information Centre

The significant species lists provided extensive information on significant species and wetland features that could be used to augment Wetland Evaluation information. Occurences of rare flora and fauna compiled by the NHIC were used to identify rare species occurring in Ontario. A comprehensive list of rare species that utilize coastal wetlands was then generated, based on an extensive literature review and information provided by biologists within the NHIC and OMNR. These species are ones found in coastal areas or areas with wetlands, but are not necessarily obligate wetland species. The list includes significant plant, bird, amphibian, reptile, fish, and lepidopteran species; there are currently no provincially significant mammal species listed that use coastal wetlands in the Great Lakes basin.

Because of the potential sensitivity of rare species information, locations of specific rare species were noted only by identifying the corresponding lake or connecting channel (Appendices P to U) in this report. Wherever possible, the location of rare species based on NHIC data and OMNR Wetland Evaluations were linked to each wetland and included in the Atlas (Appendices E, G, I, K, M, O).

Species listed in the Significant Species Tables (Appendices P to U), are those rare species with a provincial rank (S Rank) of S1, S2, S3 or S3S4, as designated by the NHIC. Species of these ranks are considered to be "provincially significant". Rare species identified in these tables have been taken from various data sources, and may not qualify for points under the Special Features Component of the Ontario Wetland Evaluation System (OMNR, 1993a). As well, some records of rare species listed in OMNR Wetland Evaluations have not been confirmed by NHIC staff (Appendix U). For bird species, two S-Ranks are assigned: breeding (B) and non-breeding (N) status. For the purposes of the Atlas, the S-Rank with the highest significance or that which is the limiting habitat (usually breeding) is used.

The Significant Species Tables (Appendices P to U) also identify the status assigned to these species by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC November 2002) and the Committee on the Status of Species at Risk in Ontario (COSSARO September 2002). A more detailed explanation of the type of data found in the Significant Species Tables is in Appendix D.

3.3.1.6 Coastal Meadow Marsh Summary, Natural Heritage Information Centre

NHIC conducted a literature review to summarize knowledge of coastal meadow marshes (shoreline fens) (NHIC, 1995). Where the UTM coordinates of the Great Lakes coastal meadow marshes overlapped with Environmentally Sensitive Areas (ESA), Natural Areas and evaluated wetlands, a reference was made in the Atlas.

3.3.2 Data Compilation

The Great Lakes Environmental Sensitivity Atlases, OMNR-evaluated wetlands, and the Natural Areas with coastal wetlands were plotted and overlapping areas were identified. Where Natural Areas overlapped with evaluated wetlands or Great Lakes Environmental Sensitivity Atlas wetlands, they were recorded on the maps using red squares around the numbers of the pie and star-shaped symbols (Figures 1 to 6), and the unique Natural Area Number was recorded (Appendices E, G, I, K, M, O). Where a Natural Area with a wetland component occurred within 2 kilometres of the shoreline, and did not correspond with an evaluated or Environmental Sensitivity Atlas wetland it was added to the Unevaluated Wetland list, given a unique identifier and plotted separately using an solid red square symbol. Summaries were created for the Natural Area wetlands, GLESA wetlands and other unevaluated wetlands, providing location information, a brief description of the wetland components, and presence/absence of significant species (Appendices F, H, J, L, N).

4.0 Atlas Summary and Future Plans

4.1 Accomplishments

The Ontario Great Lakes Coastal Wetland Atlas has provided a preliminary snapshot of the state of the Great Lakes wetland ecosystem in the late 1990s, in terms of the quality and quantity of both the available wetland-related information and the wetland resource itself. These results will help direct future wetland evaluation, protection, and management efforts. Several of the goals outlined in the Introduction have been achieved laying the groundwork for the basin-wide, interactive digital database. Coastal wetlands have been geo-referenced, therefore a consistent naming system exists, ultimately reducing duplication and confusion of sites.

4.2 Key Findings

4.2.1 General Information/Significance

The Ontario Great Lakes Coastal Wetland Atlas provides general information on the number, extent and type of wetlands in the basin. There are 236 OMNR-evaluated coastal wetlands and over 170 unevaluated coastal wetlands identified from the Natural Areas Database, Great Lakes Environmental Sensitivity Atlases, and other sources (Table 1). When evaluated and unevaluated wetlands are considered, all four wetland types (marsh, swamp, bog and fen) are found on virtually every lake and connecting channel. Lake Erie has the largest total area of evaluated wetlands at almost 20,000 ha.

Individual evaluated wetlands range in size from 2.0 ha (on Lake Ontario) to 13,465 ha (on Lake Erie) and cover over 53,600 ha (Table 2). Marshes make up over three-quarters of evaluated coastal wetland area in the basin; the majority of the remaining area is swamp. The majority of the over 170 unevaluated coastal wetlands also appear to be marsh. The Lake Huron/St. Marys River region has the most unevaluated wetlands, with 151 recorded (Table 3).

Despite the losses and impacts to Great Lakes coastal wetlands, they have retained remarkable characteristics such as the complex vegetation communities of Lake Huron and the great biodiversity of Lake Erie. Three-quarters of the evaluated wetlands are provincially significant (in part because priority was given to evaluating those wetlands thought to be provincially significant), and many evaluated and unevaluated wetlands also have special designations such as Areas of Natural and Scientific Interest or International Biological Program Sites (Table 1). Many of the unevaluated wetlands are suspected to be provincially significant, and could potentially contain species at risk or their habitat. This has implications for setting goals for biodiversity conservation and species at risk recovery.

4.2.2 Stressors

The Atlas outlines the major stressors encountered in the coastal wetlands of the lakes and connecting channels (Appendices E, G, I, K, M, O). All areas have been affected by agricultural drainage, urban encroachment, shoreline modification and sediment contamination. Wetlands in connecting channels that serve as shipping lanes are further affected by wave action and dredging. The lower Great Lakes are particularly impacted by industry outfalls, sewage treatment, and toxic chemical contamination. The more northerly sites, such as Lakes Superior and Huron and the St. Marys River, are comparatively less affected by human stressors. The relative remoteness of these northern wetlands results in more gaps in evaluation data, but less urgency to identify the provincially significant sites that could be afforded protection from human impacts. Significant gaps in evaluation data in the highly pressured lower lakes and channels were also found (Table 1, Appendix M).

4.2.3 Significant Species

A significant number of provincially significant species are found in Great Lakes coastal wetlands (Table 4), including a number of species at risk (Appendices P to U). Of the 45 species listed in the Atlas as at risk by COSEWIC, one-third are endangered, one-third are threatened, one species is vulnerable, and one-third are species of special concern. Provincially, of the 30 species listed in Atlas that are considered at risk by COSSARO, eight are endangered, 13 are threatened and 12 are vulnerable (Appendices P to U).

Lake Erie and Lake Huron are the most biodiverse and important sites in terms of significant species: approximately half of the provincially significant plants, birds, herptiles, fish and lepidoptera of coastal wetlands are found in these two lakes. These significant species were found in a large number of wetlands, many of which are provincially significant, which affords them some protection. There is great potential that species at risk or their habitats will be found in unevaluated coastal wetlands once they are evaluated, particularly in ANSIs. Since the time of many wetland evaluations (mid-1980s) and preparation of the Atlas (late 1990s), a number of wetland dependent species have suffered dramatic population declines (e.g., King Rail, Fowler's Toad, Bluehearts, and Pugnose Shiner).

4.2.4 Identification of Information Gaps

One goal of the Atlas was to identify information gaps. Gaps in wetland evaluations are more extensive in the upper lakes and connecting channels where Natural Area and Great Lakes Environmental Sensitivity Atlas information has identified that numerous wetlands exist in these areas that have yet to be evaluated by the OMNR Wetland Evaluation System (see Table 1 and Appendix G). It is suspected that there may also be provincially significant and other important wetlands that have not yet been identified, evaluated, or assigned a conservation status designation. In addition, results of the OMNR District/Area Offices survey identified numerous wetlands that had not yet been evaluated (Table 3).

There are also temporal gaps in the available data, in that the state of existing wetlands may differ from what the Atlas is able to show. Well over half of the OMNR evaluations took place prior to 1988, within the first five years of implementation of the evaluation system (OMNR 1984). These evaluations may no longer be accurate, especially in light of changing water levels and the incremental losses that continue to occur throughout the basin. The extent of both the historical widespread and current incremental losses of wetlands reported for all lakes and connecting channels is unknown. Wetland evaluations can now become more strategic due to the awareness of both spatial and temporal gaps in wetland identification and evaluation.

4.3 Future Plans

The information contained in the Atlas is not exhaustive; rather it presents a late 1990s snapshot in time and an overview of the type of information that can be obtained from such a database. Although this is the most comprehensive map of coastal wetlands in Ontario, and serves as a framework for future data collection, analysis and application, it is already dated and further work would be required to complete the database described in the goals. The completion and electronic maintenance of the coastal wetland database and expansion to include inland wetlands may become part of OMNR's Natural Resource Values Information System and Wetland Evaluation Information Management System. Ideally, all the wetlands in the basin would be identified and evaluated using the OMNR evaluation system. Further analysis of existing mapping is also a priority, as is the collection of additional biophysical information (such as water level fluctuations and near-shore bathymetry) and the creation of wetland polygons with links to summary information. Remote sensing will continue to be examined as a possible technique for tracking changes to wetland area. Different methods are currently being evaluated. Ultimately, the Ontario Great Lakes Coastal Wetland Atlas will contribute to the development of a basin-wide, interactive map of Great Lakes basin coastal and inland wetlands.

Table 1: Summary of Ontario Great Lakes Coastal Wetland Atlas Results

Lake/ Connecting Channel	Evaluated Wetlands (1983 to 1997)[6]									Unevaluated Wetlands					
	No. Evaluated Wetlands	No. PSW[1]	No. NPSW[2]	Total area of evaluated wetlands (ha)	Designations[4] (an individual wetland may have more than one designation)					Estimated No. unevaluated wetlands[5]	Designations (an individual wetland may have more than one designation)				
					ANSI	PWA	NWA	IBP	Ramsar		ANSI	PWA	NWA	IBP	Ramsar
Lake Superior	11	3	8	1,080.6[3]						13	2			1	
St. Marys River	7	3	4	3,567.0[3]						54	1				
Lake Huron	48	41	7	7,459.0[3]	22	4	1	11	1	97	29			6	
St. Clair River	5	4	1	87.2											
Lake St. Clair	5	5	0	2,522.9[3]	1		1		1	1	1				
Detroit River	4	4	0	1,136.3	1										
Lake Erie	30	26	4	19,329.8	14	1	2	2	2	3	2		1	1	
Niagara River	4	4	0	84.5	1			2							
Lake Ontario	82	64	18	11,334.5[3]	30		1	5		5	2				
St. Lawrence River	40	21	19	7,017.7[3]	11	1		5		[7]					
Total	236	175	61	53,619.5	79	6	5	23	4	173	37	0	1	8	0

[1] PSW = Provincially Significant Wetlands
[2] NPSW = Non-Provincially Significant Wetland
[3] Evaluations incomplete; see also Table 3
[4] ANSI= Area of Natural and Scientific Interest; PWA= Provincial Wildlife Area; NWA= National Wildlife Area; IBP= International Biological Program Site; Ramsar=Wetland of International Importance
[5] Identified from Natural Areas Database, Environmental Sensitivity Atlas, and Other
[6] Number of evaluated wetlands counts a wetland complex as one wetland
[7] Although there are unevaluated wetlands on the St. Lawrence River (Table 3), they have not been individually identified in the Appendices nor maps, and cannot be included in this summary.

Table 2: Summary of Evaluated Ontario Great Lakes Coastal Wetlands

Lake/ Connecting Channel	Evaluated Wetlands[1]			Wetland Area				Wetland Type			
	Number Evaluated Wetlands	Number Provincially Significant	Number Non-Provincially Significant	Total Area (ha)	Mean Size (ha)	Smallest Wetland > 2 ha (ha)	Largest Wetland (ha)	Swamp (ha)	Marsh (ha)	Bog (ha)	Fen (ha)
Lake Superior[2]	11	3	8	1,080.6[2]	98.2	4.6	429.0	611.8[2]	392.5[2]	41.3[2]	32.4[2]
St. Marys River	7	3	4	3,567.0	509.6	42.0	2,275.0	1,387.6	1,724.4	0	455.0
Lake Huron[2]	48	41	7	7,459.0[2]	155.4	5.0	807.4	3,768.8[2]	3,227.7[2]	16.1[2]	447.8[2]
St. Clair River	5	4	1	87.2	17.4	3.0	44.3	6.9	80.3	0	0
Lake St. Clair	5	5	0	2,522.9	504.6	3.9	2,335.0	270.9	2,252.0	0	0
Detroit River	4	4	0	1,136.3	284.1	32.0	575.0	37.1	1,099.2	0	0
Lake Erie	30	26	4	19,329.8	644.3	3.2	13,465.0	1,446.9	17,878.9	4.0	0
Niagara River	4	4	0	84.5	21.2	4.5	37.2	39.3	45.2	0	0
Lake Ontario	82	64	18	11,334.5	138.2	2.0	2,093.0	2,518.4	8,780.7	6.3	29.1
St. Lawrence River	40	21	19	7,017.7	175.4	3.6	1,398.0	1,270.5	5,719.2	28.0	0
Total[2]	236	175	61	53,619.5[2]	227.2	2.0	13,465.0	11,358.2[2]	41,200.1[2]	95.7[2]	964.3[2]

[1] Number of evaluated wetlands counts a wetland complex as one wetland

[2] For two Lake/Connecting Channels, the total areas (ha) of swamp, marsh, bog and fen, do not add up to the total size (ha) of wetland area, due to percentage of wetland type being recorded as either under or over 100% in several original evaluations, particularly Neebing Marsh and Caldwell Lake (Lake Superior) and Wye Marsh (Lake Huron)

Table 3: Summary of Unevaluated Ontario Great Lakes Coastal Wetlands

Atlas methodologies identified a total of 173 unevaluated coastal wetlands (Estimated No. Unevaluated wetlands). The 1996 OMNR District/Area Survey identified that there are likely significantly more coastal wetlands that remain to be evaluated (Evaluation Status). Therefore, the Atlas likely under-represents the total number of coastal wetlands on the Great Lakes. Many of these wetlands have the potential to be provincially significant or contain habitat of significant species.

Lake/ Connecting Channel	**Estimated No. Unevaluated Wetlands** (from Natural Areas, Environmental Sensitivity Atlas, and Other)	**Evaluation Status** (from 1996 OMNR District/Areas Survey)
Lake Superior	13	• approximately five wetlands not evaluated in Goulais Bay/ Batchewana Bay in northeastern Lake Superior; and seven not yet evaluated in Thunder Bay, Black Bay, Nipigon Bay Area in northwestern Lake Superior; several of these wetlands are likely Provincially Significant Wetlands (A. Dupont, pers. com., 1996) • there are no wetlands larger than 2 ha along the remainder of the north shore
St. Marys River	54	• 130+ wetlands larger than 2 ha along shoreline to be evaluated and many more wetlands within 2 kilometres of the St. Marys River (A. Dupont, pers. com., 1996) • number of potential Provincially Significant Wetlands to be evaluated unknown (A. Dupont, pers. com., 1996)
Lake Huron	97	• Wingham Area: evaluations complete • Bruce Peninsula: unknown number of wetlands that need to be evaluated • Midhurst: 5 to 10 wetlands greater than 2 ha need to be evaluated and at least one is a Provincially Significant Wetland (G. Allen, pers. Com., 1996) • Parry Sound: one large wetland and many smaller wetlands (mostly marshes < 10 to 20 ha) need to be evaluated and the number of Provincially Significant Wetlands is unknown (R. Black, pers. com., 1996) • Sudbury/Manitoulin: numerous wetlands larger than 2 ha need to be evaluated, and number of Provincially Significant Wetlands is unknown (W. Selinger, pers. com., 1996)
St. Clair River		• evaluations complete
Lake St. Clair	1	• evaluations complete except Walpole Island (approximately 10,360 ha)
Detroit River		• evaluations complete
Lake Erie	3	• evaluations incomplete
Niagara River		• evaluations complete
Lake Ontario	5	• evaluations complete except Grafton Swamp West (approximately 20 ha) in Northumberland/Hastings County (B. Snider, pers. com., 1996) and several marshes in Bay of Quinte • difficult to estimate number to be evaluated in Bay of Quinte because there are many embayments with cattails or submerged vegetation (N. MacLean, pers. com.,1996) • there are probably no unevaluated Provincially Significant Wetlands on Lake Ontario
St. Lawrence River		• approximately seven wetlands larger than 2 ha have yet to be evaluated and all are probably Provincially Significant Wetlands (one on mainland, approx. 40 ha; and six associated with Akwesasne Islands, approximately 100 ha) (M. Eckersely, pers. com., 1996)
Total	173	

Table 4: Summary of Significant Species Reported in Ontario Great Lakes Coastal Wetlands

Lake/ Connecting Channel	Total Evaluated Wetlands	Total PSW[1]	Total NPSW[2]	Total area of evaluated wetlands (ha)	Number of Significant Species[4] by Group[5] (% of total listed in Atlas by group)					Total Estimated Unevaluated wetlands[6]	Total ANSIs[7] Evaluated (Unevaluated)	Total Other Designations[8] (PWA, NWA, IBP, Ramsar) Evaluated (Unevaluated)
					Plant	Bird	Reptile / Amphibian	Fish	Lepidopteran			
Lake Superior	11	3	8	1,080.6[3]	5 (5.1)	5 (20.8)	0 (0)	0 (0)	0 (0)	13	(2)	(1)
St. Marys River	48	44	11	3,567.0[3]	1 (1.0)	2 (8.3)	0 (0)	0 (0)	0 (0)	54	(1)	
Lake Huron	7			7,459.0[3]	48 (48.5)	12 (50)	6 (54.5)	5 (50)	2 (66.7)	97	22 (29)	17 (6)
St. Clair River	4	4	1	87.2	5 (5.1)	1 (4.2)	3 (27.3)	2 (20)	0 (0)			
Lake St. Clair	5	5	0	2,522.9[3]	19 (19.2)	12 (50)	4 (36.4)	6 (60)	1 (33.3)	1	1 (1)	2
Detroit River	5	4	0	1,136.3	17 (17.2)	3 (12.5)	5 (45.5)	2 (20)	2 (66.7)		1	
Lake Erie	30	26	4	19,329.8	66 (66.7)	16 (66.7)	11 (100)	7 (70)	3 (100)	3	14 (2)	7 (2)
Niagara River	82	68	18	84.5	9 (9.1)	1 (4.2)	0 (0)	3 (30)	0 (0)		1	2
Lake Ontario	4			11,334.5[3]	20 (20.2)	11 (45.8)	3 (27.3)	3 (30)	2 (66.7)	5	30 (2)	6
St. Lawrence River	40	21	19	7,017.7[3]	7 (7.1)	5 (20.8)	1 (9.1)	3 (30)	1 (33.3)		11	6
Total	236	175	61	53,619.5	197	68	33	30	11	173	79 (37)	57 (8)

[1] PSW = Provincially Significant Wetland
[2] NPSW = Non-Provincially Significant Wetland
[3] Evaluations incomplete; see also Table 3
[4] NHIC rank S1 to S3S4; see also Appendix D
[5] From Appendices P to U
[6] From Natural Areas, Environmental Sensitivity Atlases, and Other
[7] ANSI= Area of Natural and Scientific Interest
[8] PWA= Provincial Wildlife Area; NWA= National Wildlife Area; IBP= International Biological Program Site; Ramsar = Wetland of International Importance.
Note: An individual wetland may have more than one designation; see also Table 1

5.0 Results

5.1 Lake Superior

5.1.1 Setting

Lake Superior is the largest and deepest of the Great Lakes. Its surface area is 82,100 km^2 and the shoreline extends 5,105 km (International Joint Commission, 1993). St. Ignace Island and Michipicoten Island are the largest of the many islands in Lake Superior. Its drainage basin is 138,586 km^2 in size and is mostly forested (95%) with very small areas of agricultural land (1%), urban and industrial area (0.1%) and other land uses (4%) (PLUARG, 1978).

The north shore of Lake Superior is a high-energy environment with few areas of sediment deposition. Wetlands are rare here and restricted to the large sheltered embayments of Goulais Bay and Batchawana Bay in the northeast, and Thunder Bay, Black Bay and Nipigon Bay in the northwest. Due to their rarity, those that do exist are particularly important to fish and wildlife populations. Small lacustrine marshes are the most dominant wetland type, however there are also some large swamps and peatlands (e.g. Black Bay and Goulais River). The number and area of coastal wetlands along the north shore are not known. There are at least 21 coastal wetlands (Environment Canada, 1993a), but only 11 of these have been evaluated by the OMNR, mostly along the northwestern shoreline (Appendix E; Figure 1). They total approximately 1,081 ha and range in size from just under 5 ha to 429 ha (Appendix E). At least 3,500 ha of coastal wetland have not been evaluated, including several large wetlands such as the Goulais River Wetland complex and the Black Bay Peatlands (K. Cullis, pers. com., 1996; Table 3; Appendix F). The large Goulais River Wetland complex is probably the most significant wetland on the north shore of Lake Superior.

5.1.2 Significant Features

Several provincially significant plant species are found in Lake Superior coastal wetlands such as Wiegand's Sedge and Large Water Starwort in shallow water marshes, and Black Sedge in swamps. Other significant plant species found in Lake Superior coastal wetlands are Ram's-head Lady's Slipper and Water Awlwort (Appendix P).

Forty-one fish species have been identified in the coastal wetlands of Lake Superior. These wetlands provide spawning, nursery and feeding habitat for fish (Ball and Tost, 1992; Jude and Pappas, 1992; Entwistle, 1986). Northern Pike, Walleye and Yellow Perch are the primary sportfish that use Lake Superior coastal wetlands. The only Muskellunge populations along the north shore are found at the large wetlands at the mouth of the Goulais River (OMNR, 1991). No provincially significant fish species have been reported for north shore Lake Superior coastal wetlands (Sutherland, 1994; Mandrak and Crossman, 1992; Appendix Q).

Little is known about the amphibian and reptile populations utilizing Lake Superior coastal wetlands. Snapping Turtles and Bullfrogs are present in all Thunder Bay marshes (Entwistle, 1986). No provincially significant amphibian or reptile species were reported for north shore Lake Superior coastal wetlands (Appendix R).

Lake Superior coastal wetlands provide important breeding and migratory habitat for waterfowl. Along the north shore of the lake, many migrating waterfowl use the wetlands on the shores of Batchawana Island and the Goulais River Wetland complex in the southeast, and Mission Marsh in Thunder Bay in the northwest (Smith, 1987). Tens of thousands of waterfowl and other water birds pass through the Thunder Bay Harbour marshes during spring and fall migration (Entwistle, 1986). The most abundant migratory waterfowl include Canada Geese, scaup, other diving ducks, and dabbling ducks.

Colonial birds such as the Great Blue Heron, Double-crested Cormorant and American Bittern are common summer residents in these coastal wetlands and breed in Thunder Bay marshes (Entwistle, 1986). Large numbers of Great Blue Heron utilize the wetlands on the shores of Batchawana Island (Environment Canada, 1993a) and the Sandhill Crane breeds in the Goulais River Wetland complex in Goulais Bay (Environment Canada, 1993a). Provincially significant bird species reported in Lake Superior coastal wetlands during breeding season include Bald Eagle and Peregrine Falcon, while the Yellow-headed Blackbird use these areas during migration (Appendix S). Other provincially significant species breeding or using these wetlands include Northern Shoveler and Red-necked Grebe (Appendix S).

No provincially significant lepidopteran species were reported for north shore Lake Superior coastal wetlands (Appendix T).

5.1.3 Wetland Status

There are no comprehensive estimates of coastal wetland losses for Lake Superior. Along the north shore, large-scale losses have not occurred because the shoreline is remote and sparsely populated. The only reported wetland loss has occurred in Northern Wood Preservers Marsh in Thunder Bay Harbour as a result of shoreline modification and urban encroachment (K. Bray, pers. com., 1996). None of the wetlands outside the city of Thunder Bay in western Lake Superior have suffered significant wetland loss (S. Suke, pers. com., 1996).

Coastal wetlands along Lake Superior are comparatively less affected by human stressors than those of the other Great Lakes. Water level regulation is the most widespread stressor, affecting all coastal wetlands in Lake Superior. Water levels on Lake Superior have been regulated for much of the 20th century as a result of the locks at Sault Ste. Marie; thereby restricting the natural range of water levels as compared to Lake Huron and Lake Erie. Lake Superior's range of fluctuations and the cyclic nature of high and low lake levels have not been altered as significantly as Lake Ontario. However, extreme low water conditions during summer are not frequent enough to allow cyclic, regenerative processes to occur (Wilcox *et al.*, 1993).

Other, more site-specific stressors of some Lake Superior coastal wetlands are nutrient enrichment and toxic contamination of waters and sediments. Four Areas of Concern (AOC) are located along the Canadian shore but only the Thunder Bay AOC has significant areas of coastal wetland which are potentially affected. Adjacent industrial land use in the harbour has been reported to stress these wetlands (Appendix E).

Along the northwest shore, other site specific stresses on coastal wetlands include recreational use, cottaging and associated roads, especially in Cloud Bay, Pine Bay and Sturgeon Bay to the west of Thunder Bay; these may result in incremental loss of wetland area (Appendix E). Similarly, ongoing recreational use and cottage development along the northeast shore continually exert pressure on coastal wetlands (Appendix E).

5.2 St. Marys River

5.2.1 Setting

The St. Marys River extends 112 km, draining Lake Superior into Lake Huron. It drops 6.7 m along its length, mostly at the 1.2 km long St. Marys Rapids in Sault Ste. Marie (Duffy *et al.*, 1987). Several islands occur in the river, including Sugar Island, Neebish Island, St. Joseph Island and Drummond Island. The river flows through several channels around these islands and through several large lakes including Lake Nicolet, Lake George and Munuscong Lake. Its shoreline stretches 292 km on the Canadian side and 390 km on the U.S. side (International Joint Commission, 1993). The river itself has several tributaries, draining a watershed of 2,830 km^2, yet the water entering from these tributaries is only a small fraction of the drainage from Lake Superior (Kauss, 1991). Most of this watershed is forested (95%) (Kauss, 1991); the small urban and industrial areas are concentrated in Sault Ste. Marie, Ontario and Sault Ste. Marie, Michigan.

The upper river above the St. Marys Rapids has sandy and rocky shores, with emergent wetlands occurring only in sheltered areas (Duffy *et al.*, 1987). The lower river is bordered by extensive emergent marshes in the shallow areas of large lakes, bays and islands (Kauss, 1991; Duffy *et al.*, 1987). These are exposed to the river and often grade inland to palustrine wetlands, mostly marshes and swamps. Extensive emergent wetlands are found, especially along the shores of Lake George and St. Joseph Island (Environment Canada, 1994e).

Seven wetlands have been evaluated by OMNR, totalling 3,567 ha and ranging from 42 ha to 2,275 ha in size (Figure 2; Appendix G), but at least 130 other wetlands greater than 2 ha have not yet been evaluated in the lower river (Table 3). The largest evaluated wetlands include Hay Marsh along the southwest shore of St. Joseph Island (2,275 ha) and Echo Bay in Lake St. George (710 ha) (Appendix G).

5.2.2 Significant Features

The emergent wetland areas of the St. Marys River serve as spawning, nursery and feeding habitat for 44 fish species (Duffy *et al.*, 1987). Sportfish such as Northern Pike, Muskellunge, Smallmouth Bass, Largemouth Bass, Yellow Perch, and Walleye are highly dependent on these marshes (Bray, 1993). No provincially significant fish were reported for St. Marys River coastal wetlands (Appendix Q).

The St. Marys River wetlands have been identified as a significant area of waterfowl production in the Great Lakes basin (Prince *et al.*, 1992). Common breeding species include Canada Goose, Mallard, Blue-winged Teal, Black Duck and Common Merganser (Duffy *et al.*, 1987). The wetlands are also important migratory staging areas, especially for diving ducks such as Ring-necked Duck, Redhead and scaup species (Prince *et al.*, 1992; Duffy *et al.*, 1987). Provincially significant bird species breeding in these wetlands include Bald Eagle and Black Tern (Appendix S).

One provincially significant plant species, Water Awlwort was reported in St. Marys River coastal wetlands (Appendix P). No provincially significant lepidopteran species, reptile species nor amphibian species were reported for St. Marys River coastal wetlands (Appendix T; Appendix R). However, not all evaluation data on significant species were available for analysis for this area.

5.2.3 Wetland Status

Historically, wetlands along the St. Marys River shoreline have not suffered significant loss due to human influence (Bray, 1992; Williams and Lyon, 1991). The extent of human stressors affecting St. Marys River coastal wetlands is not clearly understood, but these wetlands appear to be less impacted than other connecting channels downstream. There has been site-specific loss of wetland area along the shoreline of the city of Sault Ste. Marie from dredging, filling, and sediment contamination (Bray, 1992). More recently, most of the evaluated wetlands on the Canadian side have also suffered some loss, primarily from shoreline modification, dredging, filling, channelization and cottage development (S. Jones, pers. com., 1996; Appendix G).

The entire river has been declared an Area of Concern due to elevated concentrations of contaminants in the water, localized sediment contamination, the presence of fish tumours, localized impairment of the benthos and localized high bacterial counts (Hartig and Thomas, 1988). These impacts are especially heavy along the Canadian shore, downstream of Sault Ste. Marie, Ontario to Little Lake St. George (Burt *et al.*, 1991; Kauss, 1991; Nichols *et al.*, 1991). Local wetlands in these areas are likely stressed to some degree from contaminants in the sediments, but the extent of these impacts is not clear.

Commercial shipping continues to stress wetlands in the St. Marys River. The passing of large commercial vessels in the shipping channels causes increased current speed, greater wave action, more erosion and turbidity in these coastal wetlands, affecting plant rooting and growth and associated invertebrates and other fauna (Kauss, 1991). Dredging of the river also affects the currents and sediment deposition patterns in the river, but the impacts on the sediment supply to coastal wetlands and their functions are not clear. Tributaries to the St. Marys River can also produce excessive turbidity in nearshore areas during major runoff events as a result of the fine clay soils in their watersheds, especially in Munuscong Lake in the lower river (Kauss, 1991). The excessive turbidity negatively impacts coastal wetlands by reducing water clarity, plant growth and faunal interactions.

Cottaging also produces site-specific stresses on coastal wetlands. These stresses are associated with hardening of the shoreline and dredging and channelization for boat slips and marinas (A. Dupont, pers. com., 1996).

5.3 Lake Huron

5.3.1 Setting

Lake Huron is the second largest Great Lake. Its surface area is 59,500 km^2, with a shoreline that extends 6,373 km (International Joint Commission, 1993, 1989). It has many islands ranging from large ones such as Manitoulin Island (the largest freshwater island in the world) to the many small islets of eastern Georgian Bay. Its drainage basin is 128,863 km^2 and is predominantly forested (66%), especially on the Canadian side, with lesser amounts of agricultural land (22%), residential and industrial land (10%) and other land uses (2%) (PLUARG, 1978).

Wetlands along the Canadian shore of Lake Huron are common in the sheltered environments of embayments and creek mouths and in the lee of large islands (Environment Canada, 1994b). However, an accurate estimate of wetland area along the Canadian shore of Lake Huron has not been determined. Forty-eight wetlands have been evaluated by the OMNR in Lake Huron totalling approximately 7,459 ha and ranging in size from 5 ha to 807 ha (Figure 2; Appendix G). More than 100 wetlands greater than 2 ha in size still need to be evaluated, especially in the North Channel, along Manitoulin Island and in Georgian Bay (Table 3; Appendix H). Bookhout *et al.* (1989) estimated that 12,600 ha of wetlands occurred in Georgian Bay alone. Evaluated wetlands of Lake Huron are generally smaller but more numerous than those in the southern Great Lakes and over half are wetland complexes. Marshes and swamps are equally dominant, and many have significant fen components (Figure 2; Appendix G).

Only six small coastal wetlands occur between Sarnia and Point Clark along the southeast Canadian shore, which reflect the high-energy shoreline environments. They are predominantly swamps, with a total area of approximately 340 ha (Figure 2; Appendix G). From Point Clark to the base of the Bruce Peninsula, the shoreline is mostly exposed, but 5 wetlands totalling approximately 913 ha are found in sheltered bays. They are primarily palustrine swamp and fen wetland complexes extending back from the shore (Figure 2; Appendix G).

The western shore of the Bruce Peninsula and southern shore of Manitoulin Island have exposed irregular shorelines, with wide and shallow, boulder-strewn, limestone bedrock shelves, and many small islands, reefs, and sheltered bays. The irregular coast and islands provide many sheltered, low energy and bay environments where wetlands can develop (Environment Canada, 1994b). Thirteen wetlands have been evaluated on the western side of the Bruce Peninsula, primarily large wetland complexes with swamp, marsh and fen components, totalling approximately 1,765 ha (Figure 2; Appendix G). At least 17 unevaluated wetlands occur on the southern shore of Manitoulin Island which appear to possess similar characteristics to those of the western side of the Bruce Peninsula (Natural Heritage Information Centre, 1995; Environment Canada, 1994b; Figure 2; Appendix H).

The eastern shoreline of the Bruce Peninsula in Georgian Bay is rugged with steep nearshore slopes that, prevent the development of extensive wetlands. The long, mostly sandy shore of Nottawasaga Bay also lacks wetlands except in interdunal areas, a few harbours, and river mouths. Southern Georgian Bay is rocky, but a number of shoreline marshes have developed where sheltered embayments occur (e.g., Matchedash Bay) (Figure 2; Appendix G). Twenty-one wetlands totalling approximately 3,871 ha have been evaluated between Tobermory and the French River (Figure 2; Appendix G). They are primarily lacustrine wetlands and palustrine marshes with large swamp components. A few of these wetlands also have minor bog and fen components.

The shoreline of the North Channel and northern Georgian Bay is extremely complex with bedrock outcrops, islands and bays. The mainland coast is very sheltered from wind and wave action due to numerous islands, headlands and embayments (Environment Canada, 1994b). Wetlands develop in the protected embayments of the islands and the mainland. There are at least 60 wetlands in this area, but only three have been evaluated: Marsh Bay Island 9 Wetland near Blind River (254 ha), Spanish River Delta Marsh (305 ha), and Whites Cove (11 ha) (Figure 2; Appendix G). Unevaluated wetlands occur primarily in the protected bays of St. Joseph's Channel, along the north shore of Manitoulin Island, near Barrie Island, Strawberry Island, La Cloche Islands and in McGregor Bay (Environment Canada, 1994b).

5.3.2 Significant Features

Wetlands in Lake Huron have more complex vegetation communities than those in the southern Great Lakes. The large amount of fen and swamp habitat, the diversity of wetland types, the variations in geomorphology and the calcareous soils all contribute to this complexity (Smith *et al.*, 1991). The fens which commonly occur in Lake Huron and Georgian Bay wetlands, also known as coastal meadow marshes, have been identified as globally imperilled communities (Natural Heritage Information Centre, 1995). Over 40 species of provincially significant plants have been found in coastal wetlands of Lake Huron (Wilcox, 1995; Appendix P). For example, the coastal meadow marshes of Lake Huron and Georgian Bay support some of Ontario's rarest plant species, including Bluehearts, Round-stemmed Purple False Foxglove, Twining Bartonia and Rigid Yellow Flax (Appendix P). Of those significant plant species that are found in coastal wetlands in Ontario, one species, Bluehearts, has only been reported for Lake Huron (Appendix P).

Fifty-nine fish species utilize the coastal wetlands of Lake Huron (Severn Sound Remedial Action Plan, 1993a; Prince *et al.,* 1992). Over half are permanent residents while the remainder use them on a temporary basis for feeding, shelter, spawning, nursery, and dispersal of young. Largemouth Bass, Rock Bass, Bluntnose Minnow, Pumpkinseed and Banded Killifish are the most common permanent residents (Severn Sound Remedial Action Plan, 1993a). Sportfish such as Northern Pike, Walleye, Muskellunge and Smallmouth Bass also depend on these wetlands along with many species of bait fish. The five provincially significant fish species that use these coastal wetlands are Pugnose Shiner, Lake Chubsucker, Grass Pickerel, Black Bullhead and Longear Sunfish (Appendix Q).

Lake Huron wetlands also provide important habitat for amphibians and reptiles. The amphibians use them for breeding, nursery and feeding. Reptiles nest on uplands, but many species spend the remainder of their life cycle in these wetlands. Five provincially significant reptile species have been found in the coastal wetlands of Lake Huron and Georgian Bay: Spotted Turtle, Eastern Spiny Softshell Turtle, Queen Snake, Eastern Fox Snake and Eastern Massasauga Rattlesnake (Appendix R). One provincially significant amphibian species, Jefferson Salamander, has been reported in the coastal wetlands of Lake Huron and Georgian Bay (Appendix R).

Prince *et al.* (1992) identified the marshes of Georgian Bay as significant areas of waterfowl production for the Great Lakes. At least 2,100 pairs of dabbling ducks nest in Georgian Bay wetlands. The wetlands of Lake Huron are also important during migration for provincially significant species such as the Red-necked Grebe, Northern Shoveler, and Redhead (Appendix S). In terms of coastal wetland use for breeding in Lake Huron and Georgian Bay, 12 provincially significant species have been reported, including Great Egret, Redhead, Ruddy Duck, and Little Gull. See Appendix S for complete listings.

Two provincially significant lepidopteran species, Mulberry Wing and Two-spotted Skipper, have been reported in Lake Huron coastal wetlands (Appendix T).

Lake Huron wetlands provide habitat for many fur-bearing mammals including Mink, Beaver, River Otter, Raccoon, Red Fox and Muskrat. In south and central Lake Huron, the coastal swamps provide significant winter cover for White-tailed Deer. Moose occur in the wetlands along the north shore of Georgian Bay.

5.3.3 Wetland Status

No comprehensive estimates of coastal wetland loss are available for the Canadian shore of Lake Huron. Loss of wetland habitat on a large scale has not occurred because most of the shoreline is sparsely populated and remote. Most loss tends to be concentrated around the small urban centres that dot the shore. Within the last 15 years, there has been incremental and site-specific loss of wetland area from agricultural encroachment and cottage development. More than half of the wetlands along the central coast, the western coast of the Bruce Peninsula and southern Georgian Bay have suffered recent loss of acreage (Appendix G). A study of wetland loss in Severn Sound in southern Georgian Bay indicated that wetland habitats have decreased by 68% and 18% in Penetanguishene and Hog Bay respectively since 1951 (Severn Sound Remedial Action Plan, 1993b). The main causes of these wetland losses are shoreline modification, road construction, filling for urban and cottage development and dredging and channelization associated with marina development (Severn Sound Remedial Action Plan, 1993a). The wetlands along the shoreline of the North Channel and northern Georgian Bay have only suffered small losses, however cottaging, and marina and subdivision development continually put pressure on these coastal wetlands through dredging and modification of the shoreline (W. Selinger, pers. com., 1996). Along the rest of the Canadian shore, remaining wetlands do not appear to be as impacted by human stresses as compared to the southern Great Lakes. Stresses appear to be site-specific and localized.

In addition to outright wetland loss, urban encroachment, cottaging and marinas cause multiple stresses on remnant wetlands. On the western Bruce Peninsula and southern Georgian Bay, these stressors include shoreline modification, road crossings, dredging and channelization. Shoreline modification prevents the landward migration of remnant wetlands during high water periods. Road crossings alter wetland hydrology and, along with dredging, filling and channelization, fragment the remaining wetland habitat.

Wetlands in the bays of southeastern Georgian Bay are affected by nutrient and sediment loading from watersheds. Excessive phosphate inputs to these bays originate from point and non-point sources associated with urban areas and agriculture (Severn Sound Remedial Action Plan, 1993a). Excess sediment loadings originate mostly from non-point source inputs, mainly agricultural runoff (Severn Sound Remedial Action Plan, 1993a).

Non-indigenous species including Purple Loosestrife, Common Carp and Zebra Mussels are also localized stressors in Georgian Bay. Purple Loosestrife is especially a problem in southern Georgian Bay (G. Allen, pers. com., 1996).

5.4 St. Clair River

5.4.1 Setting

The St. Clair River drains Lake Huron into Lake St. Clair. It forms a large bird-foot delta with many distribution channels and wetlands where it meets the lake. This delta is a transitional environment between the river and the lake. For the sake of clarity in the Atlas, the delta and its coastal wetlands are covered under the assessment of wetlands of Lake St. Clair. The first 43 km of the St. Clair River are considered here, between Lake Huron at Sarnia-Port Huron and the first distribution channel of the delta, Chenal Ecarte across from Algonac.

The river above the delta is a uniform channel with very few bends or meanders, no cut-off channels nor oxbow lakes and only two islands, Stag Island and Fawn Island (Edsall *et al.*, 1988). The river drops only 1.4 m between Lake Huron and the beginning of the delta at Chenal Ecarte, but it has relatively high flows, with an average flow velocity reaching 3.2 km/hr. The natural shoreline has a bank 1.5 to 5 m high (International Joint Commission, 1989), but most of this shoreline is now artificial (Ontario Ministry of the Environment and Michigan Department of Natural Resources, 1991).

The overwhelming majority of the flow in the river comes from Lake Huron. Including the delta, the drainage basin of the river is 3,368 km^2 and is mostly agricultural (69%). The urban areas are concentrated in a narrow zone along the river with the larger centres being Sarnia, Ontario and Port Huron, Michigan. Much of the industry is concentrated in Ontario in the first 14 km of the river between Sarnia and Corunna, a stretch of shoreline known as "Chemical Valley".

The lack of shoreline complexity along with the fast current, the depth of the river and wave forces generated by the passage of large commercial vessels limits wetland development along the banks of the river (Bookhout *et al.*, 1989; Edsall *et al.*, 1988). Wetlands occur primarily on the shallow submerged shoals of the river and tributary channels and consist mostly of submergent macrophyte beds (Griffiths *et al.*, 1991; Edsall *et al.*, 1988).

There is no clear estimate of the area of coastal wetland in the St. Clair River above the delta. Edsall *et al.*, (1988) identified 550 ha of coastal wetlands in the entire St. Clair River based on navigation charts and topographic maps, but did not provide the location and size of individual wetlands. Bookhout *et al.* (1989) identified 96 ha of coastal wetland along the Ontario shore. Extensive submergent macrophyte beds are known to occur in Sarnia Bay, around Stag and Fawn Islands and along the Canadian shoreline near these islands (Griffiths *et al.*, 1991). OMNR has evaluated five wetlands for the St. Clair River totalling 87 ha. In Ontario, just over 13 ha of wetlands have been evaluated along the river itself, namely the Stag Island Marsh and the Point Edward Marsh. These estimates indicate that coastal wetlands are uncommon habitats in the St. Clair River above the delta (Figure 3; Appendix I). The remaining wetlands are therefore particularly important habitats for plants, fish and wildlife in the river.

5.4.2 Significant Features

Coastal wetlands of the St. Clair River are primarily composed of submergent species, but emergent macrophytes also occur (Griffiths *et al.*, 1991; Edsall *et al.*, 1988). Five provincially significant plant species have been found in the shallow water and meadow marshes: Angled Spike-rush, Winged Loosestrife, Swamp Rose Mallow, Riddell's Goldenrod and Walter's Barnyard Grass (Appendix P).

Unlike Lake St. Clair and the delta of the river, the wetlands of the river above the delta are not important breeding sites or migration corridors for waterfowl (Bookhout *et al.*, 1989). Waterfowl overwinter in wetlands in the river when nearby marshes with less current are frozen since thermal pollution and current keep the river open. Common species are Common Merganser, Redhead, Canvasback, American Widgeon, Mallard and scaup. One provincially significant bird species, Forster's Tern, has been reported to breed in the wetlands of the St. Clair River (Appendix S).

Forty-five species of fish have been recorded using the wetlands of the St. Clair River (Mandrak and Crossman, 1992; Edsall *et al.* 1988). Over half of these are permanent residents, with Brown Bullhead, Common Carp and White Perch being the most abundant. Other species use the wetlands on a temporary basis for spawning, nursery and feeding; the most abundant species include the White Sucker, Alewife, Rainbow Smelt, Gizzard Shad and Rock Bass. Several important sport fish also use the wetlands on a temporary basis including Northern Pike, Muskellunge, Walleye, Yellow Perch, Smallmouth Bass and Largemouth Bass (Edsall *et al.*, 1988). Two provincially significant fish species, the Pugnose Minnow and Black Bullhead, have been reported in St. Clair River coastal wetlands (Appendix Q).

Many species of amphibians and reptiles, including salamanders, frogs, toads, snakes, lizards and turtles, also occur in these wetland habitats (Edsall *et al.* 1988). Three provincially significant reptiles inhabit St. Clair River wetlands, namely the Eastern Spiny Softshell Turtle, Butler's Garter Snake and the Eastern Fox Snake (Appendix R).

No provincially significant lepidopteran species were reported for St. Clair River coastal wetlands (Appendix T).

Mammals commonly found in coastal wetlands of the St. Clair River include the Virginia Opossum, Eastern Cottontail Rabbit, Muskrat, Striped Skunk and White-tailed Deer (Edsall *et al.*, 1988).

5.4.3 Wetland Status

Some wetland loss appears to have occurred along the shores of the St. Clair River above the delta, but there is no comprehensive estimate of the extent of this loss. In Ontario, there are known wetland losses of 3 ha at Point Edward Marsh and 40 ha at Stag Island Marsh.

Almost all of the U.S. shoreline and most of the Canadian shoreline have been extensively modified and consist of residential, recreational and industrial developments (Edsall *et al.*, 1988). The river is also an important port. As such, wetland loss in the river appears to be largely related to extensive bulkheading, shoreline hardening, filling, channelization and dredging along the shores of the river. These activities also fragment the few remaining wetlands along the river. Urban encroachment continues to cause wetland loss and impairment on the Canadian side (D. Hector, pers. com., 1996).

Several other human activities stress remnant wetlands along the river. The river is a busy seaway and port. Ship wakes from large commercial vessels are a significant stressor to shoreline habitats, including remnant coastal wetlands, eroding the shoreline and hampering the establishment of aquatic macrophytes.

The St. Clair River was declared an Area of Concern (AOC) as a result of the excessive levels of toxic substances in the water, contaminated sediments, impaired benthos and bacterial contamination (Hartig and Thomas, 1988). Industry is the main source, but municipal sewage treatment plants, other point and non-point sources of pollution are also concerns (St. Clair River Remedial Action Plan, 1995). Remnant wetlands are particularly vulnerable to contamination by toxic chemicals since they are located in sediment accumulation zones along the shore.

5.5 Lake St. Clair

5.5.1 Setting

Lake St. Clair is a shallow productive lake located between the St. Clair River and the Detroit River. Where the St. Clair River meets Lake St. Clair, an expansive bird-foot delta has formed which has many distribution channels, islands and wetlands. This delta is included in the assessment of the coastal wetlands of Lake St. Clair.

Lake St. Clair is heart-shaped with a surface area of 1,115 km^2 (Edsall *et al.*, 1988). The basin is very shallow and has a maximum natural depth of 6.5 m, although a commercial shipping channel has been dug across the lake to a depth of 8.5 m. The shoreline, excluding the distributary channels in the delta, extends 272 km (Edsall *et al.*, 1988). Several large tributaries flow into the lake including the Sydenham River and the Thames River in Ontario and the Clinton River in the United States. Together they drain a watershed of 12,616 km^2, in which agricultural land uses predominate (Edsall *et al.*, 1988). These tributaries contribute 2% of the flow to the lake; the remainder flows from the St. Clair River.

The St. Clair River enters the lake in the northeast forming a large delta. The topography is very flat and the river drops 0.2 m over 17 km through the delta (Edsall *et al.*, 1988). There are numerous distribution channels, but the majority of the flow (92%) passes through the western half of the delta through the North, Middle and South Channels. This delta is the product of long-term geological and riverine processes; its development is linked to the deposition of sediments as the river slows to meet the lake. Sediments that continue to feed this delta originate from the nearshore areas of southern Lake Huron (Edsall *et al.*, 1988).

Lake St. Clair and the St. Clair Delta contain some of the largest coastal wetlands in the Great Lakes. There are many estimates of the aerial extent of these wetlands. However, since the topography surrounding much of the lake and especially in the delta is almost flat, water level fluctuations greatly affect their extent and position. Large changes in wetland area are especially great between years of high and low water levels (Herdendorf *et al.*, 1986). While these changes are important for the diversity of habitat, they make it difficult to compare different estimates of wetland extent in the lake.

The largest wetlands in the lake are found in the St. Clair Delta, which harbours a vast complex of lacustrine, riverine and palustrine wetlands. On the Canadian side of the St. Clair Delta, there are at least 12,769 ha of coastal wetlands (D. Hector, pers. com., 1996). The majority of wetlands, approximately 10,360 ha are on Walpole Island First Nation and have not been evaluated (Appendix J). Five other wetlands, totalling approximately 2,522 ha, have been evaluated on the Canadian side of the delta ranging from just under 4 ha to 2,335 ha, the largest being the St. Clair Marshes Complex in Mitchell's Bay (2,335 ha) (Figure 3; Appendix I). Of these wetlands, approximately one third of the area has been diked for intensive waterfowl management (Bookhout *et al.*, 1989).

Beyond the delta, remnant lacustrine marshes occur primarily near mouths of rivers and creeks along the northern and eastern shores of the lake. Very few wetlands occur along the highly developed southern and western shores. In Ontario, three wetlands are located along the eastern shore totalling 159 ha and ranging from 4 ha to 131 ha; the largest occurs at the mouth of the Thames River (131 ha) (Figure 3; Appendix I). Many of these wetlands are diked (Herdendorf *et al.*, 1986). Only one small wetland, Ruscom Shores Marsh (29 ha), is found on the southern shore.

5.5.2 Significant Features

Herdendorf *et al.* (1986) identified twelve different wetland habitats in Lake St. Clair and the delta, each with different vegetation and environmental characteristics. As a result of this range of habitats and the size of the wetlands, Lake St. Clair has some of the most diverse wetlands in the Great Lakes for plants, fish and wildlife. These wetlands provide habitat for many common species, but also provide some of the most important habitat for rare flora and fauna.

Nineteen provincially significant plant species have been found in the coastal wetlands of Lake St. Clair (Appendix P). These include the very rare Angled Spike-rush, as well as Southern Tickseed, Honey Locust, American Lotus, Many-fruited False Loosestrife, Round-stemmed Purple False Foxglove, Pale Purple False Foxglove, Hairy Fimbristylis and the globally rare Eastern Prairie White-fringed Orchid. Of those significant plant species that are found in coastal wetlands in Ontario, one species, Hairy Fimbristylis has only been reported for Lake St. Clair (Appendix P).

Lake St. Clair marshes provide habitat for more than 65 species of fish (Mandrak and Crossman, 1992; Edsall *et al.*, 1988). Two-thirds of the fish species are permanent residents; the most common species are Rock Bass, Bluegill, Black Bullhead, Yellow Bullhead, Channel Catfish, Alewife, White Perch and Common Carp. Other fish use the wetlands on a temporary basis for spawning, nursery, shelter or feeding; common species include White Sucker, and Rainbow Smelt. Several sportfish also commonly use these wetlands, including Northern Pike, Muskellunge, Walleye, Yellow Perch, Smallmouth Bass, Crappie and sunfish species (Jude and Pappas, 1992; Edsall *et al.*, 1988). Lake St. Clair is one of only two sites with large Muskellunge populations in the Great Lakes (Edsall *et al.*, 1988). The only large spawning area left for Muskellunge in Lake St. Clair is in Anchor Bay, Michigan, while the shallow marshes of the delta are the only known nursery areas for Muskellunge in the entire St. Clair River, Lake St. Clair and Detroit River system (Edsall *et al.*, 1988). Six provincially significant fish species, Pugnose Shiner, Pugnose Minnow, Lake Chubsucker, Grass Pickerel, Black Bullhead, and Longear Sunfish, have been reported in Lake St. Clair coastal wetlands (Appendix Q).

Many species of amphibians and reptiles also occur in the wetland habitats of Lake St. Clair, including salamanders, frogs, toads, snakes, lizards and turtles (Edsall *et al.*, 1988). Four provincially significant reptile species inhabit Lake St. Clair coastal wetlands: Eastern Fox Snake, Queen Snake, Spotted Turtle and Eastern Spiny Softshell Turtle (Appendix R).

The St. Clair Delta has been identified as one of the most significant areas for waterfowl production, staging and migration in the Great Lakes (Prince *et al.*, 1992; Bookhout *et al.*, 1989). Approximately 16% of all Great Lakes coastal wetlands of importance to waterfowl are found in the St. Clair Delta (Prince *et al.*, 1992). In terms of breeding waterfowl, the highest densities of Mallard, Black Duck, Blue-winged Teal and Green-winged Teal in the Great Lakes basin are found in the wetlands of the St. Clair Delta. Redhead is the only species of diving duck that breeds regularly in the Great Lakes, and the St. Clair Delta wetlands produce up to 4,000 Redheads annually (Prince *et al.*, 1992). Delta wetlands also provide nesting habitat for Canada Geese and Ruddy Duck. Furthermore, the St. Clair Delta lies in major migration corridors of both dabbling and diving ducks (Bookhout *et al.*, 1989). They provide one of the most important staging and feeding grounds for post-breeding and migratory Canada Geese, Tundra Swans and dabbling ducks on the Great Lakes. The delta is also one of the major fall staging areas in North America for Canvasback and Redhead (Prince *et al.*, 1992).

Lake St. Clair coastal wetlands provide habitat for many other species of birds. American Coot, Herring Gull, Common Tern, Red-tailed Hawk and Northern Harrier are commonly observed in Lake St. Clair marshes (Herdendorf, 1992). Waterbirds such as Great Blue Heron, American Bittern, Least Bittern, King

Rail, Great Egret and Black-crowned Night Heron breed here. Large nesting colonies of Great Blue Heron and Black-crowned Night-Heron occur in the marshes of Walpole Island (Smith *et al.*, 1991). Walpole Island marshes also support the largest number of nesting pairs of Forster's Tern on the Great Lakes and provide nesting habitat for the Black Tern (Smith *et al.*, 1991). Eleven provincially significant bird species breed in the coastal wetlands of Lake St. Clair. Some of the more notable species include Great Egret, Canvasback, Least Bittern, King Rail, Ruddy Duck and Redhead (Appendix S). See Appendix S for complete listing.

One provincially significant lepidopteran species, Duke's Skipper, has been reported in Lake St. Clair coastal wetlands (Appendix T).

More than a dozen species of mammals use the Lake St. Clair wetlands. The Virginia Opossum, Eastern Cottontail Rabbit, Muskrat, Striped Skunk and White-tailed Deer are common (Edsall *et al.*, 1988). Muskrat and Raccoon are important furbearers and are extensively trapped in the region.

5.5.3 Wetland Status

Lake St. Clair and the St. Clair delta have been extensively studied in terms of wetland loss. Overall, these wetlands were reduced by 9,139 ha or by 41% between 1868 and 1973. The most extensive losses have occurred at the mouth of the Clinton River, in the St. Clair Delta and along the eastern shore of the lake (Herdendorf *et al.*, 1986).

Along the Ontario shoreline, 4,764 ha or 34% of coastal wetlands have been lost in the delta and the lake between 1873 and 1968 (Edsall *et al.*, 1988). In 1873, the wetlands along the eastern shoreline of the lake were approximately 2.5 km wide, but by 1968, wetlands were reduced in width to approximately 0.8 km. Much of this loss was due to large scale conversion of wetlands to agricultural land. More recently, between 1965 and 1984, wetlands along the east shore of the lake, from the mouth of the Thames River to Chenal Ecarte, further dwindled by 1,064 ha (McCullough, 1985). This loss was mostly a result of agricultural drainage (89%), but some loss was due to marina and cottage development (11%). The wetlands on the Ontario side of the St. Clair Delta are intact in many places, but shoreline development, dredging and placement of dredge spoils have taken their toll. Between 1965 and 1978, 508 ha or 4.5% of the wetlands on Walpole Island were lost (McCullough, 1982). Recently, wetland loss to agricultural and urban development has continued in the lake and delta, albeit at a slower pace (D. Hector, pers. com., 1996).

Urban, recreational and agricultural encroachment have not only caused wetland loss but also stress remaining wetlands. In many cases, shoreline hardening such as breakwaters, jetties, and bulkheading restrict the landward migration of wetland communities during high water periods. This causes a backstopping effect that reduces the size and diversity of wetland communities. Recreational and urban developments also fragment the remaining wetland area.

Another major stress is the diking of wetlands. About half of the wetlands in Lake St. Clair and the St. Clair Delta have been diked (Bookhout *et al.*, 1989). They are managed mainly for waterfowl hunting at the expense of other wetland functions (Jude and Pappas 1992; Herdendorf *et al.*, 1986). Diking isolates these wetlands from the upland and lake environments, and many wetland functions are impaired. Their use by fish for spawning, nursery or feeding is impeded or cannot take place. Many of the fish species in the lake are dependent on wetlands for part of their life cycle (Edsall *et al.*, 1988). Diking also reduces organic material inputs into the lake thereby disrupting the food chain in the lake (Herdendorf *et al.*, 1986). Other stresses to these wetlands include sediment and nutrient loading from tributaries and invasive species such as Reed Canary Grass and Purple Loosestrife (D. Hector, pers. com., 1996).

5.6 Detroit River

5.6.1 Setting

The Detroit River connects Lake St. Clair to Lake Erie. It is 51 km long and drops 0.9 m along its length (Manny *et al.*, 1988). The shoreline stretches 107 km on the Canadian side and 127 km on the U.S. side (International Joint Commission, 1989). Several islands occur in the river, with the largest, Grosse Ile, near its mouth. About 95% of the total flow in the river enters from Lake St. Clair (Manny *et al.*, 1988), and the remainder flows from tributaries and sewer systems, draining a watershed of 1,844 km^2 (Manny and Kenaga, 1991). The Canadian portion of this watershed is largely agricultural (90%), and the remainder consists of urban, residential and industrial lands, centred around Windsor in the northern reaches of the river (Manny and Kenaga, 1991). Over five million people live in the Detroit River watershed. The natural shoreline consists of clay banks, but 20% of the Canadian shoreline is now artificial with revetments and other shoreline hardening structures (International Joint Commission, 1993; Manny and Kenaga, 1991).

Along the Canadian shore, four coastal wetlands have been evaluated by OMNR, primarily in the middle reaches of the river. They total approximately 1,136 ha and range in size from 32 ha to 575 ha (Figure 3; Appendix I). The Detroit River Marshes near Fighting Island represent the largest wetland complex (575 ha). One small wetland is found on Fighting Island, and the remaining wetlands are associated with tributaries entering the river, including the large Canard River Marshes wetland complex (416 ha) and the Turkey Creek Marsh (32 ha). Approximately half of the Canard River wetland is diked for intensive waterfowl management (Manny *et al.*, 1988).

5.6.2 Significant Features

Typically, coastal wetlands along the Detroit River are riverine and river-mouth marshes, sometimes with a small swamp component (Smith *et al.*, 1991). They are often dominated by submergent macrophyte communities (Manny *et al.*, 1988). Seventeen provincially significant plants have been found in these wetlands, including Seedbox, Southern Tickseed, and Honey Locust (Appendix P). See Appendix P for complete listings.

At least 45 species of fish inhabit Detroit River wetlands, 21 of which are permanent residents (Herdendorf, 1992; Mandrak and Crossman, 1992). The most abundant species are Northern Pike, Gizzard Shad, Bowfin, Common Carp, Goldfish, Carp-Goldfish hybrids, Golden Shiner, Blacknose Shiner, White Sucker, Brook Silverside, Rock Bass, Pumpkinseed, Black Crappie and Yellow Perch (Herdendorf, 1992). Other species which use these wetlands for spawning include Lake Sturgeon, Muskellunge, Channel Catfish, Largemouth Bass, Smallmouth Bass, Bluegill and Walleye (Herdendorf, 1992). Two provincially significant species of fish, the Black Bullhead and Pugnose Minnow, also use the wetlands (Appendix Q).

Many species of reptiles and amphibians inhabit Detroit River wetlands. These coastal wetlands offer particularly important habitat since the surrounding landscape has been dramatically altered. Five provincially significant species of reptiles have been identified in these wetlands: the Eastern Fox Snake, Eastern Massasauga Rattlesnake, Queen Snake, Butler's Garter Snake and the Eastern Spiny Softshell Turtle (Appendix R).

The wetlands of the Detroit River are significant for Great Lakes waterfowl production (Prince *et al.*, 1992; Bookhout *et al.*, 1989). The marshes provide important resting and feeding grounds for post-breeding and migratory Canada Geese, Tundra Swan and dabbling ducks. These birds are especially attracted to the diked wetlands of the Canard River along the Canadian shore where there are large areas of emergent wetlands with controlled water levels and adjacent agricultural fields. Many waterfowl winter in wetlands of the Detroit River because they remain open from commercial navigation and thermal pollution (Manny *et al.*, 1988). For instance, about 11,700 and 4,500 ducks wintered in the wetlands of the Detroit River in 1980 and 1981, respectively (Bookhout *et al.*, 1989). The wetlands also provide habitat for many other bird species. Provincially significant bird species reported to breed in these wetlands include Least Bittern, Northern Shoveler and White-eyed Vireo (Appendix S).

Two provincially significant lepidopteran species, Two-spotted Skipper and Mulberry Wing, have been reported in Detroit River coastal wetlands (Appendix T).

5.6.3 Wetland Status

No comprehensive estimate of the extent of wetland loss along the shores of the Detroit River exists. From depth surveys of the river in the 1870s, wetlands and large submergent macrophyte beds were nearly continuous along the shores of the river in historic times. A fringe of emergent vegetation occurred all along the shores of the river in waters 0.3 m to 2.0 m deep (Manny *et al.*, 1988). Emergent marshes extended inland from these depths and were sometimes over 1 km wide, especially near the mouths of tributaries. Today, around 87% of the U.S. shoreline of the Detroit River has been filled and bulkheaded (Manny and Kenaga, 1991), and more than 20% of the Canadian shoreline is artificial with revetments and hardened shorelines, especially in the northern sections of the river in Windsor (International Joint Commission, 1993). Consequently, many of the historic coastal wetlands have been lost through dredging, bulkheading and/or backfilling. The remaining wetlands mostly occur on islands in the river. In recent years, loss of wetland along the shores has diminished, but incremental loss from agricultural conversion, shoreline modification, marina development, and urban encroachment is still a concern (Appendix I).

Many human stressors continue to impact remaining wetlands, including erosion from shipping, shoreline modification, dredging and channelization, excess nutrients, contamination of water and sediments with toxic chemicals, agricultural and urban encroachment and invasive non-indigenous species (D. Hector, pers. com., 1996; Manny and Kenaga, 1991; Manny *et al.*, 1988).

The Detroit River is one of the busiest shipping channels in the Great Lakes. Commercial and recreational vessels cause excess wave action, changes in shoreline currents, and erosion of wetlands along the shores (Manny *et al.*, 1988). Shoreline hardening is the common solution to this erosion. Where this hardening occurs adjacent to remaining wetlands, it restricts their connection to upland habitats and prevents upslope migration during high water periods, greatly reducing the diversity of habitats.

The shipping channel is dredged each year for navigation, substantially altering the river morphology (Manny *et al.*, 1988). Sediment dynamics in the river are altered, but it is not known how these changes affect the distribution and status of wetlands. Dredging and channelization associated with the numerous smaller marinas, canals and boat slips also stress remaining wetland areas through wetland loss, fragmentation, changes in sediment dynamics, and increased erosion from wave action (D. Hector, pers. com., 1996).

The busy nature of the port, the large urban areas and the numerous industries contribute to the pollution of the river and its wetlands. The Detroit River has been identified as an Area of Concern. Excessive phosphates from combined sewers and other sources have caused the eutrophication of wetland communities which reduces plant and wildlife diversity. As well, sediments in many stretches of the river are contaminated with heavy metals, oils, and PCBs, especially along the U.S. side of the river (Manny *et al.* 1991; Nichols *et al.*, 1991). Wetlands and other nearshore habitats are especially vulnerable to toxic substances since they are deposition zones for sediments (Manny *et al.*, 1988). Submergent plants have been found to concentrate these contaminants and are used as food sources for fauna, including waterfowl (Manny *et al.*, 1991). Toxic effects and bioaccumulation are therefore important stressors to wetlands in the Detroit River.

Several exotic species are present in Detroit River wetlands that affect the composition and structure of wetland communities. Invasive plant species of concern include Eurasian Water Milfoil and Curled Pondweed (Manny *et al.*, 1988). Large populations of Common Carp are also now established in the river, and destroy submergent macrophyte beds, increase turbidity and displace native fish species (Manny *et al.*, 1988).

The diking of wetlands, such as parts of the Canard River Marshes, provides high quality habitat for waterfowl and some other fauna, but unfortunately also isolates them from the river, reducing their function in the river ecosystem (Manny *et al.*, 1988).

5.7 Lake Erie

5.7.1 Setting

Lake Erie has a surface area of 25,657 km^2 and a shoreline that extends 1,402 km (Herdendorf, 1992). Several large sand spits project into the lake, including Long Point, Turkey Point, Rondeau Peninsula, and Point Pelee, and a series of small islands occurs in the western part of the lake (Figure 4). The lake basin can be naturally divided into three sub-basins: the western basin to the west of Point Pelee, the central basin between Point Pelee and Long Point, and the eastern basin to the east of Long Point. Lake Erie is the shallowest of the Great Lakes, and is particularly subject to the effects of storms, wind tides, and seiches (Bedford, 1992).

Lake Erie, together with the St. Clair River, Lake St. Clair, and the Detroit River, has a watershed of 78,769 km^2 (PLUARG, 1978). Most of this watershed is agricultural (59%); the remaining land is forested (17%), residential or industrial (15%) or under other land uses (9%) (PLUARG, 1978). Only approximately10% of the flow entering the lake comes from tributaries, while the remainder flows from the Detroit River (Herdendorf, 1987).

Extensive coastal wetlands have developed behind the large sand spits and at river and creek mouths. In total, 30 wetlands occur covering approximately 19,330 ha, and ranging in size from just over 3 ha to 13,465 ha (Figure 4; Appendix K). Over half are wetland complexes, consisting mostly of marshes with some swamp and latitudinaly rare fen and bog components. In the western basin, there are 10 coastal wetlands covering approximately 3,033 ha (D. Hector, pers. com., 1996). The largest are barrier beach marshes at Point Pelee (1,175 ha) and drowned rivermouth marshes at Big Creek (1,000 ha), Cedar Creek (250 ha) and Hillman Creek (362 ha). Smaller wetlands are also found on Pelee Island and East Sister Island. In the central basin, five wetlands totalling 1,751 ha are found along the Canadian shoreline, with the marsh and swamp in Rondeau Provincial Park (930 ha) being the largest (D. Hector, pers. com., 1996). In the eastern basin, there are 15 wetlands along the Canadian shoreline, totalling approximately 14,543 ha (Appendix K). The most important wetlands of the eastern basin are the wet meadows, forested swamps, deep-water cattail marshes and shallow-water grass and sedge marshes and ponds protected by Long Point (Prince *et al.* 1992). These Long Point wetlands encompass 13,465 ha and include more than 70% of the total wetland area along the north shore of Lake Erie (R. Thompson, pers. com., 1996). The remaining wetlands along the eastern basin occur primarily at river and creek mouths, including the Dunnville Marsh complex (518 ha) at the mouth of the Grand River (G. Birch, pers. com., 1996).

5.7.2 Significant Features

The coastal wetlands of Lake Erie support the largest diversity of plant and wildlife species in the Great Lakes. As reported in the wetland evaluations, of all the Great Lakes and connecting channels in Ontario, the greatest number of provincially significant reptile and amphibian, vascular plant, lepidopteran, bird and fish species have been reported for Lake Erie wetlands (Appendices P to U).

The moderated climate and more southern latitude of Lake Erie allow many species not found along the northern Great Lakes to exist here. For instance, over 300 species of plants have been identified in the aquatic and wetland habitats of western Lake Erie (Herdendorf, 1992). In the open water of the lake and larger bays, submergent species predominate, including several species important to wildlife, such as Wild Celery, and Sago Pondweed (Herdendorf, 1992). Water lilies such as the White Water Lily, Yellow

Water Lily and the American Lotus are not common, but where they do grow they form extensive colonies.

The diverse coastal wetlands of Lake Erie provide habitat for many rare species of flora. Rare wetland communities such as coastal meadow marsh (fen) occur at several locations including Long Point. Sixty-one provincially significant plant species have been found in the coastal wetlands of Lake Erie (Appendix P). Some of the most provincially significant species found in these wetlands include, Scarlet Ammannia, Winged Oval Sedge, Small White Lady's Slipper, Engelmann's Spike-rush, Rigid Yellow Flax and Bayberry (Appendix P). Other examples of provincially significant plant species include Crested Arrow-head, American Lotus, Riverbank Sedge, Swamp Rose Mallow, and Eastern Prairie White Fringed Orchid (Appendix P). Of those significant plant species that are in coastal wetlands in Ontario, seven species, Winged Oval Sedge, Horsetail Spike-rush, Knee Spike-rush, Leafy Blue-flag, American Water-willow, Spotted Pondweed, and Bushy Cinquefoil have only been reported for Lake Erie (Appendix P).

Coastal wetlands of Lake Erie are important to fish production because they provide spawning and nursery habitat for many wetland dependent species, cover for juvenile and forage fish, and feeding areas for predatory fish. Many are important recreational or commercial fish species. Forty-six species of fish have been captured in Lake Erie wetlands. An additional eighteen species captured in open water are known to use wetlands during some part of their life cycle (Jude and Pappas, 1992). The most abundant permanent residents of Lake Erie coastal wetlands are White Crappie, Gizzard Shad, Black Bullhead, White Perch, White Bass, Log Perch and Freshwater Drum. Other species such as White Sucker, Common Carp, Emerald Shiner, Spottail Shiner and Yellow Perch are abundant temporary residents (Jude and Pappas, 1992). Many fish species in these wetlands are rare in the Great Lakes, including the provincially significant Spotted Gar, Lake Chubsucker, Longear Sunfish, Grass Pickerel and Warmouth (Appendix Q).

Many species of snakes, turtles, frogs and salamanders are dependent on Lake Erie wetlands. Twenty-eight species of amphibians and twenty-seven species of reptiles inhabit the Lake Erie region, most of which are found in coastal wetlands for part of their life cycle (Herdendorf, 1992). The four provincially significant amphibian species found in Lake Erie coastal wetlands are Jefferson's Salamander, Smallmouth Salamander, Fowler's Toad, and Blanchard's Cricket Frog (Herdendorf, 1992; Appendix R); the last three species are restricted in Canada to the shores of Lake Erie. Seven provincially significant reptile species have also been found: Spotted Turtle, Eastern Spiny Softshell Turtle, Queen Snake, Eastern Fox Snake, Lake Erie Water Snake, Butler's Garter Snake and Eastern Massasauga Rattlesnake (Appendix R). In Canada, the Lake Erie Water Snake is confined to the western shores of Lake Erie, and is predominantly found in coastal wetlands.

Wetlands of Lake Erie support a wide diversity of bird life. Waterfowl, wading birds, shore birds, gulls and terns, raptors and perching birds use Lake Erie wetlands for migration, nesting and feeding. The wetlands in western Lake Erie from the mouth of the Detroit River to Sandusky Bay, and those at Point Pelee, Rondeau Bay and Long Point have been identified as some of the most important waterfowl habitat complexes in the Great Lakes (Prince *et al.*, 1992). For instance, large numbers of post-breeding dabbling ducks and Canada Geese, and thousands of Tundra Swans stop annually in southwestern Lake Erie coastal wetlands (Prince *et al.*, 1992). As well, Long Point is one of the major staging areas in North America for Canvasback and Redhead. Wetlands of southwestern Lake Erie and Long Point also provide a major stop over point for diving ducks such as migrating Bufflehead, Common Goldeneye, Red-breasted Merganser, Common Mergansers and Ruddy Duck (Prince *et al.*, 1992). The provincially significant Ruddy Duck and Northern Shoveler have also been reported to nest in the coastal wetlands of Lake Erie (Appendix S).

In terms of other bird species, wetlands adjacent to the large sand spits such as Point Pelee, Rondeau Peninsula, and Long Point attract many migratory species that cross the lake. Several provincially significant bird species also occur in Lake Erie coastal wetlands. Bald Eagles nest near these wetlands, feed in them and also use them during migration. The swampy woodlands associated with the marshes support rare species such as Great Egret and Black-crowned Night-Heron. Other provincially significant birds nesting in Lake Erie wetlands include Wilson's Phalarope, King Rail, Least Bittern, Little Gull, Forster's Tern, Black Tern, Acadian Flycatcher, White-eyed Vireo, Prothonotary Warbler, Louisiana Waterthrush, Cerulean Warbler and Yellow-headed Blackbird (Appendix S).

Three provincially significant lepidopteran species, Mulberry Wing, Two-spotted Skipper and Duke's Skipper, have been reported in the coastal wetlands of Lake Erie (Appendix T).

About 20 species of mammals utilize Lake Erie marshes (Herdendorf, 1992). Furbearers such as Raccoon and Mink can be found near the marshes where they feed, and Muskrat are common throughout. White-tailed Deer are common around the upland edges of many of these wetlands.

5.7.3 Wetland Status

Estimates of the loss of coastal wetlands along the Canadian side of Lake Erie are not comprehensive. Loss of coastal wetland area has mostly occurred in the vicinity of the large sand spits such as Point Pelee. Agricultural land drainage has been identified as the most significant factor in the decline of these wetlands (Lynch-Stewart, 1983). For instance, the area of Point Pelee Marsh declined by 71%, from 3,878 ha in 1880 to 1,126 ha in the mid-1970s (Rutherford, 1979). The bulk of wetland drainage occurred in the 1890s when 50% of Point Pelee Marsh was converted into agricultural lands. Large portions of remaining coastal marshes are either parks or are privately-owned and managed for waterfowl hunting. Although wetland loss has slowed in recent decades, site-specific incremental loss is still a concern.

In addition to actual loss of coastal wetland acreage along the shores of Lake Erie, the quality of many remaining wetlands has been degraded by numerous stressors, especially excessive loading of sediment and nutrients, contaminants, shoreline modification, changes in sediment budgets, exotic species and diking of wetlands.

Turbidity and excessive suspended solids are significant stressors to coastal wetlands of Lake Erie (Herdendorf ,1992, 1987; Jude and Pappas, 1992; Appendix K). The waters of marshes and many bays have become turbid in the last century as a result of erosion from agriculture, dredging, diking and drainage of many large wetlands, shoreline modification and the introduction of Common Carp (Herdendorf, 1987). Many tributaries of western Lake Erie have watersheds dominated by clay soils; the western basin is consequently more turbid than the waters of the rest of the lake (Herdendorf, 1987). Although there have been efforts to reduce suspended solid inputs from tributaries, especially in western Lake Erie, they have not declined significantly in the last two decades (Richards and Baker, 1993).

Excessive nutrient loading is a common stressor in Lake Erie coastal wetlands in the U.S. and Canada (Herdendorf, 1987; Appendix K). Excess phosphorus is associated with excess inputs of suspended solids. Loadings of phosphorus to the watershed from point and non-point sources have reduced over the last two decades due to control measures (Dolan, 1993; Richards and Baker, 1993). However, nitrogen loading from non-point sources, mainly agricultural runoff, has increased in several watersheds (Richards and Baker, 1993). Wetlands with inflowing tributaries and barrier beaches are particularly prone to stresses from excess nutrients.

Pesticide loading from agricultural runoff has been identified as a significant stressor (Herdendorf, 1987); however, the impacts on coastal wetlands are not clear. As with phosphorus, it is associated with the suspended sediment load. Coastal wetlands with inflowing tributaries and barrier beaches are especially vulnerable.

Shoreline modification is a site-specific stressor when adjacent to coastal wetlands along the north shore of Lake Erie. It is mostly associated with urban encroachment and cottage development (Appendix K). Remaining wetlands suffer a backstopping effect from these revetments and dikes since the wetland communities cannot migrate upslope during high water years. This reduces wetland area during high water years and can also reduce the extent and long-term diversity of wetland communities. Storms during high water years can aggravate this problem by removing large areas of remaining wetland.

The extensive use of revetments, groynes and other structures which protect shoreline properties has also limited the supply of sediments in the littoral drift in western Lake Erie. Barrier beaches protecting wetlands must be replenished with these sediments. The few remaining natural wetlands with barrier beaches and sand spits are now losing this protection, as losses to erosion cannot be replenished from littoral sediment drift. As a result, these wetlands are becoming increasingly exposed to wave erosion.

Most of the remaining marshes along the U.S. shoreline are encompassed by dikes, while on the Canadian side, relatively few are diked. While diking allows for more intensive management for waterfowl and other fauna, it also isolates the wetland from the lake, impairing many wetland functions. For instance, many fish species, such as Northern Pike, which require wetlands for part of their life cycle can no longer access these wetlands. Other common stresses in coastal wetlands of Lake Erie are invasive non-indigenous species such as Purple Loosestrife, Zebra Mussels and Common Carp (R. Thompson, pers. com., 1996; D. Hector, pers. com., 1996).

5.8 Niagara River

5.8.1 Setting

The Niagara River drains Lake Erie into Lake Ontario. The river is 56 km long and drops 100 m along its course, most of which is at Niagara Falls. Its shoreline extends 60 km on the Canadian side and is much longer on the U.S. side, extending 112 km, as a result of the complex shoreline along Grand Island (International Joint Commission, 1993). The natural shoreline of the river consists of low banks in the upper portion of the river and a deep gorge cut through sedimentary deposits in the lower river below Niagara Falls.

Several tributaries flow into the river from the U.S. and Canada, draining a watershed of 3,251 km^2 (New York State Department of Environmental Conservation, 1994; Envirosearch Limited, 1992), but they contribute only a small fraction of flow to the river. On the Canadian side, land uses are dominated by agriculture (32%), abandoned agricultural land (23%), urban land (23%) and forests (16%) (Envirosearch, 1992).

The fast flow of the river has precluded the development of wetlands in many reaches of the river. However, wetlands and beds of submergent macrophytes are present in the upper reaches of the river (G. Birch, pers. com., 1996; Herdendorf, 1992). There are four small wetlands totalling approximately 85 ha, ranging in size from just under 5 ha to just over 37 ha (Figure 5; Appendix M). They are riverine in nature, sometimes with large palustrine components, and all but the Navy Island Marsh are associated with creek mouths. All have swamp and marsh components. The largest wetlands are the marshes on Navy Island (26 ha) and at Black Creek Wetland (37 ha).

5.8.2 Significant Features

Most of the provincially significant plant species found in the Niagara River are associated with coastal wetlands. Examples include Arrow-arum, Red-rooted Nut Sedge, Smith's Tufted Bulrush, Honey Locust, Swamp Rose Mallow, and Swamp Star Sedge (Appendix P). See Appendix P for complete listings.

Fifty-nine species of fish that use coastal wetlands on a permanent or temporary basis have been reported from the Niagara River (Mandrak and Crossman, 1992). The submerged vegetation of the wetlands in the upper river provides important spawning and nursery grounds for sportfish such as Muskellunge, Northern Pike, and Smallmouth Bass (Herdendorf, 1992). Niagara River wetlands also provide year-round habitat for three provincially significant fish species: Lake Chubsucker, Grass Pickerel and Black Bullhead (Appendix Q).

Coastal wetlands of the Niagara River also provide habitat for a wide range of amphibians, reptiles, birds and mammals. The only reported occurrence in Ontario for the Northern Dusky Salamander was in these wetlands (D. Sutherland, pers. com., 1996). No provincially significant amphibian, reptile or lepidopteran species were reported in Niagara River coastal wetlands (Appendix R; Appendix T).

Wetlands in the upper river are not heavily used by waterfowl for breeding or migration, but waterfowl numbers increase as winter approaches, when other wetlands in Lake Erie and Lake Ontario freeze (Bookhout, *et al.*, 1989). The dominant waterfowl include merganser species, Canvasback, Common Goldeneye, scaup species, Bufflehead, Mallard and Black Duck (Herdendorf, 1992; Bookhout *et al.*, 1989). Niagara River wetlands also provide important nesting habitat for the provincially significant Black-crowned Night-Heron (Appendix S).

5.8.3 Wetland Status

There is no comprehensive information on wetland loss on the Niagara River. Recent loss has been reported from one location, the Black Creek Wetland, as a result of urban development (Appendix M). Loss and stress on wetlands from shoreline modification and urban encroachment continue to be of concern (G. Birch, pers. com., 1996).

Several other human stressors affect remaining wetlands. The Niagara River and a tributary, the Buffalo River, have been declared Areas of Concern (AOCs) as a result of excessive toxic chemicals in the water, sediment contamination, fish edibility restrictions, the incidence of tumours in fish, degraded benthos and elevated phosphorus levels (Hartig and Thomas, 1988). Sources include industry outfalls, sewage treatment plants, other point sources, and non-point sources (New York State Department of Environmental Conservation, 1994; Envirosearch Limited, 1992; Hartig and Thomas, 1988). Wetlands near these sources are vulnerable to eutrophication and contamination from toxic chemicals. Nearshore areas, including wetlands, are deposition zones for sediments in the river and are therefore especially susceptible to sediment contamination (New York State Department of Environmental Conservation, 1994).

Water taking is another stress to coastal wetlands. More than half of the flow of the Niagara River is diverted for power production, causing dewatering of some marsh areas (G. Birch, pers. com., 1996; New York State Department of Environmental Conservation, 1994). This is exacerbated in some areas by road crossings that restrict wetland hydrology.

5.9 Lake Ontario

5.9.1 Setting

Lake Ontario is the smallest of the Great Lakes with a surface area of 18,960 km^2. However, it is relatively deep, and its average depth is second only to Lake Superior. The shoreline extends for 1,168 km and is particularly complex in the eastern third of the lake, with many embayments and peninsulas. The only islands occur in the eastern end of the lake, near its outlet. The Lake Ontario drainage basin is 60,600 km^2 (Fuller *et al.*, 1995) and is dominated by forests and agriculture, with lesser amounts of urban area and other land uses (PLUARG, 1978). Large urban centres occur in the western end of the lake around Toronto and Hamilton and at Rochester, New York. Water levels are controlled by dams and locks in the St. Lawrence Seaway along the St. Lawrence River.

Eighty-two coastal wetlands have been evaluated by the OMNR for Lake Ontario (not including wetlands in the Niagara River), totalling approximately 11,335 ha (Figure 5; Appendix M). There are several other significant wetland areas that have not been evaluated, but have been identified by OMNR biologists as in need of evaluation or that have been designated by the NHIC as Natural Areas (Figure 5; Appendix N). Wetlands are most abundant in the eastern portions of the lake. They occur at river mouths, nestled in embayments, and behind bars and barrier beaches. Wetlands are typically emergent and submergent marshes with a swamp or shrub-scrub component along the upland margins. The distribution of wetlands varies with shoreline geomorphology; from Niagara to Toronto, the shoreline consists mostly of bluffs and low cliffs except in the Hamilton area which has low-lying beaches (International Joint Commission, 1989). In this area, coastal wetlands are restricted to protected locations behind barrier beaches and in drowned mouths of rivers and creeks (Whillans *et al.*, 1992). Jordan Station Marsh in the Niagara Region is the largest wetland of western Lake Ontario, covering 136 ha. Many wetlands in western Lake Ontario have suffered from extensive filling and loss, especially around Toronto (Whillans, 1982). Remaining wetlands in the Toronto region include the Credit River Marshes (14 ha), Humber River Marshes (26 ha) and Rouge River Marshes (68 ha) at river mouths, and Rattray Marsh (10 ha) and the Toronto Island Wetland (22 ha) behind barrier beaches.

East of Toronto to Presqu'ile Point, the shore is steep with few wetlands; the majority that are present are found at river mouths and behind barrier beaches. Typical river mouth wetlands in this stretch of shoreline are Lynde Creek Marsh (110 ha) and Grafton Creek Swamp (62 ha); and Cranberry Marsh (32 ha) and Oshawa Second Marsh (105 ha) are examples of barrier beach wetlands.

From Presqu'ile Point to the mouth of the St. Lawrence River, the shoreline is complex with several channels, embayments, headlands, and islands. As a result, there are many protected sites suitable for wetlands. Thirty-nine wetlands (excluding Wolfe Island) totalling approximately 8,877 ha are found here; roughly 80% of the wetland area along the Canadian shore of the lake. The wetland along Presqu'ile Point (970 ha) is especially diverse, consisting mostly of marsh with some swamp and a small coastal meadow marsh (fen). The southwest shores of Prince Edward Peninsula are dominated by large marshes that are protected by bay mouth bars and are connected landward to lagoons. The largest areas include Wellers Bay Complex (363 ha), Pleasant Bay (299 ha), Hucyks Bay (245 ha), West Lake (706 ha) and East Lake (230 ha). The Bay of Quinte along the north and east side of the peninsula has a very complex shoreline, and extensive marshes have developed in the many sheltered bays, around islands and at creek mouths. The most notable are Pleasant Bay Marsh (299 ha), Big Island Marsh (858 ha), Sawguin Creek (2,093 ha), Dead Creek Marsh (359 ha), Big Marsh (400 ha), and Hay Bay Marsh (1,333 ha). Some of these coastal wetlands also include large palustrine components (Appendix M).

5.9.2 Significant Features

Coastal wetlands of Lake Ontario consist mostly of submergent and emergent marshes, swamps, and a few rare coastal meadow marsh (fen) communities (Natural Heritage Information Centre, 1995; Smith *et al.*, 1991; Herdendorf *et al.*, 1981). In many marsh, wet meadow or submergent habitats, invasive plant species are often introduced, such as Purple Loosestrife, Eurasian Water-milfoil, Reed Canary Grass and hybrid cattail (International Joint Commission, 1993; Wilcox *et al.*, 1993). Despite these problems, twenty provincially significant species of plants have been found in Lake Ontario's coastal wetlands (Appendix P). For instance, Hidden-fruited Bladderwort, Winged Loosestrife, Branching Burreed and Yellow Pond Lily occur in several marsh habitats. Other significant species such as Bushy Aster, Low Nut Rush and Smith's Tufted Bulrush are found in the few coastal meadow marshes along Lake Ontario (Appendix P).

Sixty-eight species of fish use the coastal wetlands of Lake Ontario, two thirds of which are permanent residents (Jude and Pappas, 1992; Stephenson, 1990). Gizzard Shad, White Perch and Freshwater Drum are the most abundant permanent residents, while Alewife, Rainbow Smelt, White Sucker, Smallmouth Bass, Spottail Shiner, Johnny Darter, Trout-Perch, Walleye and Yellow Perch are common species which use coastal wetlands on a temporary basis for spawning, nursery or feeding. Several important sportfish use these wetlands including Common Carp, Northern Pike, Muskellunge, Pumpkinseed, Bluegill, Black Crappie, White Crappie, Largemouth Bass, Smallmouth Bass, Walleye, Yellow Perch, and White Bass (Glooschenko *et al.*, 1987; Herdendorf *et al.*, 1981). Several species of fish in these coastal wetlands are considered provincially significant, including Eastern Silvery Minnow, Pugnose Shiner and Grass Pickerel (Appendix Q).

Lake Ontario coastal wetlands provide important habitat for reptiles and amphibians. Bullfrog, Green Frog, Northern Leopard Frog, Spring Peeper and Grey Tree Frog are common frogs which inhabit coastal marshes and swamps (M. Oldham, pers. com., 1996; Herdendorf *et al.*, 1981). The Mudpuppy and Red-spotted Newt are other amphibians commonly found in submerged macrophyte beds in coastal wetlands. Common reptiles include Snapping Turtle, Midland Painted Turtle, Northern Water Snake, Eastern Garter Snake and Eastern Milk Snake. As well, three provincially significant species of reptiles and amphibians have also been found: the Eastern Spiny Softshell Turtle, Spotted Turtle, and Jefferson Salamander (Appendix R).

Coastal wetlands along the northeast coast around Prince Edward Peninsula have been identified as important areas for waterfowl in the Great Lakes (Prince *et al.*, 1992). Prince Edward Peninsula in particular is the third most significant region for waterfowl in Ontario, after Long Point and the St. Clair Delta (Bookhout *et al.*, 1989). Their importance is mostly linked to providing staging habitats during spring and fall migration. Large numbers of diving ducks, especially scaup and merganser species, are attracted to these wetlands in the fall (Prince *et al.*, 1992). The provincially significant Northern Shoveler, Canvasback and Redhead can also be found in large numbers in the marshes and bays along the Prince Edward Peninsula. Wetlands in western Lake Ontario near Toronto have largely disappeared, but concentrations of dabbling ducks, diving ducks and Canada Geese still are present along the shoreline during fall and winter migration. Coastal wetlands of Lake Ontario are of lesser importance for breeding waterfowl, but many dabbling ducks do nest in them, including Mallard, Black Duck, Blue-winged Teal and Wood Duck (Prince *et al.*, 1992; Bookhout *et al.*, 1989). See Appendix S for a complete listing of significant waterfowl species found in Lake Ontario coastal wetlands.

Coastal wetlands of Lake Ontario provide feeding, nesting or migration habitat for many other bird species. Several raptors nest or migrate through these coastal wetlands including Peregrine Falcon, Bald Eagle, Sharp-shinned Hawk, Cooper's Hawk, Red-shouldered Hawk, and Northern Harrier (D. Sutherland, pers. com., 1996; Herdendorf *et al.*, 1981). Several species of waterbirds such as Double-crested

Cormorant, Great Blue Heron, American Bittern, and Green Heron have been recorded in these wetlands (Glooschenko *et al.*, 1987; Herdendorf *et al.*, 1981). Common Terns regularly breed in north shore wetlands, but are uncommon along the south shore. Caspian Terns are also common in wetlands along the northeast shore. Provincially significant species of birds nest and feed in these wetlands, including Least Bittern, Black-crowned Night-Heron, Black Tern, and Short-eared Owl (Appendix S). See Appendix S for complete listings.

Significant lepidopteran species including the Mulberry Wing and Two-spotted Skipper, are found in the coastal meadow marshes of Lake Ontario (Appendix T).

The coastal wetlands of Lake Ontario are important habitats for many mammals. Beaver, Muskrat, Mink and River Otter are highly dependent on these wetlands (Glooschenko *et al.*, 1987; Herendorf *et al.*, 1981). White-tailed Deer, Red Fox, Short-tailed Weasel, Raccoon and Coyote are examples of other species using Lake Ontario wetlands.

5.9.3 Wetland Status

The wetlands of Lake Ontario have suffered severe loss over the last two centuries (International Joint Commission, 1989; Lynch-Stewart, 1983; McCullough, 1982; Whillans, 1982). The main causes are agricultural drainage and urban encroachment. Between 1789 and 1979, Whillans (1982) estimated the loss of coastal marsh along the Canadian shore west of the Bay of Quinte to be 1,920 ha or 43% of the original marsh area. Similar estimates of wetland loss were reported by McCullough (1982). This loss was greatest from Toronto to the Niagara River where an estimated 1,518 ha of coastal marsh have been lost; this represents the loss of 73% to 100% of the original marsh along these shores (Whillans, 1982). The greatest losses occurred in the late 1800s and early 1900s when large marshes were filled and dredged for shipping, industrial and urban uses. East of Toronto, less marsh area has been lost. Between Toronto and Presqu'ile, 646 ha or 32% of the original marsh has been lost, while from Presqu'ile to the Bay of Quinte, only 347 ha or 7.5% has been lost. Large losses have occurred in the Bay of Quinte. Around 12,008 ha of wooded and emergent wetlands were lost prior to the 1960s within a 3.2 km strip inland from the bay, mainly due to agricultural drainage (Ontario Ministry of the Environment *et al.*, 1990). Between 1967 and 1982, a further 412 ha of coastal wetlands were lost in this area, but since 1967, 542 ha have been reclaimed from agricultural use and restored to wetland. Wetland loss along the Canadian shores continues to be a concern as a result of urban encroachment (Appendix M).

Remaining wetlands are affected by several other human stressors. A major stressor to all coastal wetlands in Lake Ontario is water level regulation. Water levels have been regulated in the lake since construction of the St. Lawrence Seaway in 1959. Regulation of water levels seeks to reduce the occurrence of both high and low lake levels. Prior to regulation, the range of water level fluctuations during the 20th century was about 2 m. Between 1960 and 1976, following regulation, this range was reduced slightly and was further reduced to about 0.9 m after 1976 (Wilcox *et al.*, 1993). Regulation also prevented water levels from reaching record highs in 1986 as they did on all other lakes. The lack of alternating flooded and dewatered conditions at the upper and lower edges of the wetlands decreased wetland area and decreased the diversity of plant and wildlife communities (Wilcox *et al.*, 1993; Busch *et al.*, 1990; International Joint Commission, 1989). In general, upland species became more prevalent along the upper edges of the wetlands. Emergent communities declined, and submerged aquatic macrophyte beds increased. As well, invasive plants began to dominate wetland communities, for example extensive stands of cattail are now established in these wetlands, and some areas are dominated by Purple Loosestrife, Reed Canary Grass and various shrubs (Wilcox *et al.*, 1993).

High sediment loads and excess turbidity have been noted as stressors in several Lake Ontario coastal wetlands. Examples include Cootes Paradise in Hamilton (Painter *et al.*, 1989; Simser, 1979), Oshawa Second Marsh (Cecile, 1983; Morris, 1983) and wetlands of the Bay of Quinte (Crowder and Bristow, 1988). Sources are site-specific and are mostly related to urban and agricultural runoff. Common Carp are also a serious related problem in Lake Ontario as they resuspend sediments, increase turbidity and destroy aquatic macrophytes (Painter *et al.*, 1989).

Turbidity problems are compounded by excess nutrients that encourage algae and in turn decrease water clarity (Painter *et al.*, 1989). Excess nutrients have also caused the eutrophication of wetland communities, reducing the diversity of wetland vegetation in some areas. These stressors are especially evident in the Bay of Quinte (Crowder and Bristow, 1988).

Contaminants are site-specific stressors to coastal wetlands. Several sites around Lake Ontario have been declared Areas of Concern including Hamilton Harbour, Toronto, Port Hope and the Bay of Quinte. Cootes Paradise in Hamilton and the Bay of Quinte have large areas of coastal wetlands and are therefore especially susceptible to contamination by toxic chemicals. Both these areas are known to have contaminated sediments or bioaccumulation of contaminants in biota (Bishop *et al.*, 1991,1995; Crowder *et al.*, 1989).

Shoreline modification is another site-specific stressor (Appendix M). Dikes or revetments not only fill wetlands but also prevent the migration of remaining wetland communities in response to fluctuating water levels.

5.10 St. Lawrence River

5.10.1 Setting

The St. Lawrence River drains Lake Ontario and is the outlet of the Great Lakes system. It extends 870 km in length from Lake Ontario to the Gulf of St. Lawrence (Grant, 1995). For the Atlas only, the 186 km section of the river from Wolfe Island at the outlet of Lake Ontario to the Quebec border is discussed. This includes the international section of the river and the Ontario shore of Lake St. Francis. Other studies examine the state of the river and its wetlands through Quebec to the Gulf of St. Lawrence (Centre St. Laurent, 1996a, b).

Water level and flows have been regulated in this section of the St. Lawrence River since the construction of the St. Lawrence Seaway in 1959. Since then, dams and water control structures have greatly changed the character of the river and its wetlands. This section of the St. Lawrence River can be divided into four distinct sections, each displaying different physical and biological characteristics: the Thousand Islands, Middle Corridor, Lake St. Lawrence, and Lake St. Francis (Grant, 1995; Busch and Patch, 1990). The Thousand Islands section lies in the uppermost reach of the river. It has a rocky shoreline and many islands, bays, shoals and quiescent areas with extensive wetlands. The Middle Corridor extends a distance of 49 km from just west of Brockville/Morristown downstream to the Iroquois Control Dam. This is the most riverine section of the St. Lawrence, with a single deep, wide channel and a relatively uniform shoreline. Currents are very strong and wetlands are restricted to small bays, shoreline indentations, and tributary mouths. At Iroquois, the water is shallower and there are some extensive vegetated shallow areas.

Lake St. Lawrence occurs in the lower reach of the international section of the river, extending from the Iroquois Control Dam to the Moses-Saunders Dam near Cornwall. This area was changed completely from a riverine environment to a lacustrine environment following the construction of the Seaway. It has numerous islands and shoals. Wetlands are relatively common in this section, however the extent of marshes is quite small compared to that of the Thousand Islands section. Below the Moses-Saunders Dam, there is a short stretch of river with fast current leading to Lake St. Francis. Lake St. Francis is a lacustrine environment with extensive wetlands located at creek mouths, in embayments and surrounding islands (Grant, 1995).

Along the Ontario shoreline, there are 40 evaluated wetlands totalling approximately 7,018 ha (Figure 6; Appendix O). The wetlands along the Ontario shoreline range in size from just under 4 ha to 1,398 ha. They are primarily riverine marshes with relatively small swamp components. More than 60% of the wetlands both in number and area are found in the Thousand Islands section of the river. The largest wetlands in this reach include the Greater Cataraqui Marsh (504 ha), the Wolfe Island Complex (1,398 ha), the Grenadier Island Complex (868 ha). The Wolfe Island Complex has a large bog component, one of only two in the lower Great Lakes. The Morrisburg Swamp (391 ha), the only swamp found along the Ontario St. Lawrence River shoreline, and the Upper Canada Migratory Bird Sanctuary (321 ha) are large wetlands in the Lake St. Lawrence Section. Along the Ontario shores of Lake St. Francis, Charlottenburg Marsh (851 ha) and Bainsville Bay Marsh (407 ha) are among the largest wetlands.

Although seven unevaluated coastal wetlands on the St. Lawrence River (Table 3) were identified through the 1996 OMNR District/Area Office Survey, these were not identified individually through Atlas methods and are not plotted on Figure 6, nor in an Appendix.

5.10.2 Significant Features

The coastal wetlands of the St. Lawrence River are predominantly cattail marshes with areas of submergents and floating plants such as water lilies, and occasional swamp components, mostly dominated by willow (Smith *et al.*, 1991). Seven provincially significant plant species have been recorded in the shallow water marshes and adjacent wet beaches of the St. Lawrence River including, Arrow-arum, Narrow-leaved Water-plantain and Smith's Tufted Bulrush (Appendix P). The other provincially significant plant species found in these coastal wetlands are Follicle Sedge, Eastern Prairie White Fringed Orchid, and Branching Bur-reed.

At least 64 species of fish inhabit wetlands of the St. Lawrence River, 42 of which are permanent residents (Jude and Pappas, 1992; Mandrak and Crossman, 1992; Patch and Busch, 1984). The major recreational fisheries support Muskellunge, Northern Pike, Brown Bullhead, Smallmouth Bass, Yellow Perch, Walleye and a variety of panfishes (Grant, 1995). Wetlands in the St. Lawrence River support one of only a few large self-sustaining populations of Muskellunge in North America (Grant, 1995). Three provincially significant species also depend on these wetlands to provide habitat on a permanent basis: the Eastern Silvery Minnow, Pugnose Shiner and Grass Pickerel (Sutherland, 1994; Mandrak and Crossman, 1992; Appendix Q).

The coastal shoreline wetlands of the St. Lawrence River provide notable migration and staging habitat for waterfowl. The St. Lawrence Lowlands have been identified in the North American Waterfowl Management Plan as an important waterfowl staging area. Wolfe Island in the Thousand Islands section of the river is located on a northwest-southeast migration route, and is surrounded by abundant shallow water areas with submerged vegetation. Large numbers of waterfowl, primarily scaup and teal, are found in this area during spring and fall migration (Ross, 1984). The Wolfe Island area is also an important staging area for Canada Geese. Provincially significant waterfowl species such as Northern Shoveler, and Redhead breed in St. Lawrence River wetlands (Appendix S). The St. Lawrence River wetlands also provide important nesting habitat for many colonial waterbirds. Several provincially significant bird species breed here including Least Bittern, Black Tern, and Short-eared Owl (Appendix S).

Many reptiles and amphibians use the wetlands along the St. Lawrence River for nesting and spawning, nursery and/or feeding sites. However, only one provincially significant species, the Spotted Turtle, has been recorded (M. Oldham, pers. com., 1996; Appendix R).

One provincially significant lepidopteran species, the Two-spotted Skipper, has been recorded for St. Lawrence River coastal wetlands (Appendix T).

A variety of mammal species are found in St. Lawrence River wetlands including Muskrat, Mink, Red Fox and Coyote. Several wetlands also provide regionally significant winter habitat for White-tailed Deer.

5.10.3 Wetland Status

The St. Lawrence River has experienced a wide variety of environmental disturbances since the first channel modifications in the late 18th century (Grant, 1995). The largest disturbance was associated with the construction and operation of the St. Lawrence Seaway. Prior to the construction of the Seaway, the river resembled a large riverine estuary in the Thousand Island section. The Middle Corridor and Lake St. Lawrence (to Cornwall) were part of a riverine system with many islands and shoals, and many rapids in the lower reaches of the international section (Busch and Patch, 1990). The creation of Lake St. Lawrence and the dredging for navigation and power production greatly altered these habitats. These changes have been monitored along the U.S. side (Busch and Patch, 1990; Patch and Busch, 1984).

Wetland change on the Canadian side of the St. Lawrence River is not well documented.

After the initial opening of the Seaway, relatively little change occurred along the U.S. shore in the Thousand Islands section of the river, but changes became apparent at Galop Island in the Middle Corridor section of the river and were most dramatic in Lake St. Lawrence. Numerous islands and shoals were flooded along with several major rapids, and there were large increases in deepwater and littoral habitat. Wetland habitats changed greatly. By 1962, shortly after the construction of the Seaway, there was a decline in wetland area of 11.4% along the U.S. shore (Patch and Busch, 1984). However by 1988, wetland area had increased by 106 ha (9%) along the U.S. shoreline as compared to the pre-Seaway condition (Busch and Patch, 1990). Between 1955 and 1988, there were losses of 174 ha (18%) and 10ha (40%) along the U.S. shoreline in the Thousand Islands and Middle Corridor sections, respectively. There was an increase of 448 ha (64%) on the U.S. side of Lake St. Lawrence. There were also large changes in wetland community structure as compared to the pre-Seaway condition as a result of the regulation and stabilization of water levels. Emergent wetlands decreased in area to the benefit of broad-leaved forested and shrub-scrub wetlands. In Lake St. Lawrence, wetlands are relatively common, but emergent wetland area is small, primarily due to the regulation of water levels.

Changes in wetland area and stressors due to human activities tend to be smaller scale and site-specific. In the Thousand Islands section of the river, recreational activities dominate with many cottages, picnic and camping areas, and marinas. Along the Canadian side of the river, shoreline modification from these activities has historically resulted in irretrievable direct losses of wetland habitat, although these losses have been greatly reduced over the last five to six years (R. Cholmondeley pers. com., 1996). Recent losses have been limited to development of occasional shoreline protection structures, although some illegal dredging still occurs. Pressures from these activities are ongoing, as marinas try to expand or dredge deeper to allow for bigger boats, or to compensate for fluctuating water levels. Recreational boats also stress remaining wetlands by creating wakes that disturb and erode emergent and submergent vegetation.

Wetland losses on the Canadian shore of the Thousand Islands section of the river have also resulted from the construction of roads such as the St. Lawrence Parkway as well as from numerous access roads to cottages (Appendix O). Not only is there wetland loss from filling of the roadbed, but many segments of wetlands are cut off from the river and wetland hydrology is altered.

Other stressors affect remaining wetlands in the Thousand Islands section. Water level regulation, which draws down water levels in the fall, and leads to plant freeze-out or ice scour has dramatically affected wetland vegetation (Appendix O). Nutrient and sediment loading from tributary creeks also act as local wetland stressors in the Thousand Islands. In addition, non indigenous species have taken their toll on wetlands. For instance, Zebra Mussels have increased water clarity, but there appears to be a consequent reduction in aquatic plant growth and an increase in filamentous algae (R. Cholmondeley, pers. com., 1996).

The Middle Corridor has been substantially altered by dredging and filling which accompanied the construction of the Iroquois Control Dam, locks and the Seaway navigational channel between 1954 and 1959 (Grant, 1995; Busch and Patch, 1990). Residential and industrial developments are common along the Canadian shoreline and few wetlands remain in this reach. Dredging and filling have been associated with these activities, resulting in direct losses in the past as well as concerns regarding future losses (R.

Grant, pers. com., 1996). These activities have also stressed remaining wetlands through sediment and nutrient loading (R. Grant, pers. com., 1996).

In Lake St. Lawrence, the greatest stresses relate to the construction and operation of the Moses-Saunders Dam, as detailed above. Stresses on existing wetlands in this area are site specific, and are primarily related to industrial, commercial, residential and recreational development (M. Eckersley, pers. com., 1996).

In Lake St. Francis, modifications to the hydrological regime have resulted in an increase of 36 cm in the mean water level, and there are no longer any annual water level fluctuations (Jean and Bouchard, 1991). The stable water levels mean that spring flooding does not occur in many wetlands (M. Eckersley, pers. com., 1996). Additionally, over the past 40 years, wetlands of Lake St. Francis have been subject to extensive urban, recreational and agricultural development (M. Eckersley, pers. com., 1996; Jean and Bouchard 1991). Between 1946 and 1983, a relatively low 7% of wetlands in Lake St. Lawrence disappeared. However wetlands in certain areas have been reduced by as much as 41% through conversion primarily to urban lands (Jean and Bouchard, 1991).

The Lake St. Francis section of the St. Lawrence River downstream of Cornwall, Ontario and Massena, New York has contaminant problems of concern for coastal wetlands. Bioaccumulation of PCBs has been observed to be very high in Red-Winged Blackbirds and Tree Swallows from coastal wetlands in Akwesasne downstream of Cornwall/Massena (Bishop *et al.*, 1995). This section of the river has been declared an Area of Concern as a result of excessive toxic substances in the water, contaminated sediments at Cornwall and at the mouth of the Grass River near Massena, fish consumption advisories, tumours in fish near Cornwall, degraded benthos, elevated faecal coliform bacteria counts, and eutrophication from elevated phosphorus downstream of Cornwall.

6.0 References

Allen, G. 1996. Personal communication. Midhurst District Office, Ontario Ministry of Natural Resources, Midhurst, Ontario.

Bakowsky, W. 1996. *Natural Heritage Resources of Ontario: Vegetation Communities of Southern Ontario.* Natural Heritage Information Centre, Ontario Ministry of Natural Resources, Peterborough, Ontario.

Bakowsky, W. 1996. Personal communication. Natural Heritage Information Centre, Ontario Ministry of Natural Resources, Peterborough, Ontario.

Ball, H. and J. Tost. 1992. *Summary of Small Fish Surveys Conducted in Rivers Entering Thunder Bay Harbour, 1990.* Technical Report #11, North Shore of Lake Superior Remedial Action Plans.

Bedford, K.W. 1992. The physical effects of the Great Lakes on tributaries and wetlands, a summary. *Journal of Great Lakes Research* **18**, 571-589.

Birch, G. 1996. Personal communication. Fonthill Area Office, Ontario Ministry of Natural Resources, Fonthill, Ontario.

Bishop, C.A., R.J. Brooks, J.H. Carey, P. Ng, R.J. Norstrom, and D.R.S. Lean. 1991. The case for a cause-effect linkage between environmental contamination and development in eggs of the common snapping turtle (*Chelydra serpentina serpentina*) from Ontario. *Canadian Journal of Toxicology and Environmental Health* **33**, 521-548.

Bishop, C.A., M.D. Koster, A.A. Chek, D.J.T. Hussell, and K. Jock. 1995. Chlorinated hydrocarbons and mercury in sediments, red-winged blackbirds (*Agelaius phoeniceus*) and tree swallows (*Tachycineta bicolor*) from wetlands in the Great Lakes-St. Lawrence River Basin. *Environmental Toxicology and Chemistry* **14**, 401-501.

Black, R. 1996. Personal communication. Parry Sound District Office, Ontario Ministry of Natural Resources, Parry Sound, Ontario.

Bookhout, T.A., K.E. Bednarik, and R.W. Kroll. 1989. The Great Lakes marshes. In: Smith, L.M., Pederson, R.L., Kaminski, R.M. (eds.). *Habitat Management for Migrating and Wintering Waterfowl in North America*. pp. 131-156. Texas Tech University Press, Lubbock, Texas.

Bray, K. 1996. Personal communication. Lake Superior Remedial Action Plan Program Office, Thunder Bay, Ontario.

Bray, K. 1993. *Preliminary Inventory and Evaluation of Littoral Habitat in the St. Marys River Area of Concern.* St. Marys River Remedial Action Plan Report.

Bray, K. 1992. *Changes to Fish Habitat of the St. Marys River: a Retrospective Analysis.* M.Sc. Thesis, Trent University, Peterborough, Ontario.

Burt, A.J., P.M. McKee, D.R. Hart, and P.B. Kauss. 1991. Effects of pollution on benthic invertebrate communities of the St. Marys River, 1985. *Hydrobiologia* **219**, 63-81.

Busch, W.-D.N., R.O. Osborn, and G.T. Auble. 1990. The effects of water levels on two Lake Ontario wetlands. In: Kusler, J. and R. Smardon (eds.) *Proceedings: International Symposium on Wetlands of the Great Lakes*. pp. 92-96, Association of State Wetland Managers, Niagara Falls, N.Y.

Busch, W.-D.N. and S. Patch. 1990. Human-Caused Habitat Changes in the International Section of the St. Lawrence River. In: Kusler, J. and R. Smardon (eds.). *Proceedings: International Symposium on Wetlands of the Great Lakes*, pp. 141-155. The Association of State Wetland Managers, Niagara Falls, N.Y.

Cadman, M.D. 1996. Personal communication. Canadian Wildlife Service, Environment Canada, Guelph, Ontario.

Cadman, M.D., P.F.J. Eagles, and F.M. Helleiner. 1987. *Atlas of the Breeding Birds of Ontario.* Federation of Ontario Naturalists and Long Point Bird Observatory. University of Waterloo, Waterloo ,Ontario.

Cecile, C. P. 1983. *Oshawa Second Marsh Baseline Study, Final Report.* Environment Canada.

Centre St. Laurent. 1996a. *The State of the Environment Report on the St. Lawrence River, Vol. 1: The St. Lawrence Ecosystem.* Environment Canada & Editions MultiMondes, Montreal, Quebec.

Centre St. Laurent. 1996b. *The State of the Environment Report on the St. Lawrence River, Vol. 2: The State of the St. Lawrence.* Environment Canada & Editions MultiMondes, Montreal, Quebec.

Cholmondeley, R. 1996. Personal communication. Brockville Area Office, Ontario Ministry of Natural Resources, Brockville, Ontario.

COSEWIC, 2002. Canadian Species at Risk, November 2002. Committee on the Status of Endangered Wildlife in Canada. [http://www.cosewic.gc.ca].

COSSARO, 2002. Vulnerable, Threatened, Endangered, Extirpated or Extinct Species of Ontario, September, 2002. Ontario Ministry of Natural Resources, Committee on the Status of Species at Risk in Ontario. [http://www.mnr.gov.on.ca/mnr/VTEEElist_2002_eng.pdf].

Crowder, A.A. and J.M. Bristow. 1988. The future of waterfowl habitats in the Canadian lower Great Lakes wetlands. *Journal of Great Lakes Research* **14**, 115-127.

Crowder, A., W.T. Dushenko, J. Greig, and J.S. Poland. 1989. Metal contamination in sediments and biota of the Bay of Quinte, Lake Ontario, Canada. *Hydrobiologia* **188/189**, 337-343.

Cullis, K. 1996. Personal communication. Lake Superior Remedial Action Plan Program Office, Thunder Bay, Ontario.

Dextrase, A. 1996. Personal communication. Lands and Natural Heritage, Ontario Ministry of Natural Resources, Peterborough, Ontario.

Dolan, D. M. 1993. Point source loadings of phosphorus to Lake Erie: 1986-1990. *Journal of Great Lakes Research* **19**, 212-223.

Duffy, W.G., T.R. Batterson, and C.D. McNabb. 1987. *The St. Marys River, Michigan: An Ecological Profile.* U.S. Fish and Wildlife Service, Biological Report 85 (7.10).

Dupont, A. 1996. Personal communication. Sault Ste. Marie District Office, Ontario Ministry of Natural Resources, Sault Ste. Marie, Ontario.

Eckersley, M. 1996. Personal communication. Lower St. Lawrence Planning Region. Ontario Ministry of Natural Resources, Peterborough, Ontario.

Edsall, T.A., B.A. Manny, and C.N. Raphael. 1988. *The St. Clair River and Lake St. Clair, Michigan: An Ecological Profile.* U.S. Fish and Wildlife Service, Biological Report 85 (7.3).

Entwistle, J. 1986. *Thunder Bay Marshes Study, Summary Report 1982-1985.* Lakehead Region Conservation Authority Report, Thunder Bay, Ontario.

Environment Canada. 1995. *A Catalogue of Wetland Databases and Inventories for the Canadian Great Lakes Basin.* Great Lakes Wetlands Conservation Action Plan 1994 - 2000.

Environment Canada. 1994a. *Environmental Sensitivity Atlas for Lake Erie (Including the Welland Canal) and Niagara River Shorelines.* Prepared By: Environment Canada - Environmental Protection Branch (Ontario Region), Canadian Coast Guard, United States Coast Guard - District 9, United States National Oceanic and Atmospheric Administration. Minister of Supply and Services Canada, Toronto.

Environment Canada. 1994b. *Environmental Sensitivity Atlas for Lake Huron's Canadian Shoreline (Including Georgian Bay.* Prepared By: Environment Canada - Environmental Protection Branch (Ontario Region). Minister of Supply and Services Canada, Toronto.

Environment Canada. 1994c. *Environmental Sensitivity Atlas for the St. Clair River, Lake St. Clair and Detroit River Shoreline.* Prepared By: Environment Canada - Environmental Protection Branch (Ontario Region), Canadian Coast Guard, United States Coast Guard - District 9, United States National Oceanic and Atmospheric Administration. Minister of Supply and Services Canada, Toronto.

Environment Canada. 1994d. *Environmental Sensitivity Atlas for the St. Lawrence River Shorelines.* Prepared By: Environment Canada - Environmental Protection Branch (Ontario Region), Canadian Coast Guard, United States Coast Guard - District 9, United States National Oceanic and Atmospheric Administration. Minister of Supply and Services Canada, Toronto.

Environment Canada. 1994e. *Environmental Sensitivity Atlas for the St. Marys River Shorelines.* Prepared By: Environment Canada - Environmental Protection Branch (Ontario Region), Canadian Coast Guard, United States Coast Guard - District 9, United States National Oceanic and Atmospheric Administration. Minister of Supply and Services Canada, Toronto.

Environment Canada. 1993a. *Environmental Sensitivity Atlas for Lake Superior's Canadian Shoreline.* Prepared By: Environment Canada - Environmental Protection Branch (Ontario Region). Minister of Supply and Services Canada, Toronto.

Environment Canada. 1993b. *Environmental Sensitivity Atlas for Lake Ontario's Canadian Shoreline.* Prepared By: Environment Canada - Environmental Protection Branch (Ontario Region). Minister of Supply and Services Canada, Toronto.

Envirosearch Limited (ed.). 1992. *Niagara River Remedial Action Plan, Stage 1: Environmental Conditions and Problem Definition.* Ontario Ministry of the Environment, Environment Canada, Ontario Ministry of Natural Resources, Fisheries and Oceans Canada.

Fuller, K., H. Shear, and J. Wittig. (eds.). 1995. *The Great Lakes: An Environmental Atlas and Resource Book*. Government of Canada and United States Protection Agency.

Glooschenko, V., B. Parker, L. Coo, R. Kent, C. Wedeles, A. Mason, J. Dawson, D. Herman, and P. Smith. 1987. *Provincially and Regionally Significant Wetlands in Southern Ontario*. Interim Report, Wildlife Branch, Ontario Ministry of Natural Resources, Toronto, Ontario.

Grant, R. 1996. Personal communication. Brockville Area Office, Ontario Ministry of Natural Resources, Brockville, Ontario.

Grant, R.E. 1995. *Canada Ports Corporation 1994 Fisheries Act Violation, Port of Prescott*. Ontario Ministry of Natural Resources, Brockville, Ontario.

Griffiths, R.W. 1991. Environmental quality assessment of the St. Clair River as reflected by the distribution of benthic macroinvertebrates in 1985. *Hydrobiologia* **219**, 143-164.

Griffiths, R.W., S. Thornley, and T.A. Edsall. 1991. Limnological aspects of the St. Clair River. *Hydrobiologia* **219**, 97-123.

Harding, J.H. 1997. *Amphibians and Reptiles of the Great Lakes Region.* University of Michigan Press. Ann Arbour, Michigan.

Hartig, J.H. and R.L. Thomas. 1988. Development of plans to restore degraded areas in the Great Lakes. *Environmental Management* **12**, 327-347.

Hector, D. 1996. Personal communication. Chatham Area Office, Ontario Ministry of Natural Resources, Chatham, Ontario.

Herdendorf, C.E. 1992. Lake Erie coastal wetlands: an overview. *Journal of Great Lakes Research* **18**, 533 551.

Herdendorf, C.E. 1987. *The Ecology of the Coastal Marshes of Western Lake Erie: A Community Profile.* U.S. Fish and Wildlife Service, Biological Report 85 (7.9).

Herdendorf, C.E., S.M. Hartley, and M.D. Barnes (eds.). 1981. *Fish and Wildlife Resources of the Great Lakes Coastal Wetlands within the United States, Vol. 2: Lake Ontario.* U.S. Fish and Wildlife Service, FWS/OBS-81/02-v2.

Herdendorf, C.E., C.N. Raphael, and W.G. Duffy. 1986. *The Ecology of Lake St. Clair Wetlands: A Community Profile.* National Wetlands Research Center, Fish and Wildlife Service, U.S. Department of the Interior, Washington, D.C.

International Joint Commission. 1993. *Levels Reference Study, Great Lakes St. Lawrence River Basin. Annex Landuse and Management.* International Joint Commission.

International Joint Commission. 1989. *Living With the Lakes: Challenges and Opportunities. Annex B. Environmental Features, Processes and Impacts: An Ecosystem Perspective on the Great Lakes - St. Lawrence River System.* Functional Group 2, International Joint Commission, Water Levels Reference Study.

Jalava, J. 1996. Personal communication. Natural Heritage Information Centre, Ontario Ministry of Natural Resources, Peterborough, Ontario.

Jean, M. and A. Bouchard. 1991. Temporal changes in wetland landscapes of a section of the St. Lawrence River, Canada. *Environmental Management* **15**, 241-250.

Jones, S. 1996. Personal communication. Sault Ste. Marie District Office, Ontario Ministry of Natural Resources, Sault Ste. Marie, Ontario.

Jude, D.J. and J. Pappas. 1992. Fish utilization of Great Lakes coastal wetlands. *Journal of Great Lakes Research* **18**, 651-672.

Kauss, P.B. 1991. Biota of the St. Marys River: habitat evaluation and environmental assessment. *Hydrobiologia* **219**, 1-35.

Lane , J. A., C.B. Portt, and C.K. Minns. 1996. *Nursery Habitat Characteristics of Great Lakes Fishes.* Canadian Manuscript Report of Fisheries and Aquatic Sciences No. 2338.

Lee, D.S., C.R. Gilbert, C.H. Hocutt, R.E. Jenkins, D.E. McAllisstar, and J.R. Stauffer, Jr. 1980. ET SEQ. *Atlas of North American Freshwater Fishes.* North Carolina State Museum, Raleigh, N.C., U.S.A.

Lynch-Stewart, P. 1983. *Land Use Change on Wetlands in Southern Canada: Review and Bibliography.* Working Paper No. 26, Lands Directorate, Environment Canada. Ottawa, Ontario.

MacLean, N. 1996. Personal communication. Napanee Area Office, Ontario Ministry of Natural Resources, Napanee, Ontario.

Mandrak, N.E. and E.J. Crossman. 1993. Status of the lake chubsucker (*Erimyzon sucetta*) in Canada. In report to the Committee on the Status of Endangered Wildlife in Canada. Ottawa, Ontario.

Mandrak, N. and E.J. Crossman. 1992. *A Checklist of Ontario Freshwater Fishes.* Royal Ontario Museum, Toronto, Ontario.

Manny, B.A. and D. Kenaga. 1991. The Detroit River: effects of contaminants and human activities on aquatic plants and animals and their habitats. *Hydrobiologia* **219**, 269-279.

Manny, B.A., T.A. Edsall, and E. Jaworski. 1988. *The Detroit River, Michigan: an Ecological Profile.* U.S. Fish and Wildlife Service Biological Report 85(7.17).

Manny, B.A., Nichols, S.J. and D.W. Schoesser. 1991. Heavy metals in aquatic macrophytes drifting in a large river. *Hydrobiologia* **219**, 333-344.

Maynard, L. and D. Wilcox. 1997. *State of the Lakes Ecosystem Conference 1996 Background Paper: Coastal Wetlands.* Canadian Wildlife Service, Environment Canada Guelph, Ontario & Great Lakes Science Centre, National Biological Service, Ann Arbor, Michigan.

McCullough, G.B. 1985. Wetland threats and losses in Lake St. Clair. In: Prince, H.P. and F.M. D'Itri (eds.). *Coastal Wetlands.* pp. 201-208. Lewis Publishing. Co. Chelsea, Michigan.

McCullough, G.B. 1982. Wetland losses in Lake St. Clair and Lake Ontario. In: Champagne, E. (ed.), *Proceedings of the Ontario Wetlands Conference*. Federation of Ontario Naturalists, Don Mills, Ontario.

Meredith, G.N. and J.J. Houston. 1988. Status of the longear sunfish (*Lepomis megalotis*) in Canada. *Canadian Field Naturalist* **102**, 277-285.

Minns, C.K., J.D. Meisner, J.E. Moore, L.D. Greig, and R.G. Randall. 1995. *Defensible Methods for Pre- and Post- Development Assessment of Fish Habitat in the Great Lakes. I. A Prototype Methodology for Headlands and Offshore Shuchues.* Canadian Manuscript Report of Fisheries and Aquatic Sciences No. 2328.

Morris, R. 1983. *Erosion and Sedimentation Processes Throughout the Oshawa Second Marsh Watershed.* Lands Directorate, Environment Canada, Burlington, Ontario.

Natural Heritage Information Centre. 1995. Rare communities of Ontario: Great Lakes coastal meadow marshes. *Ontario Natural Heritage Information Centre Newsletter* **2**, 4-5.

Nichols, S.J., B.A. Manny, D.W. Schloesser, and T.A. Edsall. 1991. Heavy metal concentrations of sediments in the upper connecting channels of the Great Lakes. *Hydrobiologia* **219**, 307-315.

New York State Department of Environmental Conservation. 1994. *Niagara River Remedial Action Plan, Summary.* Division of Water, New York State Department of Environmental Conservation, Buffalo, N.Y.

Oldham, M.J. 1996. *Natural Heritage Resources of Ontario: Rare Vascular Plants.* Natural Heritage Information Centre, Ontario Ministry of Natural Resources, Peterborough, Ontario.

Oldham, M.J. 1996. Personal communication. Natural Heritage Information Centre, Ontario Ministry of Natural Resources, Peterborough, Ontario.

Ontario Ministry of Natural Resources. 1993a. *Ontario Wetland Evaluation System. Southern Manual. Third Edition.* Ontario Ministry of Natural Resources, Peterborough, Ontario.

Ontario Ministry of Natural Resources. 1993b. *Ontario Wetland Evaluation System. Northern Manual. First Edition.* Ontario Ministry of Natural Resources, Peterborough, Ontario.

Ontario Ministry of Natural Resources. 1991. *Sault Ste. Marie District Shoreline Management Plan, 1991-2000.* Ontario Ministry of Natural Resources, Sault Ste. Marie, Ontario.

Ontario Ministry of Natural Resources and Environment Canada. 1984. *An Evaluation System for Wetlands of Ontario - South of the Precambrian Shield. Second Edition.* Wildlife Branch, Outdoor Recreation Group, OMNR, and Canadian Wildlife Service, Ontario Region, Environmental Conservation Service, Environment Canada.

Ontario Ministry of Natural Resources and Environment Canada. 1983. *An Evaluation System for Wetlands of Ontario South of the Precambrian Shield. First Edition.* Wildlife Branch, Outdoor Recreation Group, Ontario Ministry of Natural Resources, and Canadian Wildlife Service, Ontario Region, Environmental Conservation Service, Environment Canada.

Ontario Ministry of the Environment and Michigan Department of Natural Resources. 1991. *The St. Clair River Area of Concern Environmental Conditions and Problem Definitions.* Stage 1 Remedial Action Plan.

Ontario Ministry of the Environment, Environment Canada, Department of Fisheries and Oceans, Ontario Ministry of Natural Resources and Ontario Ministry of Agriculture and Food. 1990. *Bay of Quinte Remedial Action Plan Stage I: Environmental Setting and Problem Definition.* Belleville, Ontario.

Painter, D.S., K.J. McCabe, and W.L. Simser. 1989. *Past and Present Limnological Conditions in Cootes Paradise Affecting Aquatic Vegetation.* Technical Bulletin No. 13. Royal Botanical Gardens, Hamilton, Ontario.

Parker, B. and P. McKee. 1984. Status of the spotted gar (*Lepisosteus oculatus*) in Canada. *Canadian Field Naturalist* **98**, 80-85.

Parker, B., P. McKee, and R.R. Campbell. 1987a. Status of the pugnose shiner (*Notropis anogenus*) in Canada. *Canadian Field Naturalist* **101**, 203-207.

Parker, B., P. McKee, and R.R. Campbell. 1987b. Status of the pugnose minnow (*Notropis emiliae*) in Canada. *Canadian Field Naturalist* **101**, 208-212.

Patch, S.P. and W.-D.N. Busch. 1984. *The St. Lawrence River-Past and Present: A Review of Historical Natural Resource Information and Habitat Changes in the International Section of the St. Lawrence River.* U.S. Fish and Wildlife Service, Cortland, N.Y.

PLUARG. 1978. *Environmental Management Strategy for the Great Lakes System.* International Reference Group on Great Lakes Pollution from Land Use Activities, International Joint Commission, Windsor, Ontario.

Prince, H.H., P.I. Padding, and R.W. Knapton. 1992. Waterfowl use of the Laurentian Great Lakes. *Journal of Great Lakes Research* **18**, 673-699.

Riley, J. and E. Snell. 1997. Remote Sensing and Wetlands: Workshop Proceedings and Analysis. 30 pp.

Richards, R.P. and D.B. Baker. 1993. *Trends in Nutrient and Suspended Sediment Concentrations in Lake Erie Tributaries, 1975-1990.* Journal of Great Lakes Research **19**, 200-211.

Ross, R.K. 1984. Migrant waterfowl use of major shorelines of eastern Ontario. In: Curtis, S.G., Dennis, D.G., Boyd, H. (eds.). *Waterfowl studies in Ontario, 1973-1981.* pp. 53-62. Occasional Paper No. 54, Canadian Wildlife Service, Environment Canada, Ottawa, Ontario.

Rutherford, L.A. 1979. *The Decline of Wetlands in Southern Ontario.* Unpublished B.E.S. Thesis. Department of Environmental Studies. University of Waterloo, Waterloo, Ontario.

Scott, W.B. and E.J. Crossman. 1973. *Freshwater fishes of Canada.* Fisheries Research Board of Canada Bulletin 184.

Selinger, W. 1996. Personal communication. Espanola Area Office, Ontario Ministry of Natural Resources, Espanola, Ontario.

Severn Sound Remedial Action Plan. 1993a. *Severn Sound Remedial Action Plan Stage 2 Report.*

Severn Sound Remedial Action Plan. 1993b. *An Interim Fish Habitat Management Plan for the Severn Sound.*

Simser, W.L. 1979. *Changes in Aquatic Biota of Cootes Paradise Marsh.* Technical Bulletin No. 12, Royal Botanical Gardens, Hamilton, Ontario.

Smith, P. 1987. *Towards the Protection of Great Lakes Natural Heritage Areas.* Technical Paper, Heritage Resources Centre, University of Waterloo, Waterloo, Ontario.

Smith, P.G.R., V. Glooschenko, and D.A. Hagen. 1991. Coastal wetlands of the three Canadian Great Lakes: inventory, current conservation initiatives and patterns of variation. *Canadian Journal of Fisheries and Aquatic Sciences* **48**, 1581- 1594.

Snider, B. 1996. Personal communication. Peterborough District Office, Ontario Ministry of Natural Resources, Peterborough, Ontario.

St. Clair River Remedial Action Plan. 1995. *The St. Clair River Area of Concern: Water Use Goals, Remedial Measures and Implementation Strategy.* Ontario Ministry of Environment and Energy, Michigan Department of Natural Resources. Sarnia, Ontario and Lansing, Michigan.

Stephenson, T.D. 1990. Fish reproductive utilization of coastal marshes of Lake Ontario near Toronto. *Journal of Great Lakes Research* **16**, 71-81.

Suke, S. 1996. Personal communication. Lakehead Region Conservation Authority, Thunder Bay, Ontario.

Sutherland, D.A. 1996. Personal communication. Natural Heritage Information Centre, Ontario Ministry of Natural Resources, Peterborough, Ontario.

Sutherland, D.A. 1994. *Natural Heritage Resources of Ontario: Freshwater Fishes.* Natural Heritage Information Centre. Peterborough, Ontario.

Thompson, R. 1996. Personal communication. Long Point Area Office, Ontario Ministry of Natural Resources, Simcoe, Ontario.

Whillans, T.H. 1982. Changes in marsh area along the Canadian shore of Lake Ontario. *Journal of Great Lakes Research* **8**, 570-577.

Whillans, T.H., R.C. Smardon and W.-D.N. Busch. 1992. *Status of Lake Ontario Wetlands.* Great Lakes Research Consortium.

Williams, D.C. and J.G. Lyon. 1991. Use of a Geographic System Database to Measure and Evaluate Wetland Changes in the St. Marys River, Michigan. *Hydrobiologia* **219**, 83-95.

Wilcox, D.A., J.E. Meeker, and J. Elias. 1993. *Impacts of Water-level Regulation on Wetlands of the Great Lakes.* Phase 2 Report to Working Committee 2, International Joint Commission, Great Lakes Water Levels Reference Study.

Wilcox, S.A. 1995. *Bird and Nature Conservation Planning: A Financial and Human Ecological Approach, the Case of Long Point.* M.A. Thesis. School of Urban and Regional Planning. University of Waterloo, Waterloo, Ontario.

Appendix A: 1996 Ontario Ministry of Natural Resources District/Area Office Survey

(sent to all Ministry of Natural Resources District/Area Offices bordering the Great Lakes in May 1996)

COASTAL WETLAND SURVEY

District/Area: ____________________

Contact: ________________________

PART I. **Complete the following table for each evaluated coastal wetland in your district/area. Please make additional copies of the table if you need more space.**

Wetland Name	UTM Grid Reference	Wetland Characteristics						PSW[4] (Y/N)	Evaluation Date (Year)	Ownership			Protective Designation[5]	Is the wetland getting smaller? (loss in ha if known)	Major Stresses[6]
		Single wetland area or complex	Total wetland area (ha)	Type(s)[1]	FA[2]	Site types[3]	FA			FA in trust	FA public	FA private			

[1] 1=swamp 2=marsh 3=bog 4=fen

[2] FA=fractional area

[3] 1= isolated 2=palustrine (permanent or intermittent flow) 3=riverine 4= riverine (at rivermouth) 5=lacustrine (at rivermouth) 6= lacustrine (on enclosed bay with barrier beach) 7=lacustrine (exposed to lake)

[4] PSW=Provincially Significant Wetland

[5] 1=Ramsar Site 2=National Park 3=National Wildlife Area 4=Provincial Park 5=Conservation Area 6=Provincial Wildlife Area 7=Migratory Bird Sanctuary 8=Federation of Ontario Naturalists and Local Nature Club Properties 9=ANSI 10=Other (please specify)

[6] 1=water level regulation 2=shoreline modification 3=dike construction 4=road construction 5=water quality (5a=chemical contamination, 5b=nutrient loading, 5c=sediment loading) 6=urban encroachment (filling/draining) 7=agricultural encroachment (clearing/draining) 8=dredging/channelization (e.g. marina development) 9=non-indigenous species 10=water taking 11=other (please specify)

Appendix A (cont.'d): 1996 Ontario Ministry of Natural Resources District/Area Office Survey

PART II:
If the following information is readily available, please complete the following questions:

1a. Approximately how many coastal wetlands (>2 ha) are left in your district/area to be evaluated (please identify the names, size and type if known)?

b. How many coastal wetlands left to be evaluated may be provincially significant wetlands?

2. Wetlands <2 ha are not normally evaluated using the Ontario Wetland Evaluation System, unless they are part of a wetland complex. How many coastal wetlands do you have that are <2 ha and not part of a wetland complex (ballpark number would be fine)?

3. Generally, what are the main stresses on coastal wetlands in your area/district (see table footer above for the categories of stresses)?

Appendix B: A Guide to Appendices E, G, I, K, M, O: Evaluated Coastal Wetlands

* indicates information taken from Wetland Evaluations.

COLUMN/FIELD	INFORMATION / INTERPRETATION
MAP ID:	A number that identifies the evaluated wetland on the map. Numbers increase from east to west along the shoreline. Numbers are not unique across maps/figures.
WETLAND NAME*:	The official name identified by the local Ontario Ministry of Natural Resources (OMNR) District and recorded on the Wetland Evaluation. The Natural Heritage Information Centre (NHIC) has provided each evaluated wetland with a Natural Area number. This number is in brackets following the wetland name.
OMNR DISTRICT:	The OMNR District Office responsible for maintaining and updating the Wetland Evaluation.
UTM ZONE: (Universal Trans Mercator)	A two digit numeric character that identifies zones that run north and south between lines of longitude. Within each zone a metric mercator grid is defined that enables you to describe the geographical location of any wetland. Ontario consists of three numbered zones, 16, 17 and 18.
MAP SHEET:	An alphanumeric character that identifies the national topographic map that shows the UTM grid where the wetland is located.
(UTM) EASTING*:	A six-digit number that describes the point location of a wetland on the north-south lines of the UTM grid on the topographic map.
(UTM) NORTHING*:	A seven-digit number that describes the point location along the east-west lines of the UTM grid on the topographic map.
SINGLE/COMPLEX*:	A character that identifies whether the evaluation is of a single wetland (S) or a complex (C) of wetlands. Single refers to an evaluation of one wetland unit with distinct boundaries. Within a single wetland area (i.e., contiguous wetland) one may find very different ecological circumstances for example, an open water marsh, a spring fed swamp forest, a shoreline fen, etc. Complex refers to an evaluation of a group of wetlands. Major functional discontinuities such as uplands or open water lakes may subdivide the area into a number of distinctive wetland units, yet the entire suite of wetland units is evaluated as a single complex. Wetlands are grouped when delineation of the wetland units into individually recognized wetlands is not ecologically or functionally sound. For example, the St. Clair Marshes is a complex made up of 20 individual wetlands.
AREA*:	Size of the evaluated wetland in hectares.
WETLAND TYPES*:	Four types of wetland ecosystems are identified based on main vegetation composition: swamp, marsh, bog and fen. Wetland types differ in their typical form, in the number and kinds of plant and animal species they support and in their relative rates of primary productivity. Any particular wetland may be comprised of one or more wetland types. The definitions come from the Ontario Wetland Evaluation System: Southern Manual (OMNR, 1993). **SWAMP:** The proportion of the wetland that is swamp. Swamps are wooded wetlands with 25% cover or more of trees or tall shrubs. In swamps, standing to gently flowing waters occur seasonally or persist for long periods on the surface. Swamps include both forest and thicket swamps. **MARSH:** The proportion of the wetland that is marsh. Marshes are areas

	periodically inundated with standing or slowly moving water, and/or permanently inundated characterized by robust emergents, and to a lesser extent, anchored floating plants and submergents. **BOG:** The proportion of the wetland that is bog. Bogs are peat-covered areas or peat-filled depressions with a high water table and a surface carpet of mosses, chiefly *Sphagnum* spp. **FEN:** The proportion of the wetland that is fen. Fens are peatlands characterized by surface layers of poorly to moderately decomposed peat, often with well-decomposed peat near the base. They are covered by a dominant component of sedges, although grasses and reeds may be associated in local pools.
SITE TYPES*:	The physiographic position of a wetland in the landscape defines site type. There are four fundamentally different site types: isolated, palustrine, riverine and lacustrine; riverine and lacustrine are further subdivided because of the location of a wetland on a lake or river has a bearing on nutrient concentrations of the water and hence upon productivity. The definitions come from the Ontario Wetland Evaluation System: Southern Manual (OMNR, 1993). **ISOLATED:** The proportion of the wetland that has no surface outflow. **PALUSTRINE:** The proportion of the wetland that either has absent or intermittent inflow and either intermittent or permanent outflow. **RIVERINE:** The proportion of the wetland that includes the channel of continuously moving water to the 2 m depth, as well as adjacent wetlands and normal flood plains of rivers and permanent streams. **RIVERINE RIVERMOUTH:** The proportion of a wetland that has formed where a river stream enters one of Ontario's five large rivers (Ottawa, St.Lawrence, St. Clair, Detroit and Niagara). **LACUSTRINE RIVERMOUTH:** The proportion of a wetland that has formed where a river or a river or stream enters a lake. **LACUSTRINE BAY:** The proportion of a wetland that is separated from a lake by a barrier beach by which lake waters may from time to time be sealed off. **LAKE LACUSTRINE:** The proportion of a wetland on a lake where a barrier beach is not present, and the wetland is exposed to lake forces.
CLASS*:	Prior to 1993, wetlands were assigned a class from 1 to 7 based on the number of points scored in the evaluation. Wetlands classed as 1 or 2 were considered Provincially Significant, wetlands classed 3 were considered Regionally Significant. Post 1993, class designations no longer exist; instead wetlands are classified as either Provincially Significant or Other (Not Provincially Significant). Following District practices, formerly Regionally Significant wetlands are now considered Provincially Significant, in accordance with their score, under the new system. PSW refers to a Provincially Significant Wetland and NPSW refers to a Non Provincially Significant Wetland. Numbers from 1 to 7 in the Appendices refer to the class prior to 1993.
PROVINCIALLY SIGNIFICANT WETLAND:	T (True) indicates that the wetland is a Provincially Significant Wetland (PSW), and F (False) indicates that the wetland is not a PSW.
YEAR OF EVALUATION DATA:	The year of evaluation from which the information in the table was recorded. The most recent evaluation was used wherever possible. "DU" after the year of evaluation indicates a desktop update edition of the evaluation. In the case of desktop updates, previous edition evaluation information is transferred to a new edition evaluation without new

	fieldwork or research. Thus a 1996DU evaluation does not have the same currency of information that an evaluation actually carried out in 1996 would. As well, evaluations carried out over the years fall under three different editions of the Wetland Evaluation manual, and thus represent a diversity of data quality and quantity.
PUBLIC OWNERSHIP*:	An estimate of the proportion of public land contained in the wetland.
PRIVATE OWNERSHIP*:	An estimate of the proportion of private land contained in the wetland.
CORRESPONDING NATURAL AREA:	Natural areas that overlap or are adjacent to an evaluated wetland. The natural area number and classification assigned by the NHIC is provided. The existence of a corresponding natural areas is shown on the map using a red square around the wetland MAP ID. The following codes are used in the table to identify Natural Areas:

Code	NHIC Designated Natural Area Description
ANSI-ES-P	Provincial Earth Science Area of Natural and Scientific Interest
ANSI-LS-C	Candidate Life Science Area of Natural and Scientific Interest
ANSI-LS-P	Provincial Life Science Area of Natural and Scientific Interest
ANSI-LS-R	Regional Life Science Area of Natural and Scientific Interest
CAA	Conservation Authority Area
ES	Earth Science Site
ESA	Environmentally Sensitive Area
IBP	International Biological Program Site
LS	Life Science Site
MACRO	Macro Site (Unofficial NHIC Designation)
MP	Municipal Park
NGO	Non-Government Organization Nature Reserve
NP	National Park
NWA	National Wildlife Area
OHF	Ontario Heritage Foundation Trust Property
PP	Provincial Park
PP-NE	Provincial Park- Natural Environment
PP-NEZ	Provincial Park- Natural Environment Zone
PP-NR	Provincial Park- Nature Reserve
PP-NRZ	Provincial Park- Nature Reserve Zone
PP-R	Provincial Park- Recreational
PWA	Provincial Wildlife Area

OTHER COMMENTS:	The comment field contains additional information about a wetland including those provided by OMNR biologists in the OMNR District/Area Office Survey. The codes used in the Comments field are:

Code	Comment Description
CAA	Conservation Authority Owned Land
DU	Ducks Unlimited
EHJV	Eastern Habitat Joint Venture
ESA	Environmentally Sensitive Area
FON	Federation of Ontario Naturalists Property

MBS	Migratory Bird Sanctuary
MNR	Ministry of Natural Resources
MP	Municipal Property
NP	National Park
NR	Nature Reserve
PP	Provincial Park
PWA	Provincial Wildlife Area
RAMSAR	Ramsar Convention – Wetlands of International Importance (especially for waterfowl habitat)
SCPC	St. Clair Parks Commission
SLPC	St. Lawrence Parks Commission

LOSS:	A T/F character indicates whether loss of wetland area was reported by OMNR district biologists in the OMNR District/Area Office Survey. T (True) means that recent loss was reported and F (False) means that recent loss was not reported. A blank means that no information was provided.
STRESSES:	A numeric character that indicates what stress(es) were reported by OMNR district biologists in the OMNR District/Area Office Survey. A blank means that no information was provided. The numeric characters and corresponding stresses are: 1. water level regulation 2. shoreline modification 3. dike construction 4. road construction 5. water quality (5a= chemical contamination, 5b=nutrient loading, 5c= sediment loading) 6. urban encroachment (filling/draining) 7. agricultural encroachment (clearing/draining) 8. dredging/channelization (e.g. marina development) 9. non-indigenous species 10. water taking 11. other
PROVINCIALLY SIGNIFICANT SPECIES OR VEGETATION COMMUNITIES*:	A character (Y=yes) indicates whether a provincially significant vegetation community, or plant, bird, reptile/amphibian, mammal, fish or butterfly species (S-Rank S1, S2, S3 or S3S4 as designated by the NHIC as of February 1999) has been reported in the Wetland Evaluation. Blank = not reported.

Appendix C: A Guide to Appendices F, H, J, L, N: Natural Areas, Environmental Sensitivity Atlas Wetlands and Other Wetlands with an Unevaluated Coastal Wetland Component

COLUMN/FIELD	INTERPRETATION/EXPLANATION
MAP ID:	An alphanumeric character that identifies the Natural Area wetland (N#), Environmental Sensitivity Atlas wetland (A#) or other reported wetland (O#) on the map. Characters are not unique across maps.
AREA NAME:	The name assigned to the Natural Area, Environmental Sensitivity Atlas or Other wetland illustrated on the map. The Natural Heritage Information Centre (NHIC) has provided each Natural Area with a number. This number is in brackets following the area name and classification. The Natural Area classification is assigned by the NHIC. The following codes are used in the tables to identify Natural Areas:

Code	NHIC Designated Natural Area Description
ANSI-ES-P	Provincial Earth Science Area of Natural and Scientific Interest
ANSI-LS-C	Candidate Life Science Area of Natural and Scientific Interest
ANSI-LS-P	Provincial Life Science Area of Natural and Scientific Interest
ANSI-LS-R	Regional Life Science Area of Natural and Scientific Interest
CAA	Conservation Authority Area
ES	Earth Science Site
ESA	Environmentally Sensitive Area
IBP	International Biological Program Site
LS	Life Science Site
MACRO	Macro Site (Unofficial NHIC Designation)
MP	Municipal Park
NGO	Non-Government Organization
NP	National Park
NWA	National Wildlife Area
OHF	Ontario Heritage Foundation Trust Property
PP	Provincial Park
PP-NE	Provincial Park- Natural Environment
PP-NEZ	Provincial Park- Natural Environment Zone
PP-NR	Provincial Park- Nature Reserve
PP-NRZ	Provincial Park- Nature Reserve Zone
PP-R	Provincial Park- Recreational
PWA	Provincial Wildlife Area

MNR DISTRICT:	The Ontario Ministry of Natural Resource (OMNR) District Office where wetland is located.
UTM ZONE: (Universal Transverse	A two digit numeric character that identifies zones that run north and south between lines of longitude. Within each zone a metric mercator grid is

Mercator)	defined that enables you to describe the geographical location of any wetland. *Ontario consists of three numbered zones, 16, 17 and 18.*
MAP SHEET:	An alphanumeric character that identifies the national topographic map that shows the UTM grid where the wetland is located.
(UTM) EASTING:	A six-digit number that describes the point location along the north-south lines of the UTM grid on the topographic map.
(UTM) NORTHING:	A seven-digit number that describes the point location along the east-west lines of the UTM grid on the topographic map.
DESCRIPTION:	Where available, a brief summary of information that describes the unevaluated wetland. The descriptions of wetlands from the Environmental Sensitivity Atlas (ESA) include the corresponding map name and number and a correlating Natural Area # and name if applicable. The Great Lakes Environmental Sensitivity Atlases provide a visual reference intended to assist in the preparation and response to spills of oil and other hazardous materials into the Great Lakes and St. Lawrence River basin. The information contained in the Atlases was the most up-to-date available, when compiled between 1991 and 1994. The Environmental Sensitivity Atlases are broken down into seven sub-regions: Lake Superior, Lake Huron / Georgian Bay, St. Marys River, St. Clair River / Detroit River / Lake St. Clair, Lake Erie / Niagara River / Welland Canal, Lake Ontario, and St. Lawrence River. A shoreline classification scheme utilized for spill response on the Great Lakes and connecting channels identifies 19 different shoreline habitats. Data sources included species surveys, OMNR wetland evaluations, aerial video photography of the shoreline and ground surveys. A combination of geomorphologists, biologists, mapping and GIS experts, ecologists, spill response experts and others contributed to the development of the Environmental Sensitivity Atlases. Environmental Sensitivity Atlas information categories include shoreline habitats (including wetlands), biological resources, human-use resources and countermeasures. Each shoreline feature is mapped with a symbol that corresponds to a generic atlas legend.
OTHER CORRESPONDING NATURAL AREAS:	Other Natural Areas, taken from the NHIC Natural Areas Database, that overlap or are adjacent to the wetland area. Natural Areas are identified with a number and area name or classification. Classification codes are explained above in AREA NAME. If none exist, N/A is used.
SIGNIFICANT SPECIES OR VEGETATION COMMUNITIES:	A character (Y = yes) that indicates whether a provincially significant vegetation community, or plant, bird, reptile/amphibian, mammal, fish or butterfly species (S-Rank S1, S2, S3 or S3S4 as designated by the NHIC as of February, 1999) has been reported for the area.

Appendix D: A Guide to Appendices P to U: Significant Species Reported in Ontario's Great Lakes Coastal Wetlands

COLUMN/FIELD	INTERPRETATION/EXPLANATION
COMMON NAME:	The nomenclature used follows the Natural Heritage Information Centre (NHIC) current standard (February, 1999).
SCIENTIFIC NAME:	The nomenclature used follows the NHIC current standard (February, 1999). "***"indicates an Ontario species only found in coastal wetlands
NHIC S-RANK:	Provincial (or Sub-national) ranks are used by the Natural Heritage Information Centre to set protection priorities for rare species and natural communities. These ranks are not legal designations. The most important factors considered in assigning provincial ranks are the total numbers of known, extant sites in Ontario, and the degree to which they are potentially or actively threatened with destruction. Other criteria include the number of known populations considered to be securely protected, the size of the various populations, and the ability of the taxon to persist at its known sites. The taxonomic distinctness of each taxon has also been considered. Hybrids, introduced species and taxonomically dubious species, subspecies and varieties have generally not been included. By comparing the global and provincial ranks, the status, rarity, and the urgency of conservation needs can be ascertained. Provincial ranks have been assigned using the best available scientific information, and have been reviewed by a group of experts on the flora and fauna of Ontario. The NHIC evaluates provincial ranks on a continual basis . Species with ranks of S1, S2, S3, S3S4 (as of February 1999) are referred to in the text as provincially significant. In Appendices P to U, species with these ranks that are found in coastal wetlands are presented as significant species. **S1 –** Extremely rare in Ontario; usually 5 or fewer occurrences in the province or very few remaining individuals; often especially vulnerable to extirpation. **S2 –** Very rare in Ontario; usually between 5 and 20 occurrences in the province or with many individuals in fewer occurrences; often susceptible to extirpation. **S3 –** Rare to uncommon in Ontario; usually between 2 and 100 occurrences in the province; may have fewer occurrences, but with a large number of individuals in some populations; may be susceptible to large-scale disturbances. **S4 –** Common and apparently secure in Ontario; usually with more than 100 occurrences in the province. **S5-** Very common and demonstrably secure in Ontario. **SH –** Historically known from Ontario, but not verified recently (typically not recorded in the province in the last 20 years); however suitable habitat is thought to be still present in the province and there is reasonable expectation that the species may be rediscovered. **SR –** Reported for Ontario, but without persuasive documentation which would provide a basis for either accepting or rejecting the report. **SRF –** Reported falsely from Ontario. **SX –** Apparently extirpated from Ontario, with little likelihood for rediscovery. Typically not seen in the province for many decades, despite searches at known historic sites. **SE –** Exotic; not believed to be a native component of Ontario's flora or fauna. **SU –** Unranked. SU species are possibly rare in Ontario, but there is insufficient information available to assign a more accurate rank. **? –** Following a rank indicates some degree of uncertainty.

	B following a bird rank indicates breeding. **N** following a bird rank indicates non-breeding. **Rank ranges,** e.g. S3S4 indicates that the Ontario rank is either S3 or S4, but that the information currently available is insufficient to determine which rank applies.
COSEWIC:	Status assigned by the Committee on the Status of Endangered Wildlife in Canada, following the most recent COSEWIC list (November 2002) (www.cosewic.gc.ca). **EXP – Extirpated.** A species no longer existing in the wild in Canada, but occurring elsewhere. **END – Endangered.** A species facing imminent extirpation or extinction. **THR – Threatened.** A species likely to become endangered if limiting factors are not reversed. **SC – Special Concern*.** A species of special concern because of characteristics that make it particularly sensitive to human activities or natural events. **NAR – Not at Risk**.** A species that has been evaluated and found to be not at risk. **DD – Data Deficient***.** A species for which there is insufficient scientific information to support status designation. * Formerly described as "Vulnerable" from 1990 to 1999, or "Rare" prior to 1990. ** Formerly described as "Not in Any Category", or "No Designation Required." *** Formerly described as "Indeterminate" from 1994 to 1999 or "ISIBD" (insufficient scientific information on which to base a designation) prior to 1994.
COSSARO:	Status assigned by the Ontario Ministry of Natural Resources (September 2002) based on recommendations of a Ministry technical committee called the Committee on the Status of Species at Risk in Ontario (COSSARO). Endangered species (END) are protected under the province's *Endangered Species Act*. (www.mnr.gov.on.ca/mnr/VTEEElist_2002_eng.pdf) **END – Endangered.** Any indigenous species of fauna or flora that, on the basis of the best available scientific evidence, is indicated to be threatened with immediate extinction throughout all or a significant portion of its Ontario range. **END-R** – Endangered Species regulated under the provincial *Endangered Species Act*. **THR – Threatened.** Any indigenous species of fauna or flora, that on the basis of the best available scientific evidence, is indicated to be experiencing a definite non-cyclical decline throughout all or a major portion of its Ontario range, and that is likely to become an endangered species if the factors responsible for the decline continue unabated. **VUL – Vulnerable**. Any indigenous species of fauna or flora that is represented in Ontario by small but relatively stable populations, and/or that occurs sporadically, or in a very restricted area of Ontario, or at the fringe of its range, and that should be monitored periodically for evidence of a possible decline. Vulnerable should now be used for species formerly designated as Rare by MNR. **NIAC** – Not in Any category. **IND** – Indeterminate.
LAKE/ CONNECTING CHANNEL:	The main body of water associated with the location of the coastal wetland for which the occurrence of a significant species has been reported. See DATA SOURCES for occurrence information sources.
BREEDING (CONFIRMED/ PROBABLE):	**For significant bird species only (Appendix S).** The main location of breeding activity, confirmed or probable, for the occurrence of a significant bird species, or other breeding status information.

NON-BREEDING:	**For significant bird species only (Appendix S).** The main location of non-breeding activity, for the occurrence of a significant bird species, or other non-breeding status information.
HABITAT DESCRIPTION:	General habitat condition(s) preferred by species.
COMMENTS:	**For unconfirmed records of significant vascular plants only (Appendix U).** Further information on species habitat, species status, and species taxonomy, clarifying why a record may be unconfirmed according to the experts consulted.
DATA SOURCES:	**Vascular Plant Species** • **Species status and occurrence verification:** W. Bakowsky, pers. com., 1996; J. Jalava, pers. com., 1996; M.J. Oldham, pers. com., 1996; D. Sutherland, pers. com., 1996 • **Supporting Studies for species occurrence location:** OMNR Wetland Evaluations; Oldham, 1996
	Fish Species • **Species status and occurrence verification:** A. Dextrase, pers. com., 1996; D. Sutherland, pers. com., 1996 • **Supporting studies for species occurrence location or habitat descriptions:** Jude and Pappas, 1992; Lane *et al*, 1996; Lee *et al.*, 1980; Mandrak and Crossman, 1993; Mandrak and Crossman, 1992; Meredith and Houston, 1988; Minns *et al.*, 1995; Parker *et al.*, 1987a, b; Parker and McKee, 1984; Scott and Crossman, 1973 • no fish species identified in OMNR wetland evaluations
	Reptile and Amphibian Species • **Species status and occurrence verification:** J. Jalava, pers. com., 1996; M.J. Oldham, pers. com., 1996; D. Sutherland, pers. com., 1996 • **Supporting studies for species occurrence location:** OMNR Wetland Evaluations • **Supporting studies for habitat descriptions:** Harding, 1997
	Bird Species • **Species status and occurrence verification:** W. Bakowsky, pers. com., 1996; M. Cadman, pers. com., 1996; J. Jalava, pers. com., 1996; M.J. Oldham, pers. com., 1996; D. Sutherland, pers. com., 1996; • **Supporting studies for species occurrence location:** Prince *et al.*, 1992; OMNR Wetland Evaluations; Cadman *et al.*, 1987 • **Supporting studies for habitat descriptions:** Cadman *et al.*, 1987
	Lepidoptera Species • **Species status and occurrence verification:** D. Sutherland, pers. com., 1996 • **Supporting studies for species occurrence location:** OMNR Wetland Evaluations
	Unconfirmed Significant Vascular Plant Species • **Species status and occurrence verification:** W. Bakowsky, pers. com., 1996; J. Jalava, pers. com., 1996; M.J. Oldham, pers. com., 1996; D. Sutherland, pers. com., 1996; • **Supporting Studies for species occurrence location:** OMNR Wetland Evaluations; Oldham, 1996

Appendix E: Lake Superior: Evaluated Coastal Wetlands

									Wetland Type				Site Type																Significant Species						
MAP ID (Figure 1)	WETLAND NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1: 50 000)	EASTING	NORTHING	SINGLE/COMPLEX	AREA	SWAMP	MARSH	BOG	FEN	ISOLATED	PALUSTRINE	RIVERINE	RIVERINE RIVERMOUTH	LACUSTRINE RIVERMOUTH	LACUSTRINE BAY	LAKE LACUSTRINE	CLASS	PROVINCIALLY SIGNIFICANT WETLAND	YEAR OF EVALUATION DATA	PUBLIC OWNERSHIP	PRIVATE OWNERSHIP	CORRESPONDING_NATURAL AREA	OTHER COMMENTS	LOSS	STRESSES	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
1	Hurkett Cove (7128)	Thunder Bay	16	52A/16	390300	5409500	S	55.00		1.000							0.050	0.950		6	F	1988	1.000			CAA	F								
2	Northern Wood Preservers Marsh (7132)	Thunder Bay	16	52A/6	335500	5365300	C	4.56	0.040	0.960				0.600					0.400	NPSW	F	1992		1.000			T	2,6							
3	Neebing Marsh (7129)	Thunder Bay	16	52A/6	335800	5363200	S	101.10	0.400	0.490		0.100		0.200			0.200	0.600		NPSW	F	1995	0.950	0.050			F	8			Y				
4	McKellar Island Marsh (7127)	Thunder Bay	16	52A/6	336000	5361900	S	11.50		1.00									1.000	7	F	1984		1.000			F	11			Y				
5	Mission Marsh (7124)	Thunder Bay	16	52A/6	336000	5359000	S	160.20	0.430	0.570									1.000	PSW	T	1993	both	both			F	11			Y			Y	
6	Chippewa Marsh (7123)	Thunder Bay	16	52A/6	335800	5356100	S	23.00		0.900		0.100					1.000			2	T	1983	1.000		4979 LS	Protected by city	F	5b,11			Y				
7	Sturgeon Bay (North) ((7134)	Thunder Bay	16	52A/3	328800	5341300	S	72.00	0.960	0.040								0.960	0.040	NPSW	F	1994	0.960	0.040			T								
8	Sturgeon Bay (SouthWest) (7133)	Thunder Bay	16	52A/3	328200	5339800	C	42.00	0.860	0.140								0.860	0.140	NPSW	F	1994	0.140	0.860			F								
9	Caldwell Lake (7696)	Thunder Bay	16	52A/3	323200	5330900	S	166.20	0.190	0.620	0.060	0.120		0.260	0.020				0.072	PSW	T	1995	1.000						Y						
10	Cloud Bay (7126)	Thunder Bay	16	52A/3	318000	5328000	S	16.00	0.050	0.950							0.400	0.600		5	F	1988	1.000			CAA	F	4,11							
11	Pine Bay (7130)	Thunder Bay	16	52A/3;5 2A/4	311900	5322900	S	429.00	0.850	0.077	0.073				0.600		0.100		0.300	NPSW	F	1995	0.200	0.800		CAA	F	4,11	Y						
				Total Wetland Area				**1080.56**																											

Appendix F: Lake Superior: Natural Areas, Environmental Sensitivity Atlas Wetlands and Other Wetlands with an Unevaluated Coastal Wetland Component

									Significant Species						
MAP ID	AREA NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:25 000)	EASTING	NORTHING	DESCRIPTION	OTHER CORRESPONDING NATURAL AREAS	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
Natural Area Wetlands															
N1	Shore Ridges Conservation Area ANSI-LS-C (4844)	Sault Ste. Marie	16	41K/9	689000	5153500	-array of wetland community types: swamp, treed fen	N/A		Y					
N2	Batchewana Bay ANSI-LS-R (366)	Sault Ste. Marie	16	41K/15	689000	5198800	-deep & shallow shoreline marshes; behind marshes is a beach ridge swale complex with gramminoid fens	3652 ES; 3015 PP-R		Y	Y				
N3	Batchewana Island Shoreline Marshes LS (3647)	Sault Ste. Marie	16	41N/2	692000	5196000	-marshes	N/A							
N4	Michipocoten Post PP (3141)	Wawa	16	41N/15	661336	5310157		N/A							
N5	Gravel River PNR (3074)	Nipigon	16	42D/13	445050	5418244	-birds foot delta; swamps with cedar and alder thickets; shallow marsh and beach communities	N/A							
N6	Nipigon River Marshes LS (8123)	Nipigon	16	52H/1	408000	5428000	-marshes and submerged aquatic communities; also gramminoid marshes with fen affinities	8125 LS							
Environmental Sensitivity Atlas Wetlands															
A1	Goulais Bay (Goulais River Wetland complex)	Sault Ste. Marie	16	41K/8	695000	5176500	-diverse wetland communities; marshes and fens	3653 ES; 4514 IBP							
A2	Sand Bay	Sault Ste. Marie	16	41K/15	687400	5180300		N/A							
A3	Bluewater Bay	Sault Ste. Marie	16	41K/15	687200	5187200		N/A							
A4	Marlettes Bay	Sault Ste. Marie	16	41K/15	688700	5188900		N/A							
A5	Black Bay Peatlands	Nipigon	16	52A/16	308217	5322061	-bog and fen wetland communities	217 LS							
A6	North of Moose Bay	Thunder Bay	16	52A/10	385500	5392400		N/A							
A7	Blende Rivermouth Wetland	Thunder Bay	16	52A/10	369800	5381200		N/A							

Appendix G: Lake Huron and St. Marys River: Evaluated Coastal Wetlands

									Wetland Type				Site Type																Significant Species						
MAP ID (Figure 2)	WETLAND NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:50 000)	EASTING	NORTHING	SINGLE/COMPLEX	AREA	SWAMP	MARSH	BOG	FEN	ISOLATED	PALUSTRINE	RIVERINE	RIVERINE RIVERMOUTH	LACUSTRINE RIVERMOUTH	LACUSTRINE BAY	LAKE LACUSTRINE	CLASS	PROVINCIALLY SIGNIFICANT WETLAND	YEAROF EVALUATION DATA	PUBLIC OWNERSHIP	PRIVATE OWNERSHIP	CORRESPONDING_NATURAL AREA	OTHER COMMENTS	LOSS	STRESSES	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
1	Kettle Point Marsh (7161)	Aylmer	17	40O/1	417000	4780500	S	53.760	0.400	0.600									1.000	2	T	1987	'undefined	'undefined	5 ANSI-LS-R; 334 MACRO; 5654 LS		F		Y	Y					
2	Port Franks (5647)	Aylmer	17	40P/4	426500	4786700	C	123.000	0.660	0.340			0.030		0.970					1	T	1985		1.000	57 ANSI-LS-P; 2001 FEDERAL LAND; 5650 LS		F		Y	Y	Y	Y			
3	Datars-Miller Swamp (7168)	Cambridge	17	40P/5	444000	4802000	C	76.700	1.000				0.340	0.660						7	F	1987		1.000			F	7							
4	Goderich 51D (7172)	Cambridge	17	40P/12	444800	4835000	S	53.600	0.970	0.030					1.000					5	F	1983	0.400	0.600		CAA	F	7							
5	Goderich 51A (7170)	Cambridge	17	40P/12	443700	4838900	S	5.030	1.000						1.000					6	F	1983		1.000			F	7							
6	Colborne 52D (7169)	Cambridge	17	40P/13	443800	4853300	S	28.000	1.000					1.000						7	F	1983		1.000			F	7							
7	Lorne Beach Swamp (7173)	Midhurst	17	41A/4	451500	4898500	C	28.000	0.814			0.186		1.000						5	F	1989	0.300	0.700											
8	Baie Du Dore (7175)	Midhurst	17	41A/5	455700	4909200	S	95.000	0.040	0.500		0.460					0.040		0.960	2	T	1984	0.500	0.500		ANSI	T	2,4,6,8		Y	Y				
9	Scott Point Wetland Complex (7422)	Midhurst	17	41A/15	457000	4911600	C	201.800	0.934	0.033		0.033	0.006	0.949					0.045	2	T	1990	0.040	0.960	1590 ANSI-LS-P		T	2,4,6,8		Y	Y	Y			
10	MacGregor Point Wetland Complex (7492)	Midhurst	17	41/A6; 41A/5	461000	4916000	C	420.200	0.820	0.040		0.140		0.900					0.100	PSW	T	1992	0.500	0.500	3128 PP-NE; 4598 IBP; 4614 ANSI-LS-R	Stewardship Agreement	T	2,4,6,8	Y	Y	Y	Y			Y
11	Chiefs Point (7419)	Midhurst	17	41A/11	478800	4948200	C	167.600	0.740	0.080		0.180		0.990					0.010	PSW	T	1992	0.010	0.990			T	2,4,6,8	Y	Y		Y			Y
12	Oliphant Wetland (177)	Midhurst	17	41A/11	478000	4953000	S	173.000	0.090	0.720		0.190	0.050	0.040					0.910	2	T	1984	0.750	0.250	4667 IBP		F	6,8,11	Y	Y	Y	Y			
13	Fishing Islands (7181)	Midhurst	17	41A/11; 41A/14	476000	4953000	C	168.500	0.089	0.885		0.026						0.117	0.883	2	T	1987	0.857	0.143	134 ANSI-LS-P		F	2	Y	Y	Y				
14	Red Bay Wetland Complex (7457)	Midhurst	17	41A/14	476600	4958600	C	353.100	0.900	0.080		0.020		0.930					0.070	PSW	T	1992	0.040	0.960			T	2,4,6,8	Y	Y	Y				
15	Howdenvale Bay (7185)	Midhurst	17	41A/14	476700	4962800	S	36.500	0.180	0.190		0.630							1.000	2	T	1984	0.200	0.800	175 ANSI-LS-P; 2077 NGO; 4668 IBP	FON Property	T	1,2,4,6,8	Y	Y		Y			
16	Sucker Creek (Owen Sound) (7191)	Midhurst	17	41A/14	475500	4964500	S	146.000	0.560	0.250		0.190			0.840		0.110		0.050	2	T	1984	0.890	0.110	174 ANSI-LS-P	PWA; NR	F	5c,5b		Y					
17	Pike Bay (7189)	Midhurst	17	41A/14	474000	4968000	C	41.600	0.349	0.484		0.167		0.292	0.035	0.014			0.659	6	T	1985	0.300	0.700	4363 LS		T	2,6,8	Y						
18	Black Creek Swamp (7176)	Midhurst	17	41A/14	471500	4980000	S	37.900	0.570	0.430				1.000						3	T	1987	1.000		434 PP; 3020 PP-NE					Y	Y				
19	Gauley Bay Wetland Complex (7420)	Midhurst	17	41H/3;4 1H/4	468000	4983500	C	199.100	0.647	0.232		0.121	0.117	0.883						1	T	1990	0.440	0.560	4658 IBP; 5010 ANSI-LS-R		T	2,6,8	Y		Y	Y			
20	Greenough Harbour (7183)	Midhurst	17	41A/14	466600	4982100	S	27.400	0.450	0.280		0.270		0.720					0.280	2	T	1984	0.700	0.300	4557 IBP				Y	Y		Y	Y		
21	Sadler Creek Wetland Complex (7423)	Midhurst	17	41H/3	462700	4990700	C	261.400	0.601	0.318		0.081	0.020	0.466	0.504				0.010	1	T	1990		1.000	433 ANSI-LS-P; 4655 IBP		T	4,6	Y	Y	Y	Y			
22	Corisande Bay (7179)	Midhurst	17	41H/4	456000	4997700	C	59.600	0.227	0.624		0.149	0.081	0.477		0.008		0.239	0.211	3	T	1985	0.150	0.850	430 ANSI-LS-P; 4635 IBP		T	2,4,6	Y	Y		Y			
23	Dorcas Bay (7180)	Midhurst	17	41H/4	455000	5004500	C	110.100	0.590	0.196		0.214		0.877	0.041		0.055		0.027	1	T	1985	0.130	0.870	172 ANSI-LS-P; 2075 LS; 4541 IBP	NP	T	4,6	Y	Y					

Appendix G: Lake Huron and St. Marys River: Evaluated Coastal Wetlands

									Wetland Type				Site Type																Significant Species						
MAP ID (Figure 2)	WETLAND NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:50 000)	EASTING	NORTHING	SINGLE/COMPLEX	AREA	SWAMP	MARSH	BOG	FEN	ISOLATED	PALUSTRINE	RIVERINE	RIVERINE RIVERMOUTH	LACUSTRINE RIVERMOUTH	LACUSTRINE BAY	LAKE LACUSTRINE	CLASS	PROVINCIALLY SIGNIFICANT WETLAND	YEAROF EVALUATION DATA	PUBLIC OWNERSHIP	PRIVATE OWNERSHIP	CORRESPONDING_NATURAL AREA	OTHER COMMENTS	LOSS	STRESSES	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
24	Barney Lake Wetland Complex (7418)	Midhurst	17	41H/4	447000	5006400	C	151.100	0.527	0.286		0.187		0.993					0.007	3	T	1990	0.010	0.990	171 ANSI-LS-P; 1034 ANSI-ES-P; 4358, 4359 LS; 4646 IBP							Y			
25	Wingfield Basin (7163)	Midhurst	17	41H/3	476500	5009700	C	71.400	0.370	0.630			0.970					0.030		3	T	1987	0.800	0.200	158 ANSI-LS-P, 4200 LS	PWA	F			Y	Y				
26	Collingwood Shores Complex (7178)	Midhurst	17	41A/9	557500	4930000	C	166.500	0.800	0.190		0.010	0.300						0.700	2	T	1985	0.050	0.950			T	6,9		Y	Y				
27	Cranberry Lake Marsh (7446)	Midhurst	17	41A/9	559200	4929000	S	46.200	0.800	0.200				1.000						2	T	1984		1.000		Golf course with Audobon	F	5b,5,6,9		Y	Y				
28	Collingwood Harbour Marsh (7177)	Midhurst	17	41A/9	561000	4928500	S	96.400	0.090	0.890		0.020							1.000	2	T	1986	0.910	0.090			T	2,6,9			Y				
29	Wasaga Beach (7449)	Midhurst	17	41A/9;31D/12	580500	4933000	C	466.500	0.970	0.030			0.260	0.740						PSW	T	1994	0.070	0.930	156 MACRO; 1693 ANSI-LS-P; 3250 PP		T	6,9							
30	Balm Beach Swamp (5639)	Midhurst	17	31D/12	581400	4946900	C	405.500	0.927	0.060	0.009	0.004		0.520	0.480					1	T	1986		1.000	1503 ANSI-LS-R; 5760 IBP	Stewardship Agreement	F	6		Y					
31	Thunder Bay Swamp (7165)	Midhurst	17	41A/16	575000	4960000	C	71.600	0.940	0.060				1.000						3	T	1987		1.000		ANSI	F	6							
32	Awenda Shoreline Fen (7174)	Midhurst	17	31D/13	579600	4966900	S	11.200	0.850			0.150							1.000	3	T	1985	1.000		3012 PP-NE; 4383 ANSI-LS-P		F			Y					
33	Penetang Marsh (7188)	Midhurst	17	31D/13	583000	4957100	C	29.300	0.570	0.420		0.010							1.000	3	T	1985	0.350	0.650	4269 ANSI-ES-P		T	5b,8			Y				
34	Midland Swamp (7187)	Midhurst	17	31D/13	586400	4956000	C	37.600	0.930	0.070				1.000						6 or 5	F	1985	0.900	0.100		Largely a municipal water source	F	6					Y		
35	Sucker Creek (Huronia) (7167)	Midhurst	17	31D/13	588400	4960700	S	32.000	0.440	0.560				0.850	0.150					3	T	1985		1.000			F					Y			
36	Wye Marsh (7164)	Midhurst	17	31D/12	590500	4952300	C	743.900	0.414	0.554	0.004	0.030			1.000					1	T	1987	0.880	0.120	1719 ANSI-LS-R; 2015 NWA; 2179 PWA; 5758 IBP		F	9		Y	Y	Y	Y	Y	
37	Port McNicholl Marsh (7190)	Midhurst	17	31D/13	594300	4956700	S	76.900	0.100	0.900									1.000	2	T	1985	0.900	0.100			F	5b,5c,6,9			Y				
38	Hog Bay Wetland (7184)	Midhurst	17	31D/12	594500	4953500	S	31.600		1.000									1.000	3	T	1984	0.200	0.800			F	6,9			Y				
39	Victoria Harbour Marsh (7166)	Midhurst	17	31D/13	598200	4956600	S	39.000	0.177	0.823									1.000	3	T	1986	0.820	0.180			F	5b,5c,6			Y				
40	Sturgeon Bay Complex (7162)	Midhurst	17	31D/12	599700	4953600	C	192.100	0.381	0.619				0.350					0.650	2	T	1986	0.650	0.350			T	5b,5c,6			Y	Y			
41	Matchedash Bay Marsh (7186)	Midhurst	17	31D/12; 31D/13	605600	4954700	S	807.400	0.070	0.930							0.550		0.450	1	T	1986	0.930	0.070	2198 PWA; 4376 ANSI-LS-P	Ramsar	F	5c,5b,9			Y				
42	Quarry Island Wetland (7504)	Parry Sound	17	31D/13	594000	4965000	C	47.200	0.060	0.940								0.030	0.970	PSW	T	1994	0.920	0.080			F	11			Y				Y

Appendix G: Lake Huron and St. Marys River: Evaluated Coastal Wetlands

									Wetland Type				Site Type																Significant Species						
MAP ID (Figure 2)	WETLAND NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:50 000)	EASTING	NORTHING	SINGLE/COMPLEX	AREA	SWAMP	MARSH	BOG	FEN	ISOLATED	PALUSTRINE	RIVERINE	RIVERINE RIVERMOUTH	LACUSTRINE RIVERMOUTH	LACUSTRINE BAY	LAKE LACUSTRINE	CLASS	PROVINCIALLY SIGNIFICANT WETLAND	YEAROF EVALUATION DATA	PUBLIC OWNERSHIP	PRIVATE OWNERSHIP	CORRESPONDING_NATURAL AREA	OTHER COMMENTS	LOSS	STRESSES	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
43	Tobies Bay Wetland (7505)	Parry Sound	17	31D/13	596000	4966600	S	194.200	0.210	0.790				0.530					0.470	PSW	T	1994	0.650	0.350		Stewardship Agreement	F	11			Y	Y			
44	Partridge Bay Wetland (7506)	Parry Sound	17	41H/8;41H/7	559500	5023700	C	179.700	0.327	0.620	0.053			0.490	0.070				0.440	PSW	T	1993	0.500	0.500		PP	T	4	Y	Y	Y				
45	Naiscoot River Wetland (7534)	Parry Sound	17	41H/10	536000	5055800	S	125.000	0.108	0.892					1.000					PSW	T	1993	1.000				F				Y				
46	White Cove (8440)	Sudbury	17	41H/13	432500	5084600	C	11.280	0.170	0.830				0.170					0.830	NPSW	F	1994		1.000											
47	Spanish River Delta Marsh (3663)	Sudbury	17	41J/1	397000	5115000	C	304.500	0.210	0.790				0.140	0.030		0.830			PSW	T	1993	0.820	0.180	4845 ANSI-LS-C; 2024 INDIAN RESERVE		F	2,5a,5b,5			Y				
48	Marsh Bay Island 9 Wetland (7629)	Sault Ste. Marie	17	41J/2	346000	5116000	C	254.0	0.320	0.450		0.230	0.010	0.790	0.150		0.020		0.030	3	T	1994	0.800	0.200			T	2,5a,6,8	Y	Y					
				Total Wetland Area				**7458.97**																											
49	Hay Bay Wetland (7864)	Sault Ste. Marie	17	41J/5	289000	5131000	C	161.0	0.920	0.080				0.930			0.070			NPSW	F	1997		1.000							Y				
50	Hay Marsh (7630)	Sault Ste. Marie	17	41J/4	272000	5105000	S	2275.0	0.500	0.300		0.200		0.500					0.500	3	T	1993	0.100	0.900		NP, MBS	T	2,6,8							
51	Anderson Creek (7633)	Sault Ste. Marie	17	41J/5	272000	5134600	C	42.0	0.350	0.650				0.850			0.150			1	F	1993	0.100	0.900			F	2,8							
52	Maskinonge Bay (7631)	Sault Ste. Marie	16	41K/8	724000	5136800	C	63.0	0.140	0.860									1.000	1	F	1993	1.000	0.000			T	2,8,6							
53	Neebish Concession Wetland (7632)	Sault Ste. Marie	16	41K/8	723400	5125000	S	54.0	0.220	0.780									1.000	1	F	1993	0.770	0.230			T	2,8,6							
54	Lake George (3641)	Sault Ste. Marie	16	41K/8	722500	5150000	S	262.0	0.200	0.800									1.000	3	T	1993	0.200	0.800			T	2,6,7,8							
55	Echo Bay (3640)	Sault Ste. Marie	16	41K/9	726000	5153000	S	710.0	0.020	0.980							0.400	0.600		3	T	1993	0.440	0.560			T	2,8,7							
				Total Wetland Area				**3567.00**																											

Appendix H: Lake Huron and St. Marys River: Natural Areas, Environmental Sensitivity Atlas Wetlands, and Other Wetlands with an Unevaluated Coastal Wetland Component

									Significant Species						
MAP ID	AREA NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:25 000)	EASTING	NORTHING	DESCRIPTION	CORRESPONDING NATURAL AREAS	COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
Natural Area Wetlands, Lake Huron															
N1	Ipperwash Provincial Park ANSI-LS-P (4)	Aylmer	17	40P/4	422500	4785200	-graminoid coastal meadow marsh community	N/A	Y						
N2	Ipperwash Military Reserve ANSI-LS-R (5)	Aylmer	17	40P/4	424000	4784500	-graminoid coastal meadow marsh community	N/A	Y						
N3	Pinery Provincial Park ANSI-LS-P (7)	Aylmer	17	40P/5	432000	4790000	-graminoid coastal meadow marsh community	N/A	Y						
N4	Inverhuron PP (3084)	Midhurst	17	41A/5	453000	4905250		N/A							
N5	Chiefs Point Indian Reserve ANSI-LS-R (435)	Midhurst	17	41A/11	477000	4948000	-shoreline & fen vegetation typical of the Bruce Peninsula; most dominant vegetation community is cedar-white birch-balsam fir swamp	N/A							
N6	Old Woman River IBP (4669)	Midhurst	17	41A/14	473800	4980200	-shrubby cinquefoil coastal meadow marsh community	N/A	Y						
N7	Crane Island or Dane Island IBP (4632)	Midhurst	17	41A/4	469400	4977100	-graminoid coastal meadow marsh community	N/A	Y						
N8	Lyal Island ANSI-LS-P (629)	Midhurst	17	41A/14	468000	4977500	-wet active sandy beach with open to closed marsh & wet meadow communities	N/A		Y		Y			
N9	Ira Lake ANSI-LS-P (7058)	Midhurst	17	41H/3	470000	4986000	-graminoid coastal meadow marsh community	N/A	Y						
N10	Bradley Harbour IBP(4676)	Midhurst	17	41H/3	464000	4986100	-graminoid coastal meadow marsh community	N/A	Y						
N11	Johnston Harbour Pine Tree Point ANSI-LS-P (431)/ PP-NR (3090)	Midhurst	17	41H/4	459000	4994000	-well developed large shallow marsh at mouth of Crane River; marshes at Johnston Harbour	N/A		Y	Y				
N12	Hay Bay ANSI-LS-R (5008)	Midhurst	17	41H/4	445000	5009000	-shallow rocky bay which has several uncommon aquatic plant species and excellent wildlife habitat	N/A							
N13	Driftwood Cove-Little Cove IBP (4662)	Midhurst	17	41H/4	453500	5010300	-shrubby cinquefoil coastal meadow marsh community	N/A	Y						

Appendix H: Lake Huron and St. Marys River: Natural Areas, Environmental Sensitivity Atlas Wetlands, and Other Wetlands with an Unevaluated Coastal Wetland Component

									Significant Species						
MAP ID	AREA NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:25 000)	EASTING	NORTHING	DESCRIPTION	CORRESPONDING NATURAL AREAS	COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
N14	Cameron Lake Fen ANSI-LS-R (4202)	Midhurst	17	41H/4	457500	5007300	-graminoid coastal meadow marsh & shrubby cinquefoil coastal meadow marsh community	N/A	Y						
N15	Bartley Lake IBP (4672)	Midhurst	17	41H/3	461900	5006200	-graminoid coastal meadow marsh community	N/A	Y						
N16	Cabot Head (Lindsay Twp Portion) LS (4200)	Midhurst	17	41H/3	476000	5009000	-graminoid coastal meadow marsh & shrubby cinquefoil coastal meadow marsh community	N/A	Y						
N17	Lion's Head Peninsula ANSI-LS-P (4197)	Midhurst	17	41A/14	483577	4981520	-moist meadows; open ponds ringed by emergent marshes at McKay's Harbour	N/A				Y			
N18	Hope Bay Forest PNR (3082)	Midhurst	17	41A/14	486000	4973500	-at Bulls Landing has open alvars, shoreline graminoid and shrub fens, mixed and conifer seepage swamps	N/A		Y	Y				
N19	Sucker Creek - Cape Rich ANSI-LS-P (1647)	Midhurst	17	41A/10	523000	4950000	-shoreline fen occurs east of the mouth of Sucker Creek estuary; shallow water marsh on west side of Vails Point	N/A		Y	Y				
N20	Wasaga Beach PP (3250)	Midhurst	17	41A/9	578175	4927663	-graminoid coastal meadow marsh community	N/A	Y						
N21	Beausoleil and Associated Islands ANSI-LS-P (6232)	Parry Sound	17	31D/13	590000	4971000	-graminoid coastal meadow marsh; marshes occur along the coast of Beausoleil	N/A	Y			Y			
N22	McDonald Bay - Mcrae Lake ANSI-LS-C (4923)	Parry Sound	17	31D/13	595000	4975000	-dense aquatic vegetation, graminoid bog mats, bogs and marshes; also thicket and coniferous swamp	N/A		Y		Y			
N23	O'Donnell Point PNR (3172)	Parry Sound	17	41H/1	576500	4992000	-upland/lowland forests and extensive wetlands and bogs; excellent representation of shallow peatland habitats	N/A		Y		Y			
N24	Sandy Island ANSI-LS-C (4929)	Parry Sound	17	41H/8	558000	5014000	-extensive emergent shoreline complexes	N/A							
N25	Killbear Provincial Park: Natural Environment & Nature Reserve Zone ANSI-LS-C (4919)	Parry Sound	17	41H/8	563000	5022500	-deciduous & thicket swamps along the lakeshore; peat dominated bog pools inland; emergent vegetation & peatland	N/A							
N26	Franklin Island IBP (4475)	Parry Sound	17	41H/8&7	551500	5027000		N/A							

Appendix H: Lake Huron and St. Marys River: Natural Areas, Environmental Sensitivity Atlas Wetlands, and Other Wetlands with an Unevaluated Coastal Wetland Component

									Significant Species						
MAP ID	AREA NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:25 000)	EASTING	NORTHING	DESCRIPTION	CORRESPONDING NATURAL AREAS	COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
N27	Shebeshekong ANSI-LS-P (4930)	Parry Sound	17	41H/8	560000	5033000		N/A							
N28	Shawanaga Park Reserve LS (5728)	Parry Sound	17	41H/9	549000	5046000		N/A		Y					
N29	Southwest Sturgeon Bay PP-NR (3520)	Parry Sound	17	41H/9	544560	5052200	-shallow soil cover and wetlands typical of Precambrian Shield; part of a provincially significant wetland	N/A							
N30	Point Au Baril ANSI-LS-C (4927)	Parry Sound	17	41H/9	542000	5048000	-shoreline marsh vegetation; thicket swamp & bog	N/A							
N31	French River Waterway Park Delta Candidate Nature Reserve Zone ANSI-LS-C (6236)	Sudbury	17	41H/15	515000	5090000	-"ridge and swale" wetland complex with emergent aquatic vegetation and shallow water aquatic habitats	N/A		Y		Y			
N32	Boom Point-Sand Bay ANSI-LS-C (4855)	Sudbury	17	41G/14	318000	5080000		N/A							
N33	Srigley Bay-Marsh Lake ANSI-LS-C (4893)	Sudbury	17	41G/9	384700	5064000	-shoreline consists of forb and graminoid meadows; inland ridges disappear under a mantle of swamp vegetation	N/A							
N34	Deans Bay-Sand Lakes ANSI-LS-C (4861)	Sudbury	17	41G/9	397000	5060000	-cedar is prevalent species; north end of Sand Lakes are bogs; alvar also present	N/A							
N35	Fitzwilliam Island ANSI-LS-C (4865)	Sudbury	17	41H/5	438000	5035000	-wetlands associated with ponds and lakes on island	N/A							
N36	Mississagi Bay Shoreline Marsh ANSI-LS-C (3634)	Sault Ste. Marie	17	41J/3	338000	5118500	-extensive area of deep and shallow emergent shoreline marshes; also meadow marshes	N/A							
N37	Foul Bight Beach Ridge and Swale ANSI-LS-R (3632)	Sault Ste. Marie	17	41J/3	334000	5119000	-raised beach ridge swales system; cedar & black spruce swamp; wetland in swales	N/A							
N38	Dayton Beach Ridge and Swale ANSI-LS-R (3635)	Sault Ste. Marie	17	41J/3	317000	5122500	-raised beach ridge swale system; graminoid meadows; marsh, fen and thicket swamp characterize the swales	N/A							
Environmental Sensitivity Atlas Wetlands, Lake Huron															
A1	Methodist Island	Midhurst	17	31D/12	596300	4958400	Huron ESA - Map 86	N/A							
A2	Bay southwest of Kirk Creek mouth	Sudbury	17	41 I/4	458500	5100100	Huron ESA - Map 59	N/A							
A3	Kirk Creek Mouth	Sudbury	17	41I/4	458900	5101000	Huron ESA - Map 59	N/A							
A4	East Sampson Island	Sudbury	17	41I/4	455300	5101700	Huron ESA - Map 59	N/A							

Appendix H: Lake Huron and St. Marys River: Natural Areas, Environmental Sensitivity Atlas Wetlands, and Other Wetlands with an Unevaluated Coastal Wetland Component

									Significant Species						
MAP ID	AREA NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:25 000)	EASTING	NORTHING	DESCRIPTION	CORRESPONDING NATURAL AREAS	COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
A5	MacGregor Island - east channel	Sudbury	17	41I/4	455800	5102400	Huron ESA - Map 59	N/A							
A6	Middle Sampson Island	Sudbury	17	41I/4	452100	5100500	Huron ESA - Map 59	N/A							
A7	Iroquois Island	Sudbury	17	41I/4	450300	5104000	Huron ESA - Map 59	N/A							
A8	Northshore Iroquois Bay 1	Sudbury	17	41I/4	452400	5107000	Huron ESA - Map 59	N/A							
A9	Northshore Iroquois Bay 2	Sudbury	17	41I/4	450700	5107300	Huron ESA - Map 59	N/A							
A10	Jumbo Island	Sudbury	17	41I/4	450000	5100300	Huron ESA - Map 59	N/A							
A11	South shore of Little La Cloche Island	Sudbury	17	41H/13	441500	5091100	Huron ESA - Map 58	N/A							
A12	Shoal Bight, Great La Cloche Island	Sudbury	17	41H/13	438600	5092800	Huron ESA - Map 56	N/A							
A13	Bay between English Point and Stony Point (Great La Cloche Island)	Sudbury	17	41H/13	437300	5092700	Huron ESA - Map 56	N/A							
A14	Bowell Cove, Strawberry Island	Sudbury	17	41H/13	434800	5087300	Huron ESA - Map 56	N/A							
A15	Chapleau Cove, Strawberry Island	Sudbury	17	41H/13	434200	5087800	Huron ESA - Map 56	N/A							
A16	Strawberry Channel (east shore)	Sudbury	17	41H/13	432500	5087600	Huron ESA - Map 56; 4895 ANSI-LS-C; -extensive Great Lakes open water shoreline marshes; monotypic marsh backed by Red Maple swamp	N/A							
A17	Strawberry Channel (west shore)	Sudbury	17	41H/13	430900	5087100	Huron ESA - Map 56	N/A							
A18	Bays between Thompson Point, Long Point and Gibbons Point, Manitoulin Island	Sudbury	17	41H/13	430700	5089200	Huron ESA - Map 56	N/A							
A19	Bay south of Francis Point, Manitoulin Island	Sudbury	17	41G/16	405600	5087200	Huron ESA - Map 23	N/A							
A20	Rushy Cove, Manitoulin Island	Sudbury	17	41G/16	405500	5088000	Huron ESA - Map 23	N/A							
A21	Gore Bay, Manitoulin Island	Sudbury	17	41G/16	386700	5085500	Huron ESA - Map 24	N/A							
A22	Rozel's Bay, Manitoulin Island	Sudbury	17	41G/15	374500	5084600	Huron ESA - Map 25; 4337 Rozel's Bay Alvar	N/A							
A23	Bay east of Jacks Island, Manitoulin Island	Sudbury	17	41G/15	379600	5080000	Huron ESA - Map 26; 4336 Foxey Barren Alvar	N/A							
A24	Entrance to Lake Wolsley opposite Indian Point, Manitoulin Island	Sudbury	17	41G/15	379500	5077300	Huron ESA - Map 26	N/A							
A25	North bay of Lake Wolsley, Manitoulin Island	Sudbury	17	41G/15	382000	5079200	Huron ESA - Map 26; 4879 Candidate LS ANSI; -mixed forest swamp developed at head of bay; also shoreline marshes	N/A							

Appendix H: Lake Huron and St. Marys River: Natural Areas, Environmental Sensitivity Atlas Wetlands, and Other Wetlands with an Unevaluated Coastal Wetland Component

									Significant Species						
MAP ID	AREA NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:25 000)	EASTING	NORTHING	DESCRIPTION	CORRESPONDING NATURAL AREAS	COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
A26	East shore of Lake Wolsley, Manitoulin Island	Sudbury	17	41G/15	382700	5074300	Huron ESA - Map 26	N/A							
A27	South bay of Lake Wolsley, Manitoulin Island	Sudbury	17	41G/15	380000	5070800	Huron ESA - Map 26	N/A							
A28	Cooks Bay, Manitoulin Island	Sudbury	17	41G/15	359800	5082400	Huron ESA - Map 28	N/A							
A29	Bay northwest of Stony Island (Sheshegwaning First Nation), Manitoulin Island	Sudbury	17	41G/15	357600	5086800	Huron ESA - Map 29	N/A							
A30	Shigniconing Bay (Sheshegwaning First Nation), Manitoulin Island	Sudbury	17	41G/15	355500	5092700	Huron ESA - Map 29	N/A							
A31	Pitman Point, Cockburn Island	Sudbury	17	41G/14	318600	5095300	Huron ESA - Map 32; -sensitive area due to the presence of fens, marshes and swamps	N/A							
A32	Mundy's Bay, Cockburn Island	Sudbury	17	41G/14	315400	5095200	Huron ESA - Map 32	N/A							
A33	Wagosh Bay, Cockburn Island	Sudbury	17	41G/14	308800	5085600	Huron ESA - Map 33; 4902 ANSI-LS-C; -sensitive area due to presence of undisturbed wetlands, fens, bogs and marshes	N/A							
A34	Hyndman Bay, Cockburn Island	Sudbury	17	41G/14	311400	5084000	Huron ESA - Map 33	N/A							
A35	Mud Bay, Cockburn Island	Sudbury	17	41G/14	317400	5080400	Huron ESA - Map 34	N/A							
A36	Bay between Edna Point and Burnt Island, Manitoulin Island	Sudbury	17	41G/15	349600	5076000	Huron ESA - Map 36; 4339 Burnt Island Harbour	N/A							
A37	Shoreline north of Black Point, Manitoulin Island	Sudbury	17	41G/15	351200	5074300	Huron ESA - Map 36	N/A							
A38	Walkhouse Bay, Manitoulin Island	Sudbury	17	41G/15	354200	5072900	Huron ESA - Map 36	N/A							
A39	Carroll Wood Bay, Manitoulin Island	Sudbury	17	41G/15	357500	5075000	Huron ESA - Map 38	N/A							
A40	Little Huron Lake (Sand Bay), Manitoulin Island	Sudbury	17	41G/15	362000	5073600	Huron ESA - Map 38	N/A							
A41	Misery Point, Manitoulin Island	Sudbury	17	41G/15	363900	5071300	Huron ESA - Map 38	N/A							
A42	Misery Bay, Manitoulin Island	Sudbury	17	41G/15	364400	5073200	Huron ESA - Map 38; 5530 PP; 3145 PNR; -coastal meadow marsh; alvar pavement; fen complex; mixed & coniferous swamp	N/A							

Appendix H: Lake Huron and St. Marys River: Natural Areas, Environmental Sensitivity Atlas Wetlands, and Other Wetlands with an Unevaluated Coastal Wetland Component

									Significant Species						
MAP ID	AREA NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:25 000)	EASTING	NORTHING	DESCRIPTION	CORRESPONDING NATURAL AREAS	COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
A43	Mac's Bay, Manitoulin Island	Sudbury	17	41G/15	368000	5072100	Huron ESA - Map 39; 4340 Mac's Bay Alvar	N/A							
A44	Murphy's Harbour, Manitoulin Island	Sudbury	17	41G/15	369200	5071300	Huron ESA - Map 39	N/A							
A45	Dominion Point	Sudbury	17	41G/9	388300	5061300	Huron ESA - Map 40; 4335 Dominion Point Alvar; possible coastal meadow marsh	N/A							
A46	Shoreline west of Blackrock, Manitoulin Island	Sudbury	17	41G/9	399300	5057400	Huron ESA - Map 42	N/A							
A47	Hughson Bay, Manitoulin Island	Sudbury	17	41G/9	406700	5053300	Huron ESA - Map 42	N/A							
A48	Rathburn Bay, Manitoulin Island	Sudbury	17	41G/9	408400	5051700	Huron ESA - Map 42	N/A							
A49	Michaels Bay - Blue Jay Creek, Manitoulin Island	Sudbury	17	41G/9	414800	5049300	Huron ESA - Map 42; 4875 ANSI-LS-C; -raised beach ridge swale, marsh with fen, coniferous & thicket swamp; includes Blue Jay Creek wetland	N/A							
A50	Big Bay, Manitoulin Island	Sudbury	17	41G/9	415800	5047600	Huron ESA - Map 43	N/A							
A51	South of Leason Bay - North of Frood Harbour, Manitoulin Island	Sudbury	17	41H/12	423700	5049800	Huron ESA - Map 43	N/A							
A52	Leason Bay, Manitoulin Island	Sudbury	17	41H/12	424800	5050800	Huron ESA - Map 43	N/A							
A53	Bay south of Sims Bay, Manitoulin Island	Sudbury	17	41H/12	429400	5053900	Huron ESA - Map 45	N/A							
A54	Sims Bay, Manitoulin Island	Sudbury	17	41H/12	431200	5056600	Huron ESA - Map 45	N/A							
A55	Bay Southwest of Leask Point, Manitoulin Island	Sudbury	17	41H/12	433000	5057600	Huron ESA - Map 45	N/A							
A56	Leask Bay (3 wetlands), Manitoulin Island	Sudbury	17	41H/12	433800	5059200	Huron ESA - Map 45	N/A							
A57	Mouth of Riv la Cloche	Sudbury	17	41J/1	417200	5107000	Huron ESA - Map 19	N/A							
A58	Fort La Cloche	Sudbury	17	41J/1	415400	5106900	Huron ESA - Map 19	N/A							
A59	South shore of Spanish River First Nation	Sudbury	17	41J/1	412700	5107300	Huron ESA - Map 19	N/A							
A60	Long Bay, Spanish River First Nation	Sudbury	17	41J/1	403400	5111400	Huron ESA - Map 17	N/A							
A61	Bay Northwest of Graham Point, Spanish River First Nation	Sudbury	17	41J/1	398800	5111300	Huron ESA - Map 17	N/A							
Environmental Sensitivity Atlas Wetlands, St. Marys River															
A62	Joe Dollar Bay/Palideau Island	Sault St. Marie	17	41J/5	290500	5128700	Huron ESA - Map 6	N/A							

Appendix H: Lake Huron and St. Marys River: Natural Areas, Environmental Sensitivity Atlas Wetlands, and Other Wetlands with an Unevaluated Coastal Wetland Component

									Significant Species						
MAP ID	AREA NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:25 000)	EASTING	NORTHING	DESCRIPTION	CORRESPONDING NATURAL AREAS	COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
A63	Garden Bay	Sault St. Marie	17	41J/5	287300	5130800	Huron ESA - Map 5	N/A							
A64	Shoreline north of Green Island	Sault St. Marie	17	41J/5	282000	5131200	Huron ESA - Map 5	N/A							
A65	Stobie Creek	Sault St. Marie	17	41J/5	278000	5135300	Huron ESA - Map 5	N/A							
A66	Dawson Island	Sault St. Marie	17	41J/5	277200	5134100	Huron ESA - Map 5	N/A							
A67	Bay north of Lasher Island	Sault St. Marie	17	41J/5	275100	5135500	SMR ESA - Map 9	N/A							
A68	Desbarats Wetland	Sault St. Marie	17	41J/5	274500	5135600	SMR ESA - Map 9; 3644 Desbarats Marsh; -large complex of marsh and swamp; possible coastal meadow marsh	N/A							
A69	East of Kensington Point	Sault St. Marie	17	41J/5	273600	5134600	SMR ESA - Map 9	N/A							
A70	West of Kensington Point	Sault St. Marie	17	41J/5	272900	5134700	SMR ESA - Map 9	N/A							
A71	Quebec Bay	Sault St. Marie	17	41J/5	269300	5134300	SMR ESA - Map 9	N/A							
A72	Bay east of Bowen Island	Sault St. Marie	16	41K/8	726800	5133600	SMR ESA - Map 9	N/A							
A73	East shore of Bowen Island	Sault St. Marie	16	41K/8	726100	5133700	SMR ESA - Map 9	N/A							
A74	Bay opposite Sutton Island	Sault St. Marie	16	41K/8	725800	5135300	SMR ESA - Map 9	N/A							
A75	East shore of Pine Island	Sault St. Marie	16	41K/8	725400	5134800	SMR ESA - Map 9	N/A							
A76	Bay Southeast of Montague Island	Sault St. Marie	17	41J/5	270400	5132800	SMR ESA - Map 9	N/A							
A77	Bay North of Sucker Creek Mouth	Sault St. Marie	17	41J/5	272600	5130700	SMR ESA - Map 9	N/A							
A78	Sucker Creek Wetland, St. Joseph Island	Sault St. Marie	17	41J/5	272400	5130500	Huron ESA - Map 4	N/A							
A79	Campement d'Ours Island-East Shore	Sault St. Marie	17	41J/5	274100	5131700	SMR ESA - Map 9	N/A							
A80	Fisher Bay, St. Joseph Island	Sault St. Marie	17	41J/5	279200	5126400	SMR ESA - Map 10	N/A							
A81	Old Quarry Point, McMenomy Bay - north shore, St. Joseph Island	Sault St. Marie	17	41J/4	281200	5125900	SMR ESA - Map 10	N/A							
A82	West of Recollect Bay, McMenomy Bay - south shore, St. Joseph Island	Sault St. Marie	17	41J/4	281400	5023850	SMR ESA - Map 10	N/A							
A83	Moffat Bay, St. Joseph Island	Sault St. Marie	17	41J/4	283200	5121900	Huron ESA - Map 3	N/A							
A84	Mosquito Bay, St. Joseph Island	Sault St. Marie	17	41J/4	283500	5119900	Huron ESA - Map 3	N/A							
A85	Garside Bay, St. Joseph Island	Sault St. Marie	17	41J/4	284200	5117700	Huron ESA - Map 3	N/A							
A86	Richmond Bay/Richmond Point, St. Joseph Island	Sault St. Marie	17	41J/4	282700	5116600	Huron ESA - Map 2	N/A							
A87	Milford Haven Creek Mouth, St. Joseph Island	Sault St. Marie	17	41J/4	281600	5114900	Huron ESA - Map 2	N/A							

Appendix H: Lake Huron and St. Marys River: Natural Areas, Environmental Sensitivity Atlas Wetlands, and Other Wetlands with an Unevaluated Coastal Wetland Component

									Significant Species						
MAP ID	AREA NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:25 000)	EASTING	NORTHING	DESCRIPTION	CORRESPONDING NATURAL AREAS	COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
A88	Wallace Island (3 small wetlands), St. Joseph Island	Sault St. Marie	17	41J/4	284300	5114300	Huron ESA - Map 2	N/A							
A89	Tenby Bay (3 wetlands), St. Joseph Island	Sault St. Marie	17	41J/4	277500	5114700	Huron ESA - Map 1	N/A							
A90	Whiskey Bay, St. Joseph Island	Sault St. Marie	17	41J/4	276600	5108600	Huron ESA - Map 1	N/A							
A91	Elliot Point, St. Joseph Island	Sault St. Marie	17	41J/4	275300	5106700	Huron ESA - Map 1	N/A							
A92	Old Fort St. Joe - La Pointe shoreline wetlands, St. Joseph Island	Sault St. Marie	17	41J/4	272800	5105500	Huron ESA - Map 1	N/A							
A93	Shoreline north of Hay Marsh, St. Joseph Island	Sault St. Marie	17	41J/4	268800	5115000	SMR ESA - Map 13	N/A							
A94	Shoreline south of Richardson Creek Mouth (near Otter Lake), St. Joseph Island	Sault St. Marie	16	41K/1	730000	5116700	SMR ESA - Map 13	N/A							
A95	Richardson Creek Mouth, St. Joseph Island	Sault St. Marie	16	41K/1	727200	5118300	SMR ESA - Map 13	N/A							
A96	Shoreline between Court Pt. - Sunset Pt., St. Joseph Island	Sault St. Marie	16	41K/1	725700	5120400	SMR ESA - Map 13	N/A							
A97	Shoreline south of Everens Point, St. Joseph Island	Sault St. Marie	16	41K/8	724500	5123300	SMR ESA - Map 11	N/A							
A98	Sailors Encampment, St. Joseph Island	Sault St. Marie	16	41K/8	723900	5127100	SMR ESA - Map 11	N/A							
A99	Shoreline between Coyle and Reed Point, St. Joseph Island	Sault St. Marie	16	41K/8	723100	5128200	SMR ESA - Map 11	N/A							
A100	Shoreline north of Coyle Point, St. Joseph Island	Sault St. Marie	16	41K/8	722800	5129700	SMR ESA - Map 11	N/A							
A101	Stribling Point, St. Joseph Island	Sault St. Marie	16	41K/8	722400	5132900	SMR ESA - Map 11	N/A							
A102	Southwest shore of East Neebish Island	Sault St. Marie	16	41K/8	722800	5135900	SMR ESA - Map 8	N/A							
A103	East Shore of East Neebish Island	Sault St. Marie	16	41K/8	722200	5136800	SMR ESA - Map 8	N/A							
A104	Niles Point (Watson Island)	Sault St. Marie	16	41K/8	723300	5135200	SMR ESA - Map 8	N/A							
A105	Pumpkin Point	Sault St. Marie	16	41K/8	720800	5142200	SMR ESA - Map 8	N/A							
A106	Squirrel Island	Sault St. Marie	16	41K/9	721300	5155600	SMR ESA - Map 5	N/A							
A107	Little Lake George/Point Charles	Sault St. Marie	16	41K/9	716800	5157500	SMR ESA - Map 5	N/A							
A108	Bells Point, Little Lake George	Sault St. Marie	16	41K/9	714000	5158000	SMR ESA - Map 5	N/A							

Appendix H: Lake Huron and St. Marys River: Natural Areas, Environmental Sensitivity Atlas Wetlands, and Other Wetlands with an Unevaluated Coastal Wetland Component

									Significant Species						
MAP ID	AREA NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:25 000)	EASTING	NORTHING	DESCRIPTION	CORRESPONDING NATURAL AREAS	COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
A109	Whitefish Island	Sault St. Marie	16	41K/9	703000	5154300	SMR ESA - Map 3; 3660 Sault Ste. Marie NP; -wetland areas	N/A							
A110	Carpin Beach (Carp River, Leigh Bay)	Sault St. Marie	16	41K/9	696500	5153400	SMR ESA - Map 2	N/A							
A111	Marks Bay	Sault St. Marie	16	41K/8	693200	5151500	SMR ESA - Map 2; 4831 ANSI-LS-C; -alder, thicket & coniferous swamp; meadow community consisting of graminoid and forb species	N/A							
A112	Point Louise	Sault St. Marie	16	41K/8	693500	5149300	SMR ESA - Map 2	N/A							
Other Reported Wetlands, St. Marys River															
O1	Birch Point		16	41K/8	721200	5139500	-wetland undergoing evaluation; approximately 45 ha; swamp, marsh and fen; primarily palustrine	N/A							

Appendix I: Detroit River, Lake St. Clair, and St. Clair River: Evaluated Coastal Wetlands

									Wetland Type				Site Type																Significant Species						
MAP ID (Figure 3)	WETLAND NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:50 000)	EASTING	NORTHING	SINGLE/COMPLEX	AREA	SWAMP	MARSH	BOG	FEN	ISOLATED	PALUSTRINE	RIVERINE	RIVERINE RIVERMOUTH	LACUSTRINE RIVERMOUTH	LACUSTRINE BAY	LAKE LACUSTRINE	CLASS	PROVINCIALLY SIGNIFICANT WETLAND	YEAROF EVALUATION DATA	PUBLIC OWNERSHIP	PRIVATE OWNERSHIP	CORRESPONDING_NATURAL AREA	OTHER COMMENTS	LOSS	STRESSES	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
St. Clair River																																			
10	Chenal Ecarte Marshes (7113)	Aylmer	17	40J/9	383500	4712500	C	26.60	0.100	0.900					1.000					2	T	1987	0.300	0.700				2,6,7,9,			Y				
11	Snye River Marshes (7110)	Aylmer	17	40J/9	378900	4717000	C	44.30	0.095	0.905					1.000					2	T	1987	0.100	0.900						Y	Y				
12	Marshy Creek Marsh (7109)	Aylmer	17	40J/9	377800	4721100	S	3.10		1.000					1.000					3	T	1987		1.000	5661 LS	Owned by SCPC		6			Y				
13	Stag Island Marsh (7111)	Aylmer	17	40J/16	380300	4749000	S	10.20		1.000					1.000					3	T	1986		1.000							Y				
14	Point Edward Marsh (7112)	Aylmer	17	40J/16	384300	4760800	S	3.00		1.000					1.000					5	F	1987		1.000			T	6,8,11		Y	Y	Y			
				Total Wetland Area				**87.20**																											
Lake St. Clair																																			
5	Ruscom Shores Marsh (7117)	Aylmer	17	40J/7	367000	4684400	S	29.00	0.140	0.860								1.000		3	T	1985	1.000		2060 CAA; 5213 LS			5b,7,9		Y	Y	Y			Y
6	Tremblay Beach Marsh (7118)	Aylmer	17	40J/7	375400	4684800	S	24.00	0.080	0.920								1.000		3	T	1985	1.000		2066 CAA			5b,7,9			Y				
7	Paternoster Club Marsh (7116)	Aylmer	17	40J/8	378500	4684800	S	3.90		1.000								1.000		3	T	1989		1.000							Y				
8	Thames River Mouth (5704)	Aylmer	17	40J/8	382300	4686300	C	131.00	0.240	0.760							1.000			2	T	1985	0.070	0.930				4,5,6,7,8,9		Y	Y	Y		Y	Y
9	St. Clair Marshes (5088)	Aylmer	17	40J/8	383500	4699000	C	2335.00	0.100	0.900				0.100			0.010	0.530	0.360	1	T	1989	0.130	0.870	38 ANSI-LS-P; 237 MACRO; 5065 NWA; 5077 ESA; 7017 LS	Ramsar		6,7,8,9	Y	Y	Y	Y		Y	
				Total Wetland Area				**2522.90**																											
Detroit River																																			
1	Canard River Marshes (5644)	Aylmer	17	40J/3	326000	4671000	C	416.33		1.000						1.000				1	T	1984	0.660	0.340	34 ANSI-LS-R; 5199			2,5b,5c,8,9	Y	Y	Y	Y			
2	Detroit River Marshes (5683)	Aylmer	17	40J/3	326500	4676500	C	575.00	0.040	0.960					1.000					2	T	1985	0.700	0.300			T	2,5,5b,5c,6,7,8		Y	Y	Y			
3	Fighting Island (7115)	Aylmer	17	40J/3	325100	4678100	S	113.00	0.060	0.940					1.000					3	T	1985		1.000						Y	Y	Y			
4	Turkey Creek Marsh (7114)	Aylmer	17	40J/3	327300	4678900	S	32.00	0.230	0.770					0.800	0.200				3	T	1985	0.800	0.200				2,5,8		Y	Y	Y			Y
				Total Wetland Area				**1136.33**																											

Appendix J: Detroit River, Lake St. Clair and St. Clair River: Natural Areas, Environmental Sensitivity Atlas Wetlands, and Other Wetlands with an Unevaluated Coastal Wetland Component

									Significant Species						
MAP ID	**AREA NAME (NATURAL AREA NUMBER)**	**MNR DISTRICT**	**UTM ZONE**	**MAP SHEET (1:25 000)**	**EASTING**	**NORTHING**	**DESCRIPTION**	**OTHER CORRESPONDING NATURAL AREAS**	**VEGETATION COMMUNITY**	**PLANT**	**BIRD**	**REPTILE/AMPHIBIAN**	**MAMMAL**	**FISH**	**LEPIDOPTERAN**
Environmental Sensitivity Atlas Wetlands															
A1	Walpole Island Marshes, Fens and Prairies ANSI-LS-R (13)	Aylmer	17	40J/10	372000	4711000	Includes Walpole, Squirrel & St. Anne islands. Primarily cattail marsh. Wetland vegetation types include wet meadows, fen, cattail marsh & reed marsh. Approximately 10 360 ha of wetland.	N/A		Y	Y	Y		Y	

Appendix K: Lake Erie: Evaluated Coastal Wetlands

									Wetland Type				Site Type																Significant Species						
MAP ID (Figure 4)	WETLAND NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:50 000)	EASTING	NORTHING	SINGLE/COMPLEX	AREA	SWAMP	MARSH	BOG	FEN	ISOLATED	PALUSTRINE	RIVERINE	RIVERINE RIVERMOUTH	LACUSTRINE RIVERMOUTH	LACUSTRINE BAY	LAKE LACUSTRINE	CLASS	PROVINCIALLY SIGNIFICANT WETLAND	YEAROF EVALUATION DATA	PUBLIC OWNERSHIP	PRIVATE OWNERSHIP	CORRESPONDING_NATURAL AREA	OTHER COMMENTS	LOSS	STRESSES	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
1	Six Mile Creek (Fort Erie 7) (7142)	Cambridge	17	30L/14	661800	4749000	S	11.88	0.240	0.760					0.980		0.020			2	T	1987	0.150	0.850	5360 ANSI-LS-R; 5414 LS		F			Y	Y				
2	Point Abino (7160)	Cambridge	17	30L/14	655300	4746600	C	153.00	0.970	0.030				1.000						2	T	1984	0.086	0.914	15 ANSI-LS-P; 4487 IBP					Y	Y				
3	Empire Beach Backshore Basin Forest (7138)	Cambridge	17	30L/14	651500	4748500	S	61.26	0.950	0.050				1.000						3	T	1988	0.350	0.650	109 LS; 1224 ANSI-LS-P		F	11			Y				
4	Nickle Beach Woodlot & Marsh (7154)	Cambridge	17	30L/14	644200	4748600	C	58.30	0.343	0.657					0.828			0.172		3	T	1985		1.000	5458 LS		F	5a,11							
5	Erco Wetland (7140)	Cambridge	17	30L/13	616800	4746000	S	22.30	0.850	0.150						1.000				3	T	1983		1.000			T	5a,6	Y						
6	Dunnville Marshes (7137)	Cambridge	17	30L/13	616600	4748700	C	518.30	0.040	0.960						1.000				1	T	1983		1.000	1198 ANSI-LS-P; 5182 LS	EHJV purchase	T	4,5a,5b,5c, 6,8,9			Y				
7	James N. Allen Park Woodlot (7146)	Cambridge	17	30L/13	610300	4744400	C	58.52	0.650	0.350				0.900	0.080	0.020				2	T	1987	0.250	0.750	3088 PP; 5187 LS		F	11	Y	Y	Y				
8	Evans Creek Mouth (LET3) (7141)	Aylmer	17	30L/13	601300	4744200	S	3.20	0.220	0.780					1.000					NPSW	F	1996DU		1.000			F	5a,5b,5c,9							
9	Wardelle Creek Mouth (7158)	Aylmer	17	30L/13	600000	4744400	S	12.00	0.070	0.930					1.000					6	F	1986		1.000	5178 LS		F	5a,5b,5c,9		Y					
10	Gates Creek Mouth (LET1) (7144)	Aylmer	17	30L/13C	[illegible]96300	4743300	S	7.70	0.000	1.000					1.000					NPSW	F	1996DU		1.000	58 LS		F	5a,5b,5c,9							
11	Selkirk Provincial Park (SAC1) (7156)	Aylmer	17	30L/13	584500	4740200	C	32.10	0.070	0.930				0.100	0.500		0.400			PSW	T	1996DU	0.600	0.400	3210 PP		F	5a,5b,5c,6, 8,9		Y					
12	Hickory Creek (SAC3) (7145)	Aylmer	17	40I/16A	580100	4739900	C	3.70	0.140	0.860				0.350	0.650					5	F	1986		1.000			F	5a,5b,5c,9							
13	Nanticoke Creek Mouth (NC4) (7153)	Aylmer	17	40I/16	575800	4739000	S	25.70	0.000	1.000				0.110	0.890					PSW	T	1996DU	0.380	0.620	4425 IBP		F	5a,5b,5c,9			Y			Y	
14	Long Point Wetland Complex (5071)	Aylmer	17	40I/10; 40I/9	554000	4716000	C	13465.00	0.040	0.9597	0.0003		0.050				0.100		0.895	1	T	1987	0.740	0.260	184 MACRO; 2010 NWA; 2080 LS; 3126 PP; 3401 PP-NRZ; 4374 ANSI-LS-P; 5018 ESA; 5079 PP-NEZ	RAMSAR, CAA, PWA, Migratory Bird Sanctuary	T	1,2,3,5a,5b, 5c,6,8,9,11	Y	Y	Y	Y		Y	
15	Big Creek at Port Royal (BC14) (7135)	Aylmer	17	40I/9; 40I/10	541500	4716500	S	110.60	0.330	0.670				0.250	0.750					PSW	T	1996DU	0.100	0.900	2008 NWA		F	3,5a,5b,5c, 9	Y		Y				
16	Rondeau Provincial Park (5649)	Aylmer	17	40I/5	429500	4683300	C	929.58	0.196	0.804			0.097	0.490					0.413	1	T	1983	1.000		2086 PP-NEZ; 2093 PP-NRZ; 3199 PP; 4373 ANSI-LS-P			5,6,9	Y	Y	Y	Y		Y	
17	Rondeau Bay Northwest Shore (7155)	Aylmer	17	40I/5	424000	4683000	C	738.90	0.036	0.964			0.011	0.063		0.132	0.499	0.060	0.237	2	T	1989DU	0.124	0.876	1565 ANSI-LS-R			4,5a,5b,5c, 6,8,9	Y	Y	Y				
18	McGeachy Pond (7151)	Aylmer	17	40I/5	421900	4679300	S	16.00	0.060	0.940								1.000		3	T	1985	1.000			CAA		9		Y	Y				

Appendix K: Lake Erie: Evaluated Coastal Wetlands

									Wetland Type				Site Type																Significant Species						
MAP ID (Figure 4)	WETLAND NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:50 000)	EASTING	NORTHING	SINGLE/COMPLEX	AREA	SWAMP	MARSH	BOG	FEN	ISOLATED	PALUSTRINE	RIVERINE	RIVERINE RIVERMOUTH	LACUSTRINE RIVERMOUTH	LACUSTRINE BAY	LAKE LACUSTRINE	CLASS	PROVINCIALLY SIGNIFICANT WETLAND	YEAROF EVALUATION DATA	PUBLIC OWNERSHIP	PRIVATE OWNERSHIP	CORRESPONDING_NATURAL AREA	OTHER COMMENTS	LOSS	STRESSES	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
19	Wheatley Two Creeks (7159)	Aylmer	17	40J/1	380000	4660000	C	56.00	0.110	0.890								0.730	0.270	3	T	1985	1.000		2088 PP-NEZ; 2095 PP-NRZ; 3255 PP-R; 4992 ANSI-LS-P; 7016 LS			5c,9		Y	Y				
20	Muddy Creek (7152)	Aylmer	17	40J/1	378300	4658300	S	11.00	0.180	0.820					0.120				0.880	3	T	1985	0.600	0.400				2,5c,7			Y				
21	Hillman Marsh (5716)	Aylmer	17	40J/1,2	376500	4654000	S	362.00	0.200	0.800						0.200	0.800			2	T	1983	0.900	0.100	421 ANSI-LS-R; 2048 CAA; 5196 LS	ESA		2,5c	Y	Y	Y	Y			
22	Point Pelee (5646)	Aylmer	17	40G/15,16	374400	4646400	S	1175.00	0.070	0.930								1.000		1	T	1985	1.000		749 LS ANSI-LS-P; 2005 NP; 5197 LS	Ramsar		2	Y	Y	Y	Y	Y	Y	Y
23	Sturgeon Creek (7157)	Aylmer	17	40J/2	370200	4652400	S	31.00	0.160	0.840							1.000			3	T	1985	0.750	0.250				5,6,8		Y	Y				
24	Lighthouse Point (Nature Reserve) (7147)	Aylmer	17	40G/15	363800	4631400	S	85.00	0.200	0.800									1.000	3	T	1984	1.000		75 ANSI-LS-P; 3117 PP-NR; 5192 LS	ESA			Y	Y		Y			
25	Fish Point (Nature Reserve) (5682)	Aylmer	17	40G/10	361000	4621200	C	45.00	0.500	0.500			0.500					0.500		3	T	1984	1.000		470 ANSI-LS-P; 3063 PP-NR; 5191 LS	ESA			Y	Y	Y	Y			
26	Cedar Creek (5705)	Aylmer	17	40J/2	351000	4653000	S	250.00	0.050	0.950					0.900		0.100			3	T	1984	0.900	0.100	37 ANSI-LS-P; 2038 CAA; 5202 LS			5,6	Y	Y	Y			Y	
27	Fox Creek (7143)	Aylmer	17	40G/15	347300	4651000	C	19.00	0.260	0.740						1.000				3	T	1985	0.490	0.510	2044 CAA			5b,5c,6		Y	Y				
28	Lypps Beach Wetland Complex (7148)	Aylmer	17	40J/2	336800	4652000	C	12.80	0.520	0.480								1.000		3	T	1989		1.000				3,5c,7		Y	Y				
29	Big Creek Marsh (7136)	Aylmer	17	40J/3	330000	4656000	S	1000.00	0.100	0.900						1.000				1	T	1984	0.250	0.750	23 ANSI-LS-P; 5201 LS	ESA		1,5,9	Y	Y	Y	Y			
30	Man's Marsh (7149)	Aylmer	17	40J/3	326200	4657600	S	55.00	0.350	0.650								1.000		2	T	1985		1.000				1,5c,9	Y	Y	Y				
				Total Wetland Area				**19329.84**																											

Appendix L: Lake Erie: Natural Areas, Environmental Sensitivity Atlas Wetlands and Other Wetlands with an Unevaluated Coastal Wetland Component

									Significant Species						
MAP ID	AREA NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:25 000)	EASTING	NORTHING	DESCRIPTION	OTHER CORRESPONDING NATURAL AREAS	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
Natural Area Wetlands															
N1	Low Point LS (5187)	Cambridge	17	30L/13	609510	4744170	-graminoid coastal meadow marsh (specific area not included in wetland evaluation)	3088 PP		Y		Y			
N2	Mowhawk Island NWA (2011)	Cambridge	17	30L/13	620700	4743200	-vegetation is scattered with few shrubs, willow clumps, sedges, grasses and mosses	4589 ANSI-LS-R; 5184 LS							
N3	Iroquois Beach ANSI-LS-R (4584)	Aylmer	17	40I/10	515000	4721000	-wet sedge meadows, marshes and shrub thickets	3184 PP; 3505 PP-NRZ; 5128 LS; 5297 IBP; 5300 IBP		Y		Y			

Appendix M: Lake Ontario and Niagara River: Evaluated Coastal Wetlands

									Wetland Type				Site Type																Significant Species						
MAP ID (Figure 5)	WETLAND NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:50 000)	EASTING	NORTHING	SINGLE/COMPLEX	AREA	SWAMP	MARSH	BOG	FEN	ISOLATED	PALUSTRINE	RIVERINE	RIVERINE RIVERMOUTH	LACUSTRINE RIVERMOUTH	LACUSTRINE BAY	LAKE LACUSTRINE	CLASS	PROVINCIALLY SIGNIFICANT WETLAND	YEAROF EVALUATION DATA	PUBLIC OWNERSHIP	PRIVATE OWNERSHIP	CORRESPONDING NATURAL AREA	OTHER COMMENTS	LOSS	STRESSES	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
1	Parrott Bay Wetland (7035)	Peterborough	18	31C/2	364800	4897500	S	30.00		1.000					0.06				0.94	3	T	1985	1.000			CAA	F	4,11		Y	Y				
2	Bath Point Wetland (7067)	Peterborough	18	31C/2	358800	4894800	C	13.80	0.430	0.570				0.96					0.04	2	T	1991		1.000		*Unofficial Complex									
3	Long Point Marsh (7224)	Peterborough	18	31C/2	363500	4886500	C	315.00	0.390	0.590	0.02							1.00		2	T	1986	0.600	0.400	1010 ANSI-LS-P		F		Y		Y				
4	Nut Island Duck Club Marsh (7227)	Peterborough	18	31C/2	359000	4885200	S	114.00	0.600	0.400				0.35	0.15			0.50		2	T	1986	0.400	0.600							Y				
5	Wemps Bay Marsh (7226)	Peterborough	18	31C/2	357700	4886700	S	43.00	0.190	0.810								1.00		3	T	1986		1.000							Y				
6	Adolphustown Marsh (7061)	Peterborough	18	31C/3	338500	4879900	C	28.50	0.030	0.970				0.02				0.08	0.90	PSW	T	1993	0.840	0.160		SLPC	F	1,5b,5c,7,9		Y	Y				
7	Carnachan Bay Wetland Complex (7055)	Peterborough	18	31C/3	336000	4881500	C	71.00	0.030	0.970									1.00	PSW	T	1993	0.770	0.230			F	1,2,5b,5c,7,9			Y				
8	Hay Bay Marsh (5083)	Peterborough	18	31C/2	346000	4893000	C	1333.00	0.070	0.930				0.01	0.02	0.05			0.92	PSW	T	1993	0.800	0.200	1293 ANSI-LS-P	Portion in stewardship agreement	F	1,2,5b,5c,7,9		Y	Y				Y
9	Lower Napanee River (7049)	Peterborough	18	31C/2	341000	4896000	S	206.00	0.280	0.720							0.10		0.90	PSW	T	1993	0.650	0.350		Portion in stewardship agreement	F	1,2,5b,5c,6,7,9	Y		Y				Y
10	Forester's Island (7047)	Peterborough	18	31C/3	335500	4893700	S	14.50	0.030	0.970									1.00	PSW	T	1993	0.280	0.720			F	1,5b,5c,9		Y	Y				
11	Airport Creek Marsh (7062)	Peterborough	18	31C/3	332300	4893800	S	29.00	0.350	0.650				0.36					0.64	PSW	T	1993	0.060	0.940			F	1,4,5b,5c,6,7,9	Y		Y				
12	Lower Sucker Creek (7044)	Peterborough	18	31C/3	330300	4893300	C	75.60	0.420	0.580				0.03	0.29		0.47		0.21	PSW	T	1993	0.330	0.670		Navigable Water	F	1,2,5b,5c,7,9	Y	Y	Y				Y
13	Bluff Point (7054)	Peterborough	18	31C/3	325000	4892000	C	41.50	0.390	0.610				0.60					0.40	PSW	T	1993		1.000			F	1,4,5b,5c,7,9	Y	Y	Y				
14	Big Marsh (Marysville Creek) (7053)	Peterborough	18	31C/3	324000	4893500	S	400.00	0.570	0.430				0.25			0.72	0.03		PSW	T	1995		1.000	138 MACRO		F	1,5b,5c,7,9	Y	Y	Y				
15	Lower Salmon River (7052)	Peterborough	18	31C/3	320000	4894000	C	151.00	0.110	0.890				0.10	0.20		0.66		0.04	PSW	T	1993	0.390	0.610			F	1,2,5b,5c,6,7,9	Y	Y	Y				Y
16	Blessington Creek Marsh (5087)	Peterborough	18	31C/3	314600	4892600	S	75.00	0.130	0.870					0.10			0.80	0.10	PSW	T	1993	0.800	0.200	138 MACRO; 1537 LS		F	1,2,5b,5c,9		Y	Y				
17	Bell Creek Swamp Complex (7050)	Peterborough	18	31C/3	313500	4894000	C	88.10	0.580	0.420				0.43	0.56		0.01			4	F	1992	0.060	0.940			T	2,4,5b,5c,6,9							
18	Belleville Marsh (7065)	Peterborough	18	31C/3	312600	4892500	S	28.00	0.150	0.850				0.10					0.90	PSW	T	1993	0.700	0.300		Portion owned by municipality	T	1,5a,5b,5c,6,9		Y	Y				
19	Belleville Treatment Plant Marsh (7066)	Peterborough	18	31C/3	308500	4891200	S	2.50	0.200	0.800									1.00	NPSW	F	1993	1.000				T	1,2,5,6,9			Y				
20	Bayside Wetland (7063)	Peterborough	18	31C/4	299500	4887800	S	19.00	0.160	0.840				0.10					0.90	PSW	T	1993	0.500	0.500			T	1,2,6,9			Y				
21	Dead Creek Marsh (7037)	Peterborough	18	31C/4	289900	4880500	C	359.00	0.660	0.340				0.80	0.15				0.05	PSW	T	1993	0.400	0.600			F	1,2,5b,5c,6,7,8, 9		Y	Y				Y
22	Carrying Place & 12 O'Clock Point (7056)	Peterborough	18	31C/4	293300	4881200	S	62.00	0.350	0.650				0.02					0.98	PSW	T	1993	0.800	0.200			F	1,2,4,5b,5c,6,9		Y	Y				Y
23	Pine Point Wetland (7051)	Peterborough	18	31C/3	300500	4886500	S	25.80	0.280	0.720				0.14					0.86	NPSW	F	1993	0.350	0.650			F	1,2,4,5a,5b,5c,7			Y				
24	Sawguin Creek Marsh (5074)	Peterborough	18	31C/3	310300	4885700	S	2093.000	0.110	0.890					0.55		0.10		0.35	PSW	T	1993	0.450	0.550	138 MACRO; 1311 ANSI-LS-P; 4323 LS	Portion in stewardship agreement	T	1,2,4,5b,5c,7,9		Y	Y				
25	Big Island Marsh (5073)	Peterborough	18	31C/3	322000	4886000	S	858.00	0.010	0.990									1.00	PSW	T	1993	1.000		138 MACRO; 5016 ANSI-LS-R	Portion in stewardship agreement	F	1,4,5b,5c,7,9		Y	Y	Y			
26	Robinson Cove Marsh (7046)	Peterborough	18	31C/3	317500	4886700	C	9.60		1.000				0.16				0.12	0.72	PSW	T	1993	0.330	0.670			F	1,2,5b,5c,7,9		Y	Y				

Appendix M: Lake Ontario and Niagara River: Evaluated Coastal Wetlands

									Wetland Type				Site Type																Significant Species						
MAP ID (Figure 5)	WETLAND NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:50 000)	EASTING	NORTHING	SINGLE/COMPLEX	AREA	SWAMP	MARSH	BOG	FEN	ISOLATED	PALUSTRINE	RIVERINE	RIVERINE RIVERMOUTH	LACUSTRINE RIVERMOUTH	LACUSTRINE BAY	LAKE LACUSTRINE	CLASS	PROVINCIALLY SIGNIFICANT WETLAND	YEAROF EVALUATION DATA	PUBLIC OWNERSHIP	PRIVATE OWNERSHIP	CORRESPONDING NATURAL AREA	OTHER COMMENTS	LOSS	STRESSES	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
27	North Port Swamp (7045)	Peterborough	18	31C/3	326000	4889800	S	27.00	0.680	0.320					1.00					NPSW	F	1993		1.000			F	1,5b,5c,7,9		Y					
28	Cressy Swamp (7048)	Peterborough	18	31C/2	348700	4882000	S	127.70	0.540	0.460				0.95					0.05	NPSW	F	1993		1.000					Y		Y				
29	Lost Lake Swamp (7068)	Peterborough	18	31C/2	346700	4879000	S	20.00	0.550	0.450				1.00						6	F	1984			1380 ANSI-LS-P							Y			
30	Waupoos Creek Swamp (7075)	Peterborough	18	31C/3A	338100	4874500	S	48.00	1.000					1.00						7	F	1984													
31	South Bay Marsh (7078)	Peterborough	18	30N/14	336000	4864000	S	62.00	0.030	0.970									1.00	2	T	1986	0.900	0.100						Y	Y				
32	Big Sand Bay (7074)	Peterborough	18	30N/15	345000	4865000	S	122.00	0.110	0.890				0.40				0.60		3	T	1986	0.600	0.400			F			Y	Y				
33	Soup Harbour (7077)	Peterborough	18	30N/14	325000	4859500	S	69.00	0.650	0.350				0.40				0.60		5	F	1986		1.000					Y						
34	Salmon Point (7079)	Peterborough	18	30N/14	322000	4858000	S	72.00	0.350	0.650				0.40				0.60		3	T	1986		1.000							Y				
35	East Lake Marsh (7071)	Peterborough	18	30N/14	323000	4865800	S	230.20	0.050	0.950									1.00	1	T	1991	1.000								Y				
36	West Lake Wetland (7076)	Peterborough	18	30N/14	318000	4869500	S	706.00	0.160	0.840					0.60		0.10	0.30		2	T	1984	0.300	0.700	1072 ANSI-LS-R					Y	Y			Y	
37	Hucyks Bay (7069)	Peterborough	18	30N/14	302300	4868500	S	245.00	0.560	0.440					0.90		0.10			2	T	1984									Y				
38	Pleasant Bay (7080)	Peterborough	18	30N/13& 14	300000	4870400	S	299.00	0.100	0.900								1.00		2	T	1985	0.850	0.150					Y		Y				
39	Wellers Bay Wetland Complex (7222)	Peterborough	18	31C/4	293100	4877600	C	363.00	0.190	0.810					0.38				0.62	PSW	T	1993	0.240	0.760	1695 ANSI-LS-P; 2014 NWA				Y		Y				
40	Presqu'ile Bay Marsh (5766)	Peterborough	18	31C/4; 30N/13	282000	4875000	S	970.00	0.170	0.800		0.03						0.06	0.94	1	T	1986	0.800	0.200	8 ANSI-LS-P; 2092, 4365 LS; 3187 PP				Y	Y	Y				
41	Hunt & Beach Road- Popham Bay (7040)	Peterborough	18	31C/4	276500	4876600	S	51.20	0.550	0.450				0.50				0.50		PSW	T	1994		1.000			F				Y				
42	Spencer Point (7042)	Peterborough	18	31C/4	274000	4876000	S	90.40	0.720	0.280				0.79				0.21		PSW	T	1994		1.000			F	4			Y				
43	Colborne Creek (7073)	Peterborough	18	30N/13	267500	4873500	S	18.00	0.330	0.670				0.10	0.20		0.10	0.60		3	T	1986		1.000			F	11							
44	Wicklow Gravel Pit (7043)	Peterborough	18	30N/13	262600	4873300	S	2.19	0.100	0.900								1.00		5	F	1991		1.000		Adjacent to CAA	F			Y					
45	Wicklow Bay (7041)	Peterborough	18	30N/13	260800	4873300	S	47.70	0.790	0.210								1.00		2	T	1991		1.000			T	6							
46	Grafton Swamp (7070)	Peterborough	17	30M/16	740500	4873500	S	61.50	0.900	0.100				0.90			0.10			5	F	1989		1.000			F	4,7	Y	Y				Y	
47	Brookside Wetland (7034)	Peterborough	17	30M/16	734500	4872700	S	20.60	0.570	0.430				1.00						NPSW	F	1994		1.000			F	4	Y			Y			
48	Carr's Marsh (formerly Peter's Rock) (7036)	Peterborough	17	30M/16	723000	4870200	S	60.70	0.520	0.480								1.00		PSW	T	1995	0.56?	0.440	1118 ANSI-LS-P; 7334 IBP	CAA; 56% Public or Private Trust Property	F				Y				
49	Peter Rock #2 (7038)	Peterborough	17	30M/16	720800	4870500	S	2.00	0.100	0.900								1.00		5	F	1991		1.000			F	4			Y				
50	Port Britain (7039)	Peterborough	17	30M/16	711000	4867300	S	17.60	0.380	0.620				0.38		0.62				PSW	T	1994		1.000	1712 ANSI-LS-R; 5712 IBP		T	6,8			Y				
51	Crysler Point Wetland (Wesleyville Marsh) (7072)	Peterborough	17	30M/16	708500	4866100	S	6.10	0.850	0.150								1.00		6	F	1984	1.000		1696 ANSI-LS-R	Ontario Hydro owned	F								
52	Wilmot Rivermouth Wetland (3984)	Aurora	17	30M/15	693000	4863200	C	26.50	0.290	0.710					0.69		0.31			PSW	T	1994	0.490	0.510											
53	Port Darlington Marsh (Bowmanville Marsh) (3914)	Aurora	17	30M/15	687300	4862300	S	24.20		1.000					1.00					6	F	1984		1.000	4042 IBP										
54	West Side Beach Marsh (Pawson Marsh) (3974)	Aurora	17	30M/15	686400	4861800	S	35.70	0.060	0.940							1.00			2	T	1983		1.000							Y				
55	Raby Head Wetland No.2 (3921)	Aurora	17	30M/15	685500	4860800	S	3.10		1.000								1.00		7	F	1984		1.000							Y				
56	Raby Head Wetland No.1 (3920)	Aurora	17	30M/15	684900	4860500	S	4.30	0.350	0.650								1.00		6	F	1984		1.000							Y				

Appendix M: Lake Ontario and Niagara River: Evaluated Coastal Wetlands

									Wetland Type				Site Type																Significant Species						
MAP ID (Figure 5)	WETLAND NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:50 000)	EASTING	NORTHING	SINGLE/COMPLEX	AREA	SWAMP	MARSH	BOG	FEN	ISOLATED	PALUSTRINE	RIVERINE	RIVERINE RIVERMOUTH	LACUSTRINE RIVERMOUTH	LACUSTRINE BAY	LAKE LACUSTRINE	CLASS	PROVINCIALLY SIGNIFICANT WETLAND	YEAROF EVALUATION DATA	PUBLIC OWNERSHIP	PRIVATE OWNERSHIP	CORRESPONDING NATURAL AREA	OTHER COMMENTS	LOSS	STRESSES	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
57	McLaughlin Bay Wetland (3880)	Aurora	17	30M/15	677200	4859500	S	31.20	0.130	0.870									1.00	3	T	1984	1.000		3045 PP; 2085 PP-NEZ						Y				
58	Oshawa Second Marsh (3908)	Aurora	17	30M/15	675700	4860100	S	104.90	0.290	0.710							1.00			2	T	1983	1.000		1502 ANSI-LS-P	CAA				Y	Y				
59	Pumphouse Marsh (Oshawa Filtration Plant) (3918)	Aurora	17	30M/15	673600	4858300	S	2.10	0.200	0.800								1.00		3	T	1983	1.000							Y	Y				
60	Corbett Creek Mouth Marsh (3802)	Aurora	17	30M/15	669600	4857700	S	17.40	0.170	0.830							0.70	0.30		3	T	1987	0.770	0.250	4049 IBP						Y				
61	Lynde Creek Marsh (3874)	Aurora	17	30M/15	663800	4857400	S	110.00	0.400	0.600							1.00			1	T	1983	0.800	0.200	1496 ANSI-LS-R; 2053 CAA; 5713 IBP					Y	Y	Y			
62	Cranberry Marsh (LeVays Marsh) (3804)	Aurora	17	30M/15	663600	4856300	S	32.50	0.320	0.680								1.00		2	T	1983	?	?	1363 ANSI-LS-P; 2053 CAA	ESA				Y	Y				
63	Carruthers Creek Marsh (Shoal Pt Marsh) (3790)	Aurora	17	30M/15	661800	4854700	S	12.90		1.000								1.00		3	T	1983		1.000	1599 ANSI-LS-R	ESA; CAA					Y				
64	Duffin Creek (3815)	Aurora	17	30M/14	657700	4854100	S	61.00	0.100	0.900							1.00			3	T	1983		1.000	1191 ANSI-LS-R						Y				
65	Frenchman's Bay Marshes (3827)	Aurora	17	30M/14	653200	4853700	C	45.20	0.120	0.880							0.44	0.56		2	T	1983	NA	NA		ESA					Y				
66	Rouge River Marshes (3929)	Aurora	17	30M/14	651200	4850700	C	67.98	0.410	0.590							1.00			PSW	T	1995	NA	NA	105 ANSI-LS-P; 1383 ANSI-LS-R; 1569 ANSI-ES-P; 1571 LS; 1572 ANSI-ES-P; 2028 MP; 5671 LS	ESA	F	5b,5c,6,9		Y	Y				
67	Highland Creek Wetlands (3844)	Aurora	17	30M/14	649500	4847900	C	7.60	0.280	0.720				0.13	0.39		0.48			PSW	T	1994	1.000		1300 ANSI-LS-R	CAA	T	5b,5c,6,9			Y				
68	Toronto Island Wetland Complex (7282)	Aurora	17	30M/11	630500	4830300	C	21.91	0.270	0.730								0.48	0.52	PSW	T	1995	NA	NA	121 ANSI-LS-R	ESA; MP	T	2,5b,5c,6,8,9	Y	Y	Y				
69	Humber River Marshes (3851)	Aurora	17	30M/11	621900	4833100	C	26.26	0.250	0.750							1.00			PSW	T	1995	1.000		65 ESA	CAA	T	5b,5c,6,9		Y	Y				
70	Credit River Marshes (7274)	Aurora	17	30M/12	613000	4823200	C	13.43		1.000					1.00					PSW	T	1996	1.000		1159 ANSI-LS-R	Crownland	F	5c							
71	Rattray Marsh (3923)	Aurora	17	30M/12	612600	4819000	C	9.80	0.180	0.820					0.24			0.76		3	T	1984	1.000		2058 CAA; 1552 ANSI-LS-P		T	4,5,6			Y				
72	Oakville Marsh (Sixteen Mile Creek) (3906)	Aurora	17	30M/5	607500	4811100	C	3.00	0.270	0.730					1.00					5	F	1983	1.000		1492 ANSI-LS-R		F	5			Y				
73	Bronte Marsh (3781)	Aurora	17	30M/5	603900	4805100	S	6.00		1.000					1.00					6	F	1983		1.000	3029 PP-R; 3411 ANSI-LS-P; 5329 LS	ESA	F	5							
74	RBG- Hendrie Valley (Lambs Hollow) (3919)	Aurora	17	30M/5	591000	4794000	S	23.00	0.130	0.870					0.40		0.60			2	T	1984	1.000		1297 ANSI-LS-R		F	5a,5b,5c			Y	Y			
75	Cootes Paradise (7025)	Cambridge	17	30M/5	587500	4791500	C	121.70	0.810	0.190								1.00		1	T	1986	1.000		2070 ANSI-LS-P; 1148 LS			9		Y	Y	Y			
76	VanWagner's Marsh (Redhill Creek Marsh) (7033)	Cambridge	17	30M/5	600500	4789500	C	13.30	0.040	0.960				0.60	0.40					2	T	1987		1.000	5394 LS	ESA		4,5c,6,11			Y	Y			
77	Jordan Station Marsh (7032)	Cambridge	17	30M/3&6	632200	4781800	S	136.20		1.000						0.48	0.52			1	T	1985	1.000		1332 ANSI-LS-P; 5433 LS; 5546 OHF		T	4,5a,5b,5c,8,		Y	Y				
78	Sixteen & Seventeen Mile Creeks Terrace Valleys (7030)	Cambridge	17	30M/3&6	635200	4781000	S	48.10	0.030	0.970					0.98	0.02				2	T	1985	1.000		55 ANSI-LS-R		T	4,6	Y	Y	Y				

Appendix M: Lake Ontario and Niagara River: Evaluated Coastal Wetlands

									Wetland Type				Site Type																Significant Species						
MAP ID (Figure 8)	WETLAND NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:50 000)	EASTING	NORTHING	SINGLE/COMPLEX	AREA	SWAMP	MARSH	BOG	FEN	ISOLATED	PALUSTRINE	RIVERINE	RIVERINE RIVERMOUTH	LACUSTRINE RIVERMOUTH	LACUSTRINE BAY	LAKE LACUSTRINE	CLASS	PROVINCIALLY SIGNIFICANT WETLAND	YEAROF EVALUATION DATA	PUBLIC OWNERSHIP	PRIVATE OWNERSHIP	CORRESPONDING NATURAL AREA	OTHER COMMENTS	LOSS	STRESSES	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
79	Fifteen Mile Creek (7027)	Cambridge	17	30M/3&6	636900	4780400	S	31.20	0.300	0.700						1.00				1	T	1987		1.000	55 ANSI-LS-R	Niagara Parks Commission	F	4,9,11		Y		Y			
80	Martindale & Barnesdale Marsh (7031)	Cambridge	17	30M/3&6	640600	4781500	S	55.10		1.000					0.26			0.74		1	T	1985	1.000		5472 LS		T	1,2,5a,5c,5b			Y				
81	Eight Mile Creek Estuary (7026)	Cambridge	17	30M/3&6	647500	4787600	S	5.10	0.580	0.420					0.64	0.36				3	T	1986		1.000	5365 ANSI-LS-R		F	2,4,6,7	Y		Y				
82	Four Mile Creek Estuary (includes Two Mile Pond) (7028)	Cambridge	17	30M/3&6	652400	4790700	S	9.00	0.190	0.810						1.00				3	T	1986	0.600	0.400	1676 ANSI-LS-P; 5454 LS	CAA	F	7,11		Y	Y				
				Total Wetland Area				**11334.47**																											
83	Navy Island Marsh (7210)	Cambridge	17	30M/3&6	662000	4768500	C	25.78	0.310	0.690				0.27	0.73					3	T	1987		1.000	1460 ANSI-LS-P; 4485 IBP; 5442 LS		F			Y					
84	Miller's Creek (7202)	Cambridge	17	30L/15	665300	4756400	C	17.10	0.713	0.287				0.73	0.23	0.04				3	T	1985	0.100	0.900	4483 IBP		F	11	Y	Y					
85	Frenchman's Creek (7212)	Cambridge	17	30L/15	668000	4756000	S	4.50	0.330	0.670					1.00					2	T	1986		1.000	5409 LS	Friends of Frenchman's Creek	F		Y		Y				
86	Black Creek Wetland (7216)	Cambridge	17	30L/14	661300	4758300	S	37.20	0.475	0.525					1.00					2	T	1986		1.000			T	2,6,10		Y				Y	
				Total Wetland Area				**84.58**																											

Appendix N: Lake Ontario and Niagara River: Natural Areas, Environmental Sensitivity Atlas Wetlands and Other Wetlands with an Unevaluated Coastal Wetland Component

									Significant Species						
MAP ID	AREA NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:25 000)	EASTING	NORTHING	DESCRIPTION	OTHER CORRESPONDING NATURAL AREAS	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
Natural Area Wetlands															
N1	Cape Vasey Escarpment Cliffs & Coastal Wetland ANSI-LS-R (1113)	Peterborough	18	31C/2	347500	4877500	Well developed coastal escarpment, wetland (marsh) and shoreline. Partially protected by the Prince Edward Region Conservation Authority.	N/A							
N2	Sandbanks PP-NE (3204)	Peterborough	18	30N/14	318011	4864899	-graminoid coastal meadow marsh type	N/A	Y						
Environmental Sensitivity Atlas Wetlands															
A1	Joshua's Creek ANSI-ES-P (4223)	Aurora	17	30M/5	611000	4815100		N/A							
Other Reported Wetlands															
O1	Fudger's Marsh	Aurora	17	30M/12	612600	4820200	Wetland undergoing evaluation. Single wetland approximately 1.9 ha. Type: 100% marsh; Site Type: lacustrine rivermouth. Not provincially significant. 100% public ownership.	N/A							
O2	Turtle Creek - Reed Swamp	Aurora	17	30M/12	612600	4819600	Wetland undergoing evaluation. Single wetland approximately 2.8 ha. Type: 89% marsh, 11% swamp; Site Type: 86% lacustrine rivermouth, 14% lacustrine bay. Possibly provincially significant. 100% public ownership.	N/A							

Appendix O: St. Lawrence River: Evaluated Coastal Wetlands

									Wetland Type				Site Type																Significant Species						
MAP ID (Figure 6)	WETLAND NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:50 000)	EASTING	NORTHING	SINGLE/COMPLEX	AREA	SWAMP	MARSH	BOG	FEN	ISOLATED	PALUSTRINE	RIVERINE	RIVERINE RIVERMOUTH	LACUSTRINE RIVERMOUTH	LACUSTRINE BAY	LAKE LACUSTRINE	CLASS	PROVINCIALLY SIGNIFICANT WETLAND	YEAROF EVALUATION DATA	PUBLIC OWNERSHIP	PRIVATE OWNERSHIP	CORRESPONDING NATURAL AREA	OTHER COMMENTS	LOSS	STRESSES	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
1	Bainsville Bay Marsh (7535)	Kemptville	18	31G/1	548000	5002100	S	407.20	0.33	0.67					0.05	0.95				PSW	T	1995	0.500	0.400	1535 LS		T	1,2,8			Y				
2	Westleys' Creek Marsh (7536)	Kemptville	18	31G/1	544500	5000500	S	17.20	0.50	0.50					0.40	0.60				PSW	T	1995	0.420	0.580			T	1,2,8			Y				
3	Charlottenburg Marsh (7546)	Kemptville	18	31G/2	536500	4990500	C	851.00	0.10	0.90					0.93	0.70				1	T	1984	0.700	0.300		ESA; CAA; DU-EHJV Agreement	T	1			Y				
4	Hoople Bay Marsh (7572)	Kemptville	18	31G/2	504500	4987000	S	79.60	0.44	0.56					0.95	0.50				PSW	T	1994	0.350	0.650	1308 ANSI-LS-R	PWA; SLPC (50%); Ontario Hydro owns portion	T	1,2			Y				
5	Upper Canada Migratory Bird Sanctuary (7591)	Kemptville	18	31B/14	496600	4977800	C	321.00	0.29	0.71				0.050	0.87	0.08				1	T	1984	0.760	0.240	1136 ANSI-LS-R	Owned by SLPC and Ontario Hydro; MNR-DU-SLPC Management Agreement in part; MBS	F	11			Y				
6	Riverside Marsh (7599)	Kemptville	18	31B/14; 31B/11	489500	4974200	C	134.00	0.12	0.88					1.00					3	T	1985	1.000	0.000	1557 ANSI-LS-R	Most owned by SLPC	T	1			Y				
7	Morrisburg Swamp (7596)	Kemptville	18	31B/14; 31B/11	485000	4973500	C	390.90	0.96	0.04				0.890	0.11					5	F	1985	0.097	0.903			F	6							
8	Doran Creek Marsh (7553)	Kemptville	18	31B/14; 31B/11	477000	4966600	C	42.20		1.00				0.200	0.55	0.25				3	T	1984	1.000	0.000			F	5b,5c							
9	McLachlan Creek Complex (7593)	Kemptville	18	31B/14	467000	4958300	C	22.20		1.00					1.00					3	T	1984	0.000	1.000			T	2,5c,6,8							
10	Johnstown Marsh Complex (7573)	Kemptville	18	31B/14; 31B/11	465000	4956000	C	186.20		1.00					1.00					NPSW	F	1993	1.000	0.000			T	6,8							
11	Bradleys Creek Marsh (7541)	Kemptville	18	31B/12	457300	4949800	C	3.80	0.11	0.89					0.63	0.37				NPSW	F	1993	0.580	0.420			F	4,5b,5c,							
12	Grant's Creek (7554)	Kemptville	18	31B/12	443000	4935000	S	10.00		1.00					1.00					5	F	1984	0.400	0.600			T	1,2,5c,6,8							
13	Molly's Gut Complex (7595)	Kemptville	18	31B/12	441500	4933000	C	6.50	0.10	0.90					1.00					6	F	1984	1.000	0.000			F	1							
14	Jones Creek Marsh (7574)	Kemptville	18	31B/12; 31B/5	434200	4927500	S	140.00	0.05	0.95					1.00	1.00				2	T	1984	0.800	0.200	143 ANSI-LS-P		F	1		Y	Y	Y			
15	Browns Bay Complex (7544)	Kemptville	18	31B/5	434500	4925500	C	140.00		1.00					1.00					3	T	1984	0.900	0.100	2007 NP		T	1,2,4,5							
16	Grenadier Island Complex (7556)	Kemptville	18	31B/5	429000	4915000	C	868.00	0.05	0.95					1.00					1	T	1984	0.800	0.200	1272 ANSI-LS-P; 2007 NP; 5022 IBP		T	1,2,3,4,5c,6							
17	La Rue Mills Wetland Complex (7557)	Kemptville	18	31B/5	425500	4918500	C	218.00	0.31	0.69				0.240	0.76					3	T	1985	0.050	0.950			T	5b,5c,7				Y			
18	Hill Island East (7559)	Kemptville	18	31B/5	424500	4912500	S	72.00		1.00					1.00					4	F	1984	1.000	0.000	142 ANSI-LS-P; 2007 NP; 5024 IBP		F	1			Y				
19	Hill Island West (7558)	Kemptville	18	31B/5	422200	4910700	S	20.00		1.00					1.00					6	F	1984	1.000	0.000	142 ANSI-LS-P; 2007 NP; 5024 IBP		F	1,2							
20	Ivy Lea Wetland Complex (7569)	Kemptville	18	31C/8	417700	4911600	C	63.70	0.07	0.93				0.060	0.810	0.130				PSW	T	1991DU	0.760	0.240			T	1,2,3,5a,5c,6				Y			
21	Collier Island Wetland (7540)	Kemptville	18	31C/8	415500	4910900	S	16.00		1.00					1.000					5	F	1984	0.500	0.500			F	1,2							

Appendix O: St. Lawrence River: Evaluated Coastal Wetlands

									Wetland Type				Site Type																Significant Species						
MAP ID (Figure 6)	WETLAND NAME (NATURAL AREA NUMBER)	MNR DISTRICT	UTM ZONE	MAP SHEET (1:50 000)	EASTING	NORTHING	SINGLE/COMPLEX	AREA	SWAMP	MARSH	BOG	FEN	ISOLATED	PALUSTRINE	RIVERINE	RIVERINE RIVERMOUTH	LACUSTRINE RIVERMOUTH	LACUSTRINE BAY	LAKE LACUSTRINE	CLASS	PROVINCIALLY SIGNIFICANT WETLAND	YEAROF EVALUATION DATA	PUBLIC OWNERSHIP	PRIVATE OWNERSHIP	CORRESPONDING NATURAL AREA	OTHER COMMENTS	LOSS	STRESSES	VEGETATION COMMUNITY	PLANT	BIRD	REPTILE/AMPHIBIAN	MAMMAL	FISH	LEPIDOPTERAN
22	Landons Bay (7581)	Kemptville	18	31C/8	415200	4911500	S	45.70		1.00					1.000					2	T	1991	1.000	0.000	145 IBP; 1356 ANSI-LS-P; 5026 ANSI-ES-P; 5280 IBP		T	1,2,6			Y				
23	Stave Island Wetland Complex (7585)	Kemptville	18	31C/8	414500	4909700	C	8.50		1.00					1.000					5	F	1984	1.000	0.000			F	1							
24	Halsteads Creek Marsh (7563)	Kemptville	18	31C/8	414200	4911600	C	8.10	0.23	0.77					0.280	0.720				5	F	1991	0.700	0.300			F	1		Y					
25	Halsteads Bay Marsh (7561)	Kemptville	18	31C/8	412500	4910500	C	64.00		1.00					1.000					4	F	1984	0.500	0.500			T	1,2,6							
26	Gray's Creek (7555)	Kemptville	18	31C/8	411500	4910500	S	21.00		1.00					1.000					5	F	1984	0.000	1.000			T	1,6							
27	Legges Creek (7582)	Kemptville	18	31C/8	410500	4910500	S	24.50	0.05	0.95					1.000					5	F	1984	0.000	1.000			F	1							
28	Hay Island Marsh (7568)	Kemptville	18	31C/8	408500	4907000	S	59.90		1.00					1.000					6	F	1984	1.000	0.000	2007 NP		F	1							
29	Bostwick Island Complex (7537)	Kemptville	18	31C/8	406300	4906200	C	33.00		1.00					1.000					6	F	1984	0.500	0.500	2007 NP		F	1,2							
30	Willowbank Marsh (7590)	Kemptville	18	31C/8	403000	4908000	S	39.50		1.00					1.000					3	T	1984	0.300	0.700							Y				
31	Firman's Creek Wetland (7600)	Peterborough	18	31C/8	401500	4907000	S	9.75		1.00					1.00					5	F	1985	1.000	0.000											
32	Seburns Creek (7584)	Peterborough	18	31C/8	402200	4905500	S	60.00	0.08	0.92					0.770	0.230				5	F	1985	0.950	0.050						Y					
33	Johnson Bay (7571)	Peterborough	18	31C/8	398900	4904000	S	265.00	0.02	0.98					0.700	0.300				1	T	1985	0.900	0.100	1329 ANSI-LS-R						Y				
34	Grass Creek (7560)	Peterborough	18	31C/8	395300	4905300	S	40.00	0.11	0.89					0.270	0.730				5	F	1985	0.750	0.250							Y				
35	Pitts Ferry Wetland	Peterborough	18	31C/8	394200	4903500	S	3.60		1.00					1.000					5	F	1985	0.000	1.000											
36	Cassidy's Bay (7538)	Peterborough	18	31C/8	393400	4901600	S	3.71	0.39	0.61								1.000		2	T	1989	0.500	0.500							Y				
37	Madoma Marsh (7587)	Peterborough	18	31C/8	390000	4901800	S	60.00		1.00					0.650	0.350				3	T	1985	1.000	0.000											
38	Wolfe Island Wetland Complex (7588)	Peterborough	18	31C/1	390000	4895000	C	1398.00	0.20	0.78	0.02			0.010	0.690		0.010	0.250	0.040	2	T	1986	0.730	0.270	1061 ANSI-LS-P; 4016 ESA; 7772 IBP	EHJV Site			Y	Y	Y				
39	Greater Cataraqui Marsh (6230)	Peterborough	18	31C/8	380000	4903000	S	504.00		1.00						1.000				1	T	1990	1.000	0.000	137 ANSI-LS-P	ESA		2,4,5b,5c,6,11		Y	Y				
40	Little Cataraqui Marsh Complex (7592)	Peterborough	18	31C/2; 31C/7;	379000	4904000	C	359.90	0.28	0.72				0.29	0.44	0.17				1	T	1992	0.710	0.140	2181 CAA	ESA	T	4,5,6,11			Y				
				Total Wetland Area				**7817.66**																											

Appendix P: Confirmed Records of Significant Vascular Plant Species Reported in Ontario's Great Lakes Coastal Wetlands

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	LAKE/CONNECTING CHANNEL	HABITAT DESCRIPTION
Algae-like Pondweed	*Potamogeton confervoides*	S2	-	-	Lake Huron	• shallow water
American Lotus	*Nelumbo lutea*	S2	-	-	Lake Erie, Detroit River, Lake St. Clair, Lake Huron	• shallow water marsh
American Water-willow	*Justicia americana***	S2	THR	VUL	Lake Erie	• shallow water marsh
Angled Spike-rush	*Eleocharis quadrangulata***	S1	-	-	Lake Erie, Lake St. Clair, St. Clair River	• shallow water marsh
Arrow-arum	*Peltandra virginica*	S2	-	-	St. Lawrence River, Niagara River, Lake Huron	• shallow water marsh
Awnless Graceful Sedge	*Carex formosa*	S3S4	-	-	Lake Ontario, Lake Huron, Lake Erie	• mesic woods • floodplains
Bayberry	*Myrica pensylvanica*	S1	-	-	Lake Erie	• deciduous swamp
Big Shellbark Hickory	*Carya laciniosa*	S3	-	-	Niagara River, Lake Huron	• hardwood swamp
Black Sedge	*Carex atratiformis*	S2	-	-	Lake Superior	• coniferous swamp
Bluehearts	*Buchnera americana***	S1	END	-	Lake Huron	• wet meadows
Branched Bartonia (Twining Bartonia)	*Bartonia paniculata spp. paniculata*	S1	SC	-	Lake Huron	• open sedge bog
Branching Burreed	*Sparganium androcladum*	S1	-	-	Lake Erie, Lake Ontario, St. Lawrence River	• shallow water marsh
Bristly Cattail Sedge	*Carex frankii*	S2	-	-	Lake Erie	• floodplain woods • ditches
Bushy Aster	*Aster dumosus*	S2	-	-	Lake Ontario, Lake Erie, Detroit River, Lake St. Clair; Lake Huron	• meadow marsh • moist fresh tall grass prairie
Bushy Cinquefoil	*Potentilla paradoxa***	S3	-	-	Lake Erie	• wet beach
Carey's Smartweed	*Polygonum careyi*	S3S4	-	-	Lake Huron	• meadow marsh
Common Stiff Sedge	*Carex tetanica*	S3	-	-	Lake Erie, Detroit River, Lake Huron	• meadow marsh • seepages
Crested Arrow-head	*Sagittaria graminea* var. *cristata*	S3	-	-	Lake Erie, Lake Huron	• shallow water marsh
Cylindrical Blazing Star	*Liatris cylindracea*	S3	-	-	Lake Huron	• alvar • meadow marsh
Dwarf Lake Iris	*Iris lacustris*	S3	-	-	Lake Huron	• alvar • coniferous swamp • coastal meadow marsh
Eastern Prairie Orchid	*Platanthera leucophaea*	S2	SC	-	St. Lawrence River, Lake Erie, Detroit River, Lake St. Clair; Lake Huron	• meadow marsh • open sedge fen • wet tall grass prairie

Appendix P: Confirmed Records of Significant Vascular Plant Species Reported in Ontario's Great Lakes Coastal Wetlands

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	LAKE/CONNECTING CHANNEL	HABITAT DESCRIPTION
Eastern Yellow Star Grass	*Hypoxis hirsuta*	S3	-	-	Lake Erie, Lake St. Clair, Lake Huron	• mesic prairie
Engelmann's Spike-rush	*Eleocharis engelmannii*	S1	-	-	Lake Erie	• wet beach
Fogfruit	*Phyla lanceolata*	S2	-	-	Lake Erie	• meadow marsh • flood plain
Follicle Sedge	*Carex folliculata*	S3	-	-	St. Lawrence River, Lake Huron	• coniferous swamp • thicket swamp • mixed swamp • shallow water marsh
Gattinger's Agalinis (previously Round-stemmed Purple False Foxglove)	*Agalinis gattingeri*	S1	END	-	Lake St. Clair, Lake Huron	• meadow marsh • alvar • mesic prairie
Giant Ironweed	*Vernonia gigantea*	S3	-	-	Lake Erie, Detroit River, Lake St. Clair, Lake Huron	• mesic prairie
Grass Sedge	*Carex jamesii*	S3	-	-	Lake Ontario, Lake Erie	• floodplain woods • mesic woods
Grass-leaved Rush	*Juncus marginatus*	S2	-	-	Lake Erie	• mesic prairie
Gray-headed Coneflower	*Ratibida pinnata*	S2S3	-	-	Detroit River	• mesic prairie
Great Plains Ladies' Tresses	*Spiranthes magnicamporum*	S3	-	-	Lake Erie	• not a typical wetland species but sometimes found in wetlands • mesic prairie • meadow marsh
Green Dragon	*Arisaema dracontium*	S3	SC	-	Lake Erie	• wooded floodplains
Green Violet	*Hybanthus concolor*	S2	-	-	Lake Ontario	• often found in upland areas, but sometimes in wooded floodplains
Grooved Yellow Flax	*Linum sulcatum*	S3	-	-	Lake Erie	• alvar • mesic prairie
Hairy Fimbristylis	*Fimbristylis puberula***	S1	-	-	Lake St. Clair	• mesic prairie
Hidden-fruited Bladderwort	*Utricularia geminiscapa*	S3	-	-	Lake Ontario, Lake Huron	• open sedge bog • shallow water marsh
Honey Locust	*Gleditsia triacanthos*	S2	-	-	Niagara River, Lake Erie, Detroit River, Lake St. Clair	• deciduous swamp
Horsetail Spike-rush	*Eleocharis equisetoides***	S1	END	END-R	Lake Erie	• shallow water marsh
Houghton's Goldenrod	*Solidago houghtonii*	S2	-	-	Lake Huron	• wet alvar
Knee Spike-rush	*Eleocharis geniculata***	S1	-	-	Lake Erie	• wet beach

Appendix P: Confirmed Records of Significant Vascular Plant Species Reported in Ontario's Great Lakes Coastal Wetlands

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	LAKE/CONNECTING CHANNEL	HABITAT DESCRIPTION
Large Water Starwort	*Callitriche heterophylla*	S2?	-	-	Lake Erie, Lake Huron, Lake Superior	• shallow water marsh • deep water marsh
Leafy Blue-flag	*Iris brevicaulis***	S1	-	-	Lake Erie	• wooded deciduous swamp
Lizard's Tail	*Saururus cernuus*	S3	-	-	Lake Erie	• floodplains
Low Nut-rush	*Scleria verticillata*	S3	-	-	Lake Ontario, Lake Erie, Lake Huron	• coastal meadow marsh • prairie fen
Many-fruited False Loosestrife	*Ludwigia polycarpa*	S2	-	-	Lake Erie, Detroit River; Lake St. Clair, Lake Huron	• shallow water marsh • meadow marsh
Marsh St. John's-wort	*Triadenum virginicum*	S3	-	-	Lake Ontario, Lake Huron (Georgian Bay)	• shallow water marsh • open bog
Mosquito Fern	*Azolla caroliniana*	S1	-	-	St. Lawrence River, Lake Ontario	• shallow water marsh
Narrow-leaved Water-plantain	*Alisma gramineum*	S3S4	-	-	St. Lawrence River, Lake Ontario, Lake Huron	• shallow water marsh
Ohio Spiderwort	*Tradescantia ohiensis*	S2	-	-	Detroit River	• mesic prairie
Pin Oak	*Quercus palustris*	S3	-	-	Niagara River, Lake Erie	• hardwood swamp
Prairie Dropseed	*Sporobolus heterolepis*	S2	-	-	Lake Huron	• primarily upland but occasionally found in wetlands • alvar • prairie
Pumpkin Ash	*Fraxinus profunda*	S2	-	-	Lake Erie, Lake St. Clair, Lake Huron	• deciduous swamp
Purple-jointed Joe Pye Weed	*Eupatorium purpureum*	S3	-	-	Lake Erie, Lake Huron	• mesic floodplain
Ram's-head Lady's Slipper	*Cypripedium arietinum*	S3	-	-	Lake Ontario, Lake Huron, Lake Superior	• mixed swamp • conifer swamp
Red-rooted Nut Sedge	*Cyperus erythrorhizos*	S3	-	-	Lake Ontario, Niagara River, Lake Erie, Lake Huron	• meadow marsh • mud flat • wet beach
Redtop Panic Grass	*Panicum rigidulum*	S2S3	-	-	Lake Erie, Lake Huron	• meadow marsh
Riddell's Goldenrod	*Solidago riddellii*	S2S3	SC	VUL	St. Clair River, Lake Huron	• mesic prairie • meadow marsh
Rigid Yellow Flax	*Linum striatum*	S1	-	-	Lake Huron, Lake Erie	• meadow marsh
Riverbank Sedge	*Carex emoryi*	S3	-	-	Lake Erie, Lake St. Clair, Lake Huron	• meadow marsh • shallow water marsh • wet prairie • flood plains
Rough Water Horehound	*Lycopus asper*	S2	-	-	Lake Huron, Lake Erie, Lake Ontario	• shallow water marsh
Scarlet Ammannia	*Ammannia robusta*	S1	END	-	Lake Erie	• wet beach (mud)

Appendix P: Confirmed Records of Significant Vascular Plant Species Reported in Ontario's Great Lakes Coastal Wetlands

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	LAKE/CONNECTING CHANNEL	HABITAT DESCRIPTION
Seedbox	*Ludwigia alternifolia*	S1	-	-	Detroit River, Lake Erie	• shallow water marsh
Sharp-fruit Rush	*Juncus acuminatus*	S3	-	-	Lake Ontario, Niagara River, Lake Erie, Detroit River, Lake St. Clair, Lake Huron	• wet beach • meadow marsh
Skinner's Agalinis (previously Pale Purple False Foxglove)	*Agalinis skinneriana*	S1	END	-	Lake St. Clair	• meadow marsh • prairie
Slender Bulrush	*Scirpus heterochaetus*	S2	-	-	Lake Huron	• shallow water marsh
Small White Lady's Slipper	*Cypripedium candidum*	S1	END	END-R	Lake Erie	• meadow marsh • fen • mesic prairie
Small-flowered Lipocarpha (previously Dwarf Bulrush)	*Lipocarpha micrantha*	S1	END	THR	Lake Erie	• moist beach
Smith's Tufted Bulrush	*Scirpus smithii*	S2?	-	-	St. Lawrence River, Lake Ontario, Niagara River, Lake Erie, Lake Huron	• wet beach
Southern Tickseed	*Bidens coronata*	S2	-	-	Lake Erie, Detroit River, Lake St Clair, Lake Huron	• meadow marsh
Spotted Pondweed	*Potamogeton pulcher***	SH	-	-	Lake Erie	• shallow water
Stalked Water Horehound	*Lycopus rubellus*	S2	-	-	Lake Erie	• hardwood swamp
Stiff Cowbane	*Oxypolis rigidior*	S2	-	-	Lake Erie	• hardwood swamp • thicket swamp
Stiff Yellow Flax	*Linum medium var. medium*	S3	-	-	Lake Huron	• meadow marsh
Stiff Yellow Flax	*Linum medium var. texanum*	S1	-	-	Lake Erie	• meadow marsh
Swamp Agrimony	*Agrimonia parviflora*	S3S4	-	-	Lake St. Clair	• moist old fields • open mesic woods • scrub
Swamp Oval Sedge	*Carex muskingumensis*	S2	-	-	Lake Erie	• floodplain woods • mesic woods
Swamp Rose-mallow	*Hibiscus moscheutos*	S3	SC	-	Lake Ontario, Niagara River, Lake Erie, Detroit River, Lake St. Clair; St. Clair River	• shallow water marsh • meadow marsh • mixed swamp
Swamp Star Sedge	*Carex seorsa*	S2	-	-	Niagara River, Lake Erie	• deciduous swamp • hardwood swamp • mixed swamp
Tall Yellow-eyed Grass	*Xyris difformis*	S3?	-	-	Lake Huron	• meadow marsh
Thread-like Naiad	*Najas gracillima*	S2	-	-	Lake Erie, Lake Huron	• submerged aquatic • shallow water marsh

Appendix P: Confirmed Records of Significant Vascular Plant Species Reported in Ontario's Great Lakes Coastal Wetlands

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	LAKE/CONNECTING CHANNEL	HABITAT DESCRIPTION
Three-awn Grass	*Aristida longespica var. geniculata*	S2	-	-	Lake Erie	• primarily upland species but sometimes occurs in meadow marsh
Trailing Wild Bean	*Strophostyles hevula*	S3	-	-	Lake Erie, Detroit River	• moist beach
Tuberous Indian-plantain (previously Prairie Indian Plantain)	*Arnoglossum plantagineum (previously Cacalia plantaginea)*	S3	SC	-	Lake Huron	• meadow marsh • alvar • open sedge fen
Virginia Bugleweed	*Lycopus virginicus*	S2	-	-	Lake Erie	• shallow water marsh • perched prairie fen
Virginia Meadow Beauty	*Rhexia virginica*	S3S4	-	-	Lake Ontario, Lake Huron	• meadow marsh
Walter's Barnyard Grass	*Echinochloa walteri*	S3	-	-	Lake Erie, Detroit River, Lake St. Clair; St. Clair River	• meadow marsh
Water Awlwort	*Subularia aquatica*	S3?	-	-	Lake Huron, St. Marys River, Lake Superior	• shallow water marsh • deep water marsh
Wedge-fruited Oval Sedge	*Carex suberecta*	S2	-	-	Detroit River	• wet prairie • deciduous swamp
White-fringed Orchid	*Platanthera blephariglottis*	S3S4	-	-	Lake Ontario, Lake Erie, Lake Huron	• open bog • treed bog
Wicket Spike-rush	*Eleocharis rostellata*	S3	-	-	Lake Erie, Lake Huron (Georgian Bay)	• open sedge fen • coastal meadow marsh • prairie fen
Wiegand's Sedge	*Carex wiegandii*	S1	-	-	Lake Superior	• thicket swamp • treed bog
Wild Hyacinth	*Camassia scilloides*	S2	THR	THR	Lake Erie	• moist woods
Winged Loosestrife	*Lythrum alatum*	S3	-	-	Lake Ontario, Lake Erie, Detroit River, Lake St. Clair, St. Clair River	• meadow marsh • shallow water marsh • deciduous swamp • alvar grassland
Winged Monkey Flower	*Mimulus alatus*	S2	-	-	Lake Erie	• floodplain • wooded deciduous swamp
Winged Oval Sedge	*Carex alata***	S1	-	-	Lake Erie	• hardwood swamp
Wingstem	*Verbesina alternifolia*	S2S3	-	-	Lake Erie	• floodplain woods
Yellow Flat Sedge	*Cyperus flavescens*	S2	-	-	Lake Erie	• meadow marsh • wet beach
Yellow Pond Lily	*Nuphar advena*	S3	-	-	Lake Ontario, Lake Erie, Lake Huron	• shallow water marsh
Yerba de Tajo	*Eclipta prostrata*	S2	-	-	Lake Erie	• wet beach • mud flat

Appendix Q: Significant Fish Species Reported in Ontario's Great Lakes Coastal Wetlands

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	LAKE/ CONNECTING CHANNEL	HABITAT DESCRIPTION
Bigmouth Buffalo	*Ictiobus cyprinellus*	SU	SC	NIAC	Lake Erie	• spawns on flooded vegetation • resides in shallows of sluggish, clear, weedy waters
Black Bullhead	*Ameiurus melas*	S3	-	-	Niagara River, Lake Erie, Detroit River, Lake St. Clair, St. Clair River, Lake Huron	• spawn in moderate to heavy vegetation • like silty soft bottom backwaters
Eastern Silvery Minnow	*Hybognathus regius*	S2	-	NIAC	St. Lawrence River, Lake Ontario	• shallow inshore areas of large lakes and rivers • spawns in heavily vegetated backwaters over bottom muck
Grass Pickerel	*Esox americanus*	S3	-	-	St. Lawrence River, Lake Ontario, Niagara River, Lake Erie, Lake St. Clair, Lake Huron (south)	• spawns on flooded stream margins or in marshes • resides in heavily vegetated lowland streams and marshes
Lake Chubsucker	*Erimyzon sucetta*	S2	THR	THR	Niagara River, Lake Erie, Lake St. Clair, Lake Huron (south)	• spawns on submerged vegetation in marshes • resides in the shallow weedy waters of lakes, streams and backwaters
Longear Sunfish	*Lepomis megalotis*	S3	NAR	NIAC	Lake Erie, Lake St. Clair, Lake Huron (south, central, Georgian Bay)	• nests in shallow gravel areas • prefers clear shallow areas in or near aquatic vegetation
Pugnose Minnow	*Opsopoeodus emiliae*	S2	SC	VUL	Detroit River, Lake St. Clair, St. Clair River	• spawns on aquatic vegetation • prefers sluggish, clear weedy waters
Pugnose Shiner	*Notropis anogenus*	S2	END	THR	St. Lawrence River, Lake Ontario, Lake St. Clair, Lake Huron, Lake Erie	• heavily vegetated waters as well as clear streams of sand and marl • no information on spawning
Spotted Gar	*Lepisosteus oculatus*	S2	SC	THR	Lake Erie	• all life history stages associated with dense submergent vegetation
Warmouth	*Lepomis gulosus*	S1	SC	VUL	Lake Erie	• lakes, ponds and sluggish streams with dense aquatic vegetation • nests in shallow water over rubble, loose vegetation or other debris

Appendix R: Significant Reptile and Amphibian Species Reported in Ontario's Great Lakes Coastal Wetlands

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	LAKE/ CONNECTING CHANNEL	HABITAT DESCRIPTION
Reptiles						
Butler's Gartersnake	*Thamnophis butleri*	S2	THR	VUL	Lake Erie, Detroit River, St. Clair River	• wet meadows and prairies, marshy ponds and lake border and other moist grassy places • in Ontario often found in upland areas adjacent to wetlands
Eastern Fox Snake	*Elaphe vulpina gloydi*	S3	THR	THR	Lake Erie, Detroit River, Lake St. Clair, St. Clair River, Lake Huron (central, Georgian Bay)	• Great Lakes shoreline marshes and vegetated dunes and beaches
Eastern Massasauga Rattlesnake	*Sistrurus catenatus catenatus*	S3	THR	THR	Lake Erie, Detroit River, Lake Huron (central, Georgian Bay, North Channel)	• associated with damp lowlands, including river bottom woodlands, shrub swamps, bogs, fens, marsh borders, sedge meadows and moist prairies
Eastern Spiny Softshell Turtle	*Apalone spinifera*	S3	THR	THR	Lake Ontario, Lake Erie, Detroit River, Lake St. Clair, St. Clair River, Lake Huron (south)	• rivers and larger streams, inland lakes, reservoirs and protected bays and river mouths • prefer sand or mud bottom
Lake Erie Water Snake	*Nerodia sipedon insularum*	S2	END	END-R	Lake Erie (Pelee Is.)	• live on or near permanent bodies of water, rivers, streams, sloughs, lakes, ponds, bogs meadows and swamps
Queen Snake	*Regina septemvittata*	S2	THR	THR	Lake Erie, Detroit River, Lake St. Clair, Lake Huron (south, central)	• in or near warm shallow rocky bottomed streams, also inhabit edges of ponds, lakes, marshes, ditches and canals
Spotted Turtle	*Clemmys guttata*	S3	SC	VUL	St. Lawrence River, Lake Ontario, Lake Erie, Lake St. Clair, Lake Huron (south, central, Georgian Bay)	• shallow ponds, wet meadows, tamarack swamps, bogs fens, marsh channels, sphagnum seepages and slow streams
Amphibians						
Fowler's Toad	*Bufo fowleri*	S2	THR	THR	Lake Erie	• open woodlands, sand prairies, meadows and beaches
Jefferson Salamander	*Ambystoma jeffersonianum*	S2	THR	-	Lake Ontario, Lake Erie, Lake Huron	• most common in moist well drained upland forests, but breeds in ponds within or adjacent to their woodland habitats
Northern Cricket Frog (previously Blanchard's Cricket Frog)	*Acris crepitans*	SH	END	END-R	Lake Erie (Pelee Is.)	• permanent ponds, bogs, lakes and slow moving streams or rivers • abundant aquatic vegetation • sparsely vegetated mud flats and muddy or sandy shorelines are also favoured habitats • in Ontario, only marginally a wetland species
Smallmouth Salamander	*Ambystoma texanum*	S1	SC	THR	Lake Erie (Pelee Is.)	• most abundant in lowland floodplain woods • breed in shallow bodies of water; also breed in runoff ponds, flooding, river backwaters and roadside ditches

Appendix S: Significant Bird Species Reported in Ontario's Great Lakes Coastal Wetlands

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	LAKE/CONNECTING CHANNEL		HABITAT DESCRIPTION
					Breeding	**Non-Breeding**	
Acadian Flycatcher	*Empidonax virescens*	S2B	END	-	• Lake Erie • very few breeding sites along coast • no record of confirmed breeding sites outside the extreme southwest of Ontario		• breeds in river swamps, creek bottoms, well-wooded ravines, Black Ash swamps and deep shaded forests • in Ontario, fairly mature deciduous forest or edges of wood ravines are favoured • Ontario is the northern limit of its range • good quality habitat has been reduced through clearing of forests for agriculture
Bald Eagle	*Haliaeetus leucocephalus*	S3B	NAR	END-R	• breeding sites are not rare but much reduced (especially in southern Ontario) since the 1950s due to shooting, habitat destruction and DDT chemical (primary cause) • Lake Erie, St. Marys River, Lake Superior	• migratory and resting areas are very few • Lake Erie, Lake Ontario, Lake Superior	• stick nests in trees are immediately adjacent to wetlands (shallow water marshes), or on the lakeplain
Black Tern	*Chlidonias niger*	S3B	NAR	VUL	• St. Lawrence River, Lake Ontario, Lake Erie, Lake St. Clair, Lake Huron (Georgian Bay), St. Marys River		• nests amidst moderately dense emergent marsh vegetation- cattails favored • usually open water adjacent to nests • likes productive wetlands with extensive shallow areas and emergent vegetation • commonly abandon breeding sites when vegetation or water levels change • likely to have been a decrease in populations over the last few decades with extensive alteration and destruction of marsh habitat
Black-crowned Night-Heron	*Nycticorax nycticorax*	S3B	-	-	• Lake Ontario, Niagara River, Lake Erie, Lake St. Clair, Lake Huron (& Georgian Bay)		• most Ontario nests were in deciduous shrubs or trees, some in conifers and one other on matted vegetation in a cattail marsh; more usual sites were on islands, wooded river banks or swamps • declines in 1960s, early 1970s due to pesticides; the species is expanding its range but will be limited by loss of wetland habitat
Canvasback	*Aythya valisineria*	S1B	-	-	• Ontario population is very small • rare breeder in Ontario appearing mostly in larger marshes of the Great Lakes	• coastal wetlands are major non-breeding and staging areas • occasionally overwinter in coastal wetlands	• appears mostly in large fertile marshes • almost all nests are placed on emergent vegetation (cattails and bulrushes) primarily, but also sedges and bur-reed • extreme extent of its western range
Cerulean Warbler	*Dendroica cerulea*	S3B	SC	VUL	• Lake Erie, Lake St. Clair		• nest in deciduous swamps

Appendix S: Significant Bird Species Reported in Ontario's Great Lakes Coastal Wetlands

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	LAKE/CONNECTING CHANNEL		HABITAT DESCRIPTION
					Breeding	**Non-Breeding**	
Forster's Tern	*Sterna forsteri*	S3B	DD	IND	• nesting in Ontario confined to coastal wetlands of Lake Erie, Lake St. Clair, St. Clair River and Lake Huron		• deep water portion of large cattail marshes • range expansion and increased population abundance attributed to years of high water levels in Lake Erie • nests primarily in the Prairie Provinces • locally common, rare overall
Great Egret	*Casmerodius albus*	S2B	-	-	• in Ontario, Great Lakes marshes are the primary breeding sites: Lake Erie, Lake St. Clair, Lake Huron (& Georgian Bay)		• frequents marshes and open areas of water • breeds in treed swamps, thickets and occasionally low vegetation • gradual northward extension of its breeding range
King Rail	*Rallus elegans*	S2B	END	END-R	• coastal marshes of Lake Erie, Lake St. Clair, Lake Ontario, Lake Huron		• prefer large expanses of shallow water marsh that merge with scrubby swales • cattail, grass/sedge marshes, shrubby swamps and marshy borders of ponds and rivers • population declines noted in many parts of North America - drainage of wetlands contributed to their decline in Ontario
Least Bittern	*Ixobrychus exilis*	S3B	THR	VUL	• St. Lawrence River, Lake Ontario, Lake Erie, Detroit River, Lake St. Clair, Lake Huron (& Georgian Bay)		• open cattail and *Scirpus* spp. marshes • moderately dense cattail stands and dense reed beds in the larger marshes of the lower Great Lakes provide the most extensive habitat in Ontario • species has decreased in number within its range due to loss of wetlands
Little Gull	*Larus minutus*	S1S2B	-	-	• Lake Erie, Lake Huron (Georgian Bay) • all nesting sites in Great Lakes Basin are coastal • coastal lowlands of James Bay and Hudson Bay may be prime nesting area in North America		• nests on wet grassy knoll or islets of tangled floating vegetation - marshes • most southern Ontario sites are now historical; most active sites are on Lake Huron • chiefly a breeding bird of northern and central temperate Asia • number of nests in North America is slowly increasing
Louisiana Waterthrush	*Seiurus motacilla*	S3B	SC	VUL	• Lake Ontario, Lake Erie, Lake Huron • only a few of the recorded nesting sites are in protected areas		• deep forested ravines with tumbling waters, occasionally forested swamps and linear woodland sloughs • common across eastern U.S.; rare in Ontario - northern extent of the species

Appendix S: Significant Bird Species Reported in Ontario's Great Lakes Coastal Wetlands

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	LAKE/CONNECTING CHANNEL		HABITAT DESCRIPTION
					Breeding	Non-Breeding	
Northern Shoveler	*Anas clypeata*	S3S4B	-	-	• occasional breeder in Great Lakes coastal wetlands: St. Lawrence River, Lake Ontario, Lake Erie, Detroit River, Lake Huron (& Georgian Bay), Lake Superior	• primary use of coastal wetlands is during migration	• short grass areas, large highly productive marshes, sewage lagoons and water impoundments provide nesting habitat • Ontario is the eastern periphery of its breeding area
Peregrine Falcon *anatum* subspecies	*Falco peregrinus anatum*	S2B	THR	END-R	• Lake Superior	• feeds in wetlands particularly during migration	• cliffs and crags situated near water • wetlands important feeding areas during migration • suffered serious decline in 1960s due to DDT pesticide contamination
Prothonotary Warbler	*Protonotaria citrea*	S1S2B	END	END-R	• Lake Erie		• nest in deciduous swamp, Silver Maple-Red Maple swamp, Buttonbush swamp, Black Willow-maple-ash floodplains • nests made of moss - critical habitat requirement • tree cavities used as nest sites • nest nearly always closely associated with an open pool of standing or flowing water • species has a narrow habitat tolerance • number of nesting sites is declining
Redhead	*Aythya americana*	S2B	-	-	• vast majority of Ontario birds breed in Lake St. Clair coastal wetlands • also confirmed breeding sites: St. Lawrence River , Lake Ontario, Lake Erie, and Lake Huron (Georgian Bay) • most breeding sites are in Great Lakes wetlands	• important migration areas	• cattail, bulrush and fens are preferred nesting areas, but will nest on islands and in meadows • nests constructed in emergent vegetation above shallow water • most common in prairies, but may become more widespread in Ontario (like lagoons) • low population abundance across North America
Red-necked Grebe	*Podiceps grisegena*	S3B	NAR	NIAC	• rarely breed in coastal wetlands	• important during spring migration: Lake Ontario, Lake Huron (& Georgian Bay) and possibly Lake Superior	• shallow water marshes • loss of breeding habitat in southern Ontario, specifically due to shoreline development
Ruddy Duck	*Oxyura jamaicensis*	S2B	-	-	• Lake St. Clair, Lake Erie, Lake Huron		• nests are attached to cattails, bulrushes, bur-reeds and sedges • nests usually associated with shallow water (often sewage lagoons) situated in the marshy borders of waterbodies • primarily a prairie species

Appendix S: Significant Bird Species Reported in Ontario's Great Lakes Coastal Wetlands

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	LAKE/CONNECTING CHANNEL		HABITAT DESCRIPTION
					Breeding	Non-Breeding	
Short-eared Owl	*Asio flammeus*	S2N	SC	-	• St. Lawrence River, Lake Ontario • also breeds in James Bay and Hudson Bay	• overwinters in Great Lakes coastal wetlands	• frequents meadows, bogs and marshes • nest is made of dead grass placed among vegetation on wet ground • cosmopolitan - found on all continents except Australia and Antarctica; found in northern and southern Ontario, but its distribution is uneven and highly correlated with its specialized habitat requirements • destruction of wetlands by drainage for agriculture and housing is likely the single most important factor responsible for its steady decline
Tundra Swan	*Cygnus columbianus*	S3B	-	-	• no breeding locations in the Great Lakes	• species migrates across most of Ontario, particularly through the southern Great Lakes (Lake Erie & Lake St. Clair)	• breeding swans were extirpated in Ontario and Hudson Bay during the fur trade; the population is recovering and is expected to expand
White-eyed Vireo	*Vireo griseus*	S2B	-	-	• Lake Erie, Detroit River • most nesting sites in Ontario are coastal; typically in the southwest		• occupies wet and dry habitats - dense shrubbery of overgrown fields, young second growth woodlands, woodlot edges and clearings and open swampy woodlands • concentration in eastern U.S.; has been expanding its range northwards and may become more abundant in the future
Wilson's Phalarope	*Phalaropus tricolor*	S3B	-	-	• only confirmed coastal sites: eastern Lake Ontario and Lake Erie • two probable sites on Lake Huron (& Georgian Bay)		• generally breed in ponds and lakes close to wet meadow vegetation • in southern Ontario most often closely associated with sewage lagoons with lush grassy or sedgy margins or in marshes • historical records indicate this species has only recently become established in southern Ontario
Yellow Rail	*Coturnicops noveboracensis*	S3S4B	SC	VUL	• Lake Ontario • breeds primarily in coastal areas of James Bay and Hudson Bay		• large dense grass and sedge marshes • emergent marsh vegetation • historically had a widespread breeding distribution in Ontario (prior to draining of wetlands)
Yellow-headed Blackbird	*Xanthocephalus xanthocephalus*	S3B	-	-	• Lake Erie and Lake St. Clair	• occasional migrant found in Lake Superior	• permanent deep marshes or sloughs, marshy zones of lakes or river impoundments in which there are stands of cattails or *Phragmites* spp. • spreading south and west into Ontario

Appendix T: Significant Lepidopteran Species Reported in Ontario's Great Lakes Coastal Wetland Evaluations

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	LAKE/ CONNECTING CHANNEL	HABITAT DESCRIPTION
Duke's Skipper	*Euphyes dukesi*	S2	-	-	Lake Erie, Lake St. Clair	• marshy or wet areas with long grass and sedges • larval food plants: *Carex lacustris* and *Carex hyalinolepis*
Mulberry Wing	*Poanes massasoit*	S3	-	-	Lake Ontario, Lake Erie, Detroit River, Lake Huron (south & central)	• marshy or wet areas with long grass and sedges • favourite food: *Carex lacustris*
Two-spotted Skipper	*Euphyes bimacula*	S3S4	-	-	St. Lawrence River, Lake Ontario, Lake Erie, Detroit River, Lake Huron (north and Georgian Bay)	• marshy areas and damp meadows • sedge food: *Carex stricta* marshes

Appendix U: Unconfirmed Records of Significant Vascular Plant Species Reported in OMNR Wetland Evaluations for Great Lakes Coastal Wetlands

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	COMMENTS
American Beach Grass	*Ammophila breviligulata*	S3	-	-	• Non-Wetland species • Dune/ Upland species
American Chestnut	*Castanea dentata*	S3	THR	-	• Non-Wetland species • mesic or upland woods habitat
Bashful Bulrush (previously Small-flowered Clubrush	*Trichophorum planifolium* (previously *Scirpus verecundus)*	S1	END	-	• Non-Wetland species • Upland species
Biennial Gaura	*Gaura biennis*	S2	-	-	• typically a Non-Wetland species • Upland species
Black Cohosh	*Cimicifuga racemosa*	S2	-	-	• typically a Non-Wetland species
Blunt-leaved Spurge	*Euphorbia obtusata*	S1	-	-	• typically a Non-Wetland species
Carolina Gentian	*Frasera caroliniensis*	S1	-	-	• typically a Non-Wetland species
Climbing Prairie Rose	*Rosa setigera*	S3	THR	-	• Non-Wetland species • Upland species
Coast Jointweed	*Polygonella articulata*	S3	-	-	• Non-Wetland species • open sandy habitats
Common Hoptree	*Ptelea trifoliata*	S3	THR	-	• Non-Wetland species • usually Upland species
Dwarf Chinquapin Oak	*Quercus prinoides*	S2	-	-	• Non-Wetland species • Upland species • record probably a misidentification and species actually *Quercus muehlenbergii*
Fern-leaved False Foxglove	*Aureolaria pedicularia*	S3	-	-	• Non-Wetland species • Upland species
Hairy Bedstraw	*Galium pilosum*	S3	-	-	• Non-Wetland Species • usually Upland species
Hoary Tick Trefoil	*Desmodium canescens*	S2	-	-	• Non-Wetland species • Upland species
Large-flowered Purple False Foxglove	*Agalinis purpurea*	S1	-	-	• record probably a misidentification and actually *Agalinis purpurea var. paupercula,* which is not rare
Longleaf Dropseed	*Sporobolus asper*	S1S2	-	-	• Non-Wetland Species • probably non-native at this site

Appendix U: Unconfirmed Records of Significant Vascular Plant Species Reported in OMNR Wetland Evaluations for Great Lakes Coastal Wetlands

COMMON NAME	SCIENTIFIC NAME	NHIC S-RANK	COSEWIC	COSSARO	COMMENTS
Narrow-leaved Sedge	*Carex amphibola*	S2	-	-	• record probably a misidentification and actually *Carex amphibola var. turgida (=Carex grisea,)* which is not rare
Pale Avens	*Geum virginianum*	S1	-	-	• occasionally a Wetland species, probably a misidentification
Panicled Hawkweed	*Hieracium paniculatum*	S2	-	-	• usually a Non-Wetland species
Perfoliate Bellwort	*Uvularia perfoliata*	S1	-	-	• Non-Wetland species • Upland species
Pitch Pine	*Pinus rigida*	S2S3	-	-	• typically a Non-Wetland species
Rough Sand Sedge	*Cyperus schweinitzii*	S3	-	-	• Non-Wetland Species • Upland species • found in dunes and other dry, open sandy sites
Round-leaf Ragwort	*Senecio obovatus*	S3	-	-	• Non-Wetland species • Upland/Alvar species
Rue Anemone	*Senecio obovatus*	S3	-	-	• marginal Wetland Species • floodplain, mesic woods
Sand Reedgrass	*Calamovilfa longifolia*	S3	-	-	• not usually a Wetland species • Dune species
Short's Aster	*Aster shortii*	S2	NAR	NIAC	• Non-Wetland species • dryish upland woods habitat
Shrubby St. John's-wort	*Hypericum prolificum*	S2	-	-	• Non-Wetland species • Upland species
Stemless Rubberweed	*Hymenoxys herbacea*	S2	-	-	• Non-Wetland species
Stiff-leaved Goldenrod	*Solidago rigida*	S3	-	-	• Non-Wetland species • dry prairie species
Tall Joe-pyeweed	*Eupatorium altissimum*	S1	-	-	• Non-Wetland species • Upland species
Vasey's Rush	*Juncus vaseyi*	S3	-	-	• Wetland species, but probably a misidentification
Virginia Culver's-root	*Veronicastrum virginicum*	S2	-	-	• typically not a wetland species • found in moist prairie
Wood Fern	*Dryopteris X triploidea*	S3S4	-	-	• Non-Wetland species • Hybrid species

Appendix V: Scientific Names of Common (Non-Significant) Species mentioned in the Atlas

Common Name	Scientific Name
Vascular Plants	
American Lotus	*Nelumbo lutea*
Curled Pondweed	*Potamogeton crispus*
Eurasian Water Milfoil	*Myriophyllum sibiricum*
Purple Loosestrife	*Lythrum salicaria*
Reed Canary Grass	*Phalaris arundinacea*
Sago Pondweed	*Potamogeton pectinatus*
White Water Lily	*Nymphaea odorata*
Wild Celery	*Vallisneria americana*
Yellow Water lily	*Nuphar variegatum*
Fish	
Alewife	*Alosa pseudoharengus*
Banded Killfish	*Fundulus diaphanus*
Black Crappie	*Pomoxis nigromaculatus*
Blacknose Shiner	*Notropis hetrolepis*
Bluegill	*Lepomis macrochirus*
Bluntnose Minnow	*Pimephales notatus*
Bowfin	*Amia calva*
Brook Silverside	*Labidesthes sicculus*
Brown Bullhead	*Ictalurus nebulosis*
Channel Catfish	*Ictalurus punctatus*
Common Carp	*Cyprinus carpio*
Emerald Shiner	*Notropis antherinoides*
Freshwater Drum	*Aplodinotus grunniens*
Gizzard Shad	*Dorosoma cepedianum*
Golden Shiner	*Notemigonus crysoleucas*
Goldfish	*Carassius auratus*
Johnny Darter	*Etheostoma nigrum*
Lake Sturgeon	*Acipenser fulvescens*
Largemouth Bass	*Micropterus salmoides*
Log Perch	*Percina caprodes*
Muskellunge	*Esox masquinongy*
Northern Pike	*Esox lucius*
Pumpkinseed	*Lepomis gibbosus*
Rainbow Smelt	*Osmerus mordax*
Rock Bass	*Ambloplites rupestris*
Smallmouth Bass	*Micropterus dolomieui*
Spottail Shiner	*Notropis hudsonius*
Walleye	*Stizostedion vitreum*
White Bass	*Morone chrysops*
White Crappie	*Pomoxis annularis*
White Perch	*Morone americana*
White Sucker	*Catostomus commersoni*
Yellow Perch	*Perca flavescens*
Amphibians and Reptiles	
Bullfrog	*Rana catesbeiana*
Eastern Garter Snake	*Thamnophis sirtalis sirtalis*
Eastern Milk Snake	*Lampropeltis triangulum*
Green Frog	*Rana clamitans*
Grey Tree Frog	*Hyla versicolor*
Midland Painted Turtle	*Chrysemys picta marginata*
Mudpuppy	*Necturus maculosus*
Northern Dusky Salamander	*Desmognathus fuscus*
Northern Leopard Frog	*Rana pipiens*
Northern Water Snake	*Natrix sipedon sipedon*
Red-spotted Newt	*Notophthalmus viridescens*
Snapping Turtle	*Chelydra serpentina*
Spring Peeper	*Pseudacris crucifer*
Birds	
American Bittern	Botaurus lentiginosus
American Coot	*Fulica americana*
American Widgeon	*Anas americana*
Black Duck	*Anas rubripes*
Blue-winged Teal	*Anas discors*
Bufflehead	*Bucephala albeola*
Canada Goose	*Branta canadensis*
Canvasback	*Aythya valisineria*
Caspian Tern	*Sterna caspia*
Common Goldeneye	*Bucephala clangula*
Common Merganser	*Mergus merganser*
Common Tern	*Sterna hirundo*
Cooper's Hawk	*Accipiter cooperii*
Double-crested Cormorant	*Phalacrocorax auritus*
Great Blue Heron	*Ardea herodias*
Green Heron	*Butorides virescens*
Green-winged Teal	*Anas crecca*
Herring Gull	*Larus argentatus*
Mallard	*Anas platyrhynchos*
Northern Harrier	*Circus cyaneus*
Red-breasted Merganser	*Mergus serrator*
Red-tailed Hawk	*Buteo jamaicensis*
Red-winged Blackbird	*Agelaius phoniceus*
Sharp-shinned Hawk	*Accipiter striatus*
Tree Swallow	*Tachycineta bicolor*
Wood Duck	*Aix sponsa*
Invertebrates	
Zebra Mussel	*Dreissena polymorpha*
Mammals	
Beaver	*Castor canadensis*
Coyote	*Canis latrans*
Eastern Cottontail Rabbit	*Sylvilagus floridanus*
Mink	*Mustela vison*
Moose	*Alces alces*
Muskrat	*Ondatra zibethicus*
Raccoon	*Procyon lotor*
Red Fox	*Vulpes vulpes*
River Otter	*Lontra canadensis*
Short-tailed Weasel	*Mustela erminea*
Striped Skunk	*Mephitis mephitis*
Virginia Opossum	*Didelphis virginiana*
White-tailed Deer	*Odocoileus virginianus*

Lake Superior Coastal Wetlands



FORT FRANCIS DISTRICT
RAINY RIVER
THUNDER BAY DISTRICT
THUNDER BAY
NIPIGON DISTRICT
WAWA DISTRICT
ALGOMA
CHAPLEAU DISTRICT
SUDBURY
SAULT STE. MARIE DISTRICT
Lake Superior
Canada
U.S.A.
See Inset 1
Inset 1
1
2
3
4
5
6
7
8
9
10
11
A1
A2
A3
A4
A6
A7
N1
N2
N3
N4
N5
N6
LEGEND
Swamp
Marsh
Bog
Fen
Evaluated Wetland - Provincially Significant
Evaluated Wetland - Other
NHIC Natural Area Designation - Unevaluated Wetland Component
Environmental Sensitivity Area (ESA) Unevaluated Wetland
Other Unevaluated Wetland
Wetland overlaps with other Natural Area(s) in NHIC Natural Areas database
Coastline
Administrative OMNR Districts
Scale 1: 2 382 000
20
0
20
40
Kilometers
N
Natural Heritage Information Centre
Ministry of Natural Resources
Ontario
Environment Canada
© February, Queen's Printer for Ontario


Lake Huron and St. Marys River Coastal Wetlands

Detroit River, Lake St. Clair and St. Clair River Coastal Wetlands


14
13
St. Clair River
LAMBTON
12
11
10
AYLMER DISTRICT
A1
U.S.A.
Canada
Lake St. Clair
9
KENT
8
7
6
5
Detroit River
3
4
2
ESSEX
1
Lake Erie
LEGEND
Swamp
Marsh
Bog
Fen
Evaluated Wetland - Provincially Significant
Evaluated Wetland - Other
NHIC Natural Area Designation - Unevaluated Wetland Component
Environmentally Sensitive Area (ESA) Unevaluated Wetland
Other Unevaluated Wetland
Wetland overlaps with other Natural Area(s) in NHIC Natural Areas database
Coastline
Administrative (MNR) Districts
Scale 1: 1 013 000
0
5
10 Kilometres
Natural Heritage Information Centre
Ministry of Natural Resources
Ministère des Richesses naturelles
Ontario
© April, 1999 Queen's Printer for Ontario

Lake Erie Coastal Wetlands


Lake Huron
Lake Ontario
Lake Erie
CAMBRIDGE DISTRICT
NIAGARA
HALDIMAND-NORFOLK
AYLMER DISTRICT
ELGIN
KENT
ESSEX
Canada
U.S.A.
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
N1
N2
N3
LEGEND
Swamp
Marsh
Bog
Fen
Evaluated Wetland - Provincially Significant
Evaluated Wetland - Other
NHIC Natural Area Designation - Unevaluated Wetland Component
Environmental Sensitivity Area (ESA) Unevaluated Wetland
Other Unevaluated Wetland
Wetland overlaps with other Natural Area(s) in NHIC Natural Areas database
Counties
Administrative OMNR District
10
0
10
20
Kilometers
Scale 1: 1 300 000
Natural Heritage Information Centre
Ministry of Natural Resources
N
© April, 1999 Queen's Printer for Ontario

Lake Ontario and Niagara River Coastal Wetlands


LENNOX & ADDINGTON
HASTINGS
PETERBOROUGH DISTRICT
NORTHUMBERLAND
PRINCE EDWARD
See Inset 1
DURHAM
AURORA DISTRICT
METROPOLITAN TORONTO
PEEL
HALTON
HAMILTON-WENTWORTH
CAMBRIDGE DISTRICT
NIAGARA
Niagara River
Lake Ontario
Canada
U.S.A.
Inset 1
5 0 5 Kilometres
LEGEND
Swamp
Marsh
Bog
Fen
Evaluated Wetland - Provincially Significant
Evaluated Wetland - Other
NHIC Natural Area Designation - Unevaluated Wetland Component
Environmental Sensitivity Area (ESA) Unevaluated Wetland
Other Unevaluated Wetland
Wetland overlaps with other Natural Area(s) in NHIC Natural Areas database
Counties
Administrative OMNR Districts
Scale 1: 815 000
10 0 10 Kilometres
N
© April, 1998 Queen's Printer for Ontario

St. Lawrence River Coastal Wetlands



The Ontario Great Lakes Coastal Wetlands Atlas contributes to delivering the goals of the following initiatives.

Great Lakes Wetlands Conservation Action Plan (GLWCAP)

The GLWCAP is a cooperative program of the federal and Ontario governments as well as non-governmental groups to protect, rehabilitate and create wetlands on the Canadian side of the Great Lakes basin. Implementation of the plan is coordinated by representatives from the Canadian Wildlife Service of Environment Canada, the Ontario Ministry of Natural Resources, The Nature Conservancy of Canada, the Federation of Ontario Naturalists, and Ducks Unlimited Canada. The Atlas contributes to several GLWCAP strategies, including: improve wetland science, data and monitoring, and improve coordination. For more information, visit www.on.ec.gc.ca/wetlands/glwcap-e.cfm.

Eastern Habitat Joint Venture (EHJV)

The EHJV is one of 14 joint ventures established across the continent to ensure the implementation of the North American Waterfowl Management Plan. In Ontario, the EHJV is a partnership of the federal and Ontario governments, Ducks Unlimited Canada, The Nature Conservancy of Canada, and Wildlife Habitat Canada. Extensive programs are applied on a broad scale to influence land use policies and promote ecologically sound and sustainable land use practices. Intensive programs are tailored to secure, create, restore, or rehabilitate balanced habitat conditions for waterfowl and other wetland wildlife. For more information, visit www.nawmp.ca.

Species at Risk

The Government of Canada's three-part strategy to protect species at risk consists of building on the federal-provincial-territorial Accord for the Protection of Species at Risk, developing and implementing stewardship and incentive programs, and creating and passing federal endangered species legislation, the *Species at Risk Act* (SARA). For more information, visit www.speciesatrisk.gc.ca.

Land Information Ontario (LIO)

LIO is an initiative to restructure the management and use of Ontario's land information infrastructure led by the Ontario Ministry of Natural Resources (OMNR). LIO is building a new way to manage land information; one which will ensure that all of Ontario's land information assets are standardized, properly managed, widely known, accessible, and collected efficiently. At present, natural resource, environmental, and geologic themes dominate the approximately 140 layers of data contained in the warehouse. This composition is expected to change as more of LIO's stakeholders migrate their holdings to the facility. LIO provides the public and private sectors with an unprecedented opportunity to work together in the collection, management and distribution of Ontario's land information assets. For more information, visit www.lio.mnr.gov.on.ca.

The Canada-Ontario Agreement Respecting the Great Lakes Basin (COA)

The COA is a commitment by the governments of Canada and Ontario to restore and protect the Great Lakes Basin Ecosystem. Renewed in 2002, the current Agreement outlines how the two governments will cooperate and coordinate their efforts to achieve this commitment. The Agreement includes an overarching agreement to stand in perpetuity, along with four annexes, each of which specifies five-year goals for the Great Lakes Basin Ecosystem. The Lakewide Management Annex for example, includes commitments for Canada and Ontario to work towards achieving: rehabilitated, conserved, and protected fish and wildlife habitats and protected areas; and coordinated and integrated monitoring for scientific interpretative reporting, decision-making, and reporting on progress. For more information, visit www.on.ec.gc.ca/coa.

The State of the Lakes Ecosystem Conference (SOLEC)

The SOLEC conferences are hosted by the U. S. Environmental Protection Agency and Environment Canada on behalf of the two countries every two years in response to the binational Great Lakes Water Quality Agreement. The conferences are intended to provide a forum for exchange of information on the ecological condition of the Great Lakes and surrounding lands. A major purpose of this is to reach a large audience of people in the government (at all levels), corporate and not-for-profit sectors who make decisions that affect the lakes. For more information, visit www.on.ec.gc.ca/solec.